

Available online on 15.01.2017 at <http://jddtonline.info>

## Journal of Drug Delivery and Therapeutics

Open access to Pharmaceutical and Medical research

© 2016, publisher and licensee JDDT, This is an Open Access article which permits unrestricted noncommercial use, provided the original work is properly cited



### Research Article

## PHARMACOGNOSTICAL EVALUATION AND PHYTOCHEMICAL SCREENING OF THEVETIA PERUVIANA

Jitpure Rupesh\*, Yadav Savita, Kurmi Ronak, Dhaker Kehar, Bhowmick Mithun, Rathi Jagdish

NRI Institute of Pharmaceutical Sciences, Bhopal, Madhya Pradesh

### ABSTRACT

Flowers of Plant *Thevetia peruviana* (Pers.) K. Schum. were reported to possess good medicinal value in traditional system of medicine, the present investigation deals with Macroscopic, Microscopic and preliminary phytochemical investigation of flowers of *Thevetia peruviana* which includes pharmacognostical parameters, physicochemical parameters like ash values, extractive values and moisture content. The total ash, acid insoluble ash, water-soluble ash values and sulfated ash were observed to be 3.50%, 1.60%, 1.30% and 1.20% w/w respectively. Alcohol soluble, water-soluble and Ether soluble extractive values of the flowers were observed to be 8.30%, 4.80%, and 3.70% respectively. Powdered flowers were also subjected to fluorescence analysis with different chemicals. Phytochemical investigation of powdered drug revealed the presence of Alkaloids, glycosides, tannins, phenolic compounds, proteins, essential oils, gums, mucilage and fixed oils. The main aim of the present investigation is to study the macro, microscopic and some other pharmacognostic characters and physicochemical standards of flowers of *Thevetia peruviana* Pers. which could be used to prepare a monograph for the proper identification of the plant.

**Keywords:** *Thevetia peruviana*, Phytochemical, Fluorescence analysis, Physical evaluation.

### Article Info

#### Cite this article as:

Jitpure R, Yadav S, Kurmi R, Dhaker K, Bhowmick M, Rathi J, Pharmacognostical evaluation and phytochemical screening of *Thevetia peruviana*, Journal of Drug Delivery and Therapeutics. 2017; 7(1):60-64, DOI: <http://dx.doi.org/10.22270/jddt.v7i1.1378>

#### \*Address for Correspondence

Mr. Rupesh jitpure, M.S. Savita yadav, B.Pharm. Student, Department of Pharmacognosy, NRI Institute of Pharmaceutical Sciences, R.G.P.V. University, Bhopal, M.P., India-462023 Mob:- 08878830679, 8717844939, Email:- saviy262@gmail.com

## INTRODUCTION:

*Thevetia peruviana* (Pers.) K. Schum. is a small evergreen plant of 2–6 meter in height with a broad geographical and ecological distribution<sup>1</sup>, and its convinced parts are used as medicinal resources in Chinese folk remedy<sup>2</sup>. *Thevetia peruviana* is drought-tolerant evergreen plant of the Family *Apocynaceae* that originate from Mediterranean countries<sup>3</sup>. All parts of the yellow oleander plant are toxic to humans, animals and certain insects<sup>4</sup>.

This species generate derivative metabolites<sup>5</sup>, some of which are of pharmacological significance. It is usually grown-up as an ornamental plant in warm clement and subtropical regions, due to its profuse and long durable flowering and temperate durability<sup>6-7</sup>. It is used for screens, equivocation alongside highways, planting along beaches and in town areas by removing suckers and leaving just a few stems, it can also be shaped into very attractive small trees. In Northern regions it may be grown-up as an interior or terrace plant. Oleander

has supple branches with green, flat bark ultimately turning to dark grey.

On cutting or broken branches emanate a thick, white sap<sup>8-9</sup>. The leaves are 5 to 15 cm long, thin, acuminate or sharp in the apex, shortly petiolate, with a coriaceous dark-green cutting edge. Flowers are shaped in fatal heads and their colors differ from deep yellow, white, purple and orange<sup>10</sup>. Each flower is about 5 cm in diameter with five petals. The fruit consists of a thin follicle 7.5 to 18.5 cm long which opens to separate feathery seeds. Oleander can be propagated by seed<sup>11</sup> but, being allogamous and highly heterozygous, it shows huge unpredictability in seedling populations. Phytochemical Analysis revealed that *Thevetia peruviana* contains a new cardiac glycoside called Digitoxigenin<sup>12</sup>, Thevetin A and B, theveridoside, cerberin<sup>12,13</sup>, galactouronic acid, rhamnase, aucubin, ursolic acid, cardenolides<sup>13</sup>, quercetin, alpha and beeta amyryn, and lupenyl acetate<sup>14</sup>, as prime phytoconstituents.

## MATERIAL AND METHODS:

### Plant material collection and Authentication:

The flowers of plant *Thevetia peruviana* Pers. were collected from the local area Jaipur Dist. Rajasthan, India, in the month of June-July 2010 and were positively identified and confirmed by the botanist, Mr. A. Sharma, Department of Botany, University of Rajasthan, Jaipur. A voucher specimen has been deposited (RUBL 20856) in the herbarium of the botany department of the University of Rajasthan. The fresh mature flowers were used for the study of macroscopic and microscopic characters, whereas the dried uniform flower powder was used for the determination of ash value, extractive Values, loss on drying and phytochemical investigation.

### Drying and Pulverization:

Flowers of *Thevetia peruviana* were collected and cut into small pieces. It was shed dried and pulverized to mash size 22 and stored in air tight container for further use.

### PHARMACOGNOSTIC STUDY Macroscopic studies:

Whole fresh flowers of *Thevetia peruviana* was examined for color, odor, taste, shape and structure, size and touch were determined.

### Microscopic studies:

Under microscopical examination histological characters of flower were studied, different sections of

flower made and observed under Compound microscope. For the microscopical study, all flowers were collected and washed with distilled water properly and cut thin sections of flower and also different parts like Petals, Sepals, ovary etc. with the help of blade and swept the sections with the fine camel hair brush and suspended in distilled water in watch glass. After that selected thin section placed on cleaned glass slide and added one drop of glycerin and covered with cover slip carefully, wiped out excess of mountant on slide with filter paper and observed under compound microscope for various histological characters of flower parts.

Powder of the dried flowers was used for the observation of powder microscopical characters. The powdered drug was separately treated with phloroglucinol and HCl solution. Glycerin and iodine solution were used to determine organoleptic properties and the presence of Basal cells of trichomes, epidermal cells, oil globules, vein termination, fibers and starch grains<sup>15,16</sup>.

### Physiochemical Investigation:

The moisture content, total ash, water-soluble ash, acid-insoluble ash, sulphated ash, alcohol, water-soluble and ether soluble extractive values were determined as a part of its physiochemical parameters. The Powdered flowers parts were subjected to analysis under day/visible light and ultra violet light after treatment with various chemical as a part of Fluorescence analysis<sup>15</sup>.

## RESULTS:




Figure 1

Figure 2

Figure 3

Figure 1, 2, 3: Morphological Features of Flower, Petal, Sepal

Table 1: Morphological properties of flowers of *Thevetia peruviana*

S.N.	Observed For	Fresh Flower
1.	Shape & Structure	Narrow funnel shaped with swird Petals,
2.	Color	Yellowish green, Dark yellow, Yellowish
3.	Taste	Sweetish, agreeable
4.	Odour	Characteristic, Pleasant smell
5.	Touch	Smooth, Soft
6.	Size	6-7 × 2-3 cm

In the microscopic studies, the flower shows all the typical characteristics of flower (Fig. 4). The petal shows the presence of upper and lower epidermis, centrally vascular bundles as phloem surrounds with the xylem, cortex, parenchyma and also trichomes (Fig. 5). The sepal shows the presence of upper and lower epidermis, centrally vascular bundles as phloem surrounds with the xylem, cortex (Fig. 6). The cross section of ovary shows the presence of trichomes, vein trace, locule, ovule and placenta. (Fig. 7)

Microscopic study of powder flower revealed the presence of basal Cell of the Trichomes, epidermal cell, stomata, vein – islets, vein – termination. (Fig. 8)

Physiochemical tests of powdered drug of *Thevetia peruviana* (Pers.) K. Schum. Flower for the presence of secondary metabolites and color observation shows the following results when treated with different reagents. (Table 2)




Figure 4: L. S. of Flower      Figure 5: T. S. of Petal      Figure 6: T. S. of Sepal      Figure 7: T. S. of Ovary  
IE- inner epidermis, LE- lower epidermis, PH- phloem, PC- parenchyma, CL- cuticle, XY- xylem, UP- upper epidermis, LPTV- leaf parallel trace vein, C- cortex, TR- trichomes, VT- vein trace, LE- locule, OE- ovule, PA- placenta.




Figure 8, 9: Powder Microscopy of flower  
BTr – Basal cell of the Trichomes, Ec – Epidermal cells, VT – Vein termination, OG – oil globules, SG – starch grains, F – fiber

Table 2: Physiochemical analysis of powdered drug of flower of *Thevetia peruviana*

S.N.	REAGENTS	COLOUR OBSERVED
1.	Powder as such	Light black to Yellowish
2.	Powder + Conc. HCl	Brownish black
3.	Powder + Conc. HNO <sub>3</sub>	Light brownish
4.	Powder + Conc. H <sub>2</sub> SO <sub>4</sub>	Light brick red
5.	Powder + Glacial acetic acid	Light brownish
6.	Powder + 5% NaOH	Light brownish
7.	Powder + 5% KOH	Light brownish
8.	Powder + 5% Ferric chloride	Yellow-brownish
9.	Powder + Picric acid (Aq. Sol.)	Yellowish
10.	Powder + Ammonia solution	Light brownish

The physical constant evaluation of the drugs is an important parameter in detecting adulteration or improper handling of drugs. The total ash is particularly important in the evaluation of purity of drugs, i.e. the

presence or absence of foreign inorganic matter such as metallic salts and/or silica. The ash values, extractive values and moisture content of flowers were determined. The results are depicted in (Table 3).

**Table 3: Physical parameters of flowers of *Thevetia peruviana***

S. No.	PARAMETER	VALUES (%)w/w
1.	Loss on Drying	9.5
2.	Extractive Values	
	Ethanol soluble extractive	8.3
	Water soluble extractive	4.8
	Petroleum ether soluble Extractive	3.7
3.	Ash Values	
	Total Ash	3.5
	Water soluble ash	1.3
	Acid insoluble ash	1.6
	Sulphated ash	1.2

**Table 4: Fluorescence analysis of powder of flowers of *Thevetia peruviana***

S.NO.	TREATMENT	OBSERVATION (under long wavelength)	OBSERVATION (under short wavelength)
1.	Powder as such	No fluorescence	Yellowish black
2.	Powder + 1 N NaOH in methanol	No fluorescence	Greenish black
3.	Powder + 1 N NaOH in water	No fluorescence	Yellowish black
4.	Powder + 50% HCl	No fluorescence	Light yellowish
5.	Powder + 50% HNO <sub>3</sub>	Reddish black	Yellowish
6.	Powder + 50% H <sub>2</sub> SO <sub>4</sub>	No fluorescence	Dark brownish
7.	Powder + Petroleum ether	Light brown	Light yellowish
8.	Powder + Chloroform	No fluorescence	Light yellowish
9.	Powder + Picric acid	Light yellowish	Yellowish
10.	Powder + 5% Ferric chloride solution	No fluorescence	Greenish-yellowish
11.	Powder + 5% iodine solution	No fluorescence	Light brown yellowish
	Powder + Methanol	No fluorescence	Light brown
13.	Powder + HNO <sub>3</sub> + NH <sub>3</sub>	No fluorescence	Yellowish black

**Table 5: Preliminary Phytochemica screening of powder of *Thevetia peruviana* flowers**

S.NO.	TEST FOR	TESTS	POWDERED DRUG
1.	CARBOHYDRATES	Molisch's test	- VE
		Iodine test	-VE
		Fehling's test	+VE
		Barfoed's test	-VE
2.	PROTEIN & AMINO ACIDS	Ninhydrin test	+VE
		Biuret test	+VE
		Millon's test	+VE
		Test with tannic acid	+VE
3.	OILS & FATS	Spot test	+VE
		Saponification test	-VE
4.	ALKALOIDS	Dragendorff's test	+VE
		Mayer's reagent test	+VE
		Wagner's reagent test	+VE
		Hager's reagent test	+VE
5.	GLYCOSIDES	Legal's test	+VE
		Borntrager's test	+VE
		Keller-killiani's test	+VE
6.	FLAVONOIDS	Ferric chloride test	+VE
		Shinoda's test	-VE
		Fluorescence test	-VE
7.	VOLATILE OIL	By Clevenger apparatus	+VE

The result of fluorescence studies of flowers powder using different reagents is given in **Table 4**. Fluorescence is an important phenomenon exhibited by various chemical constituents present in plant material. Some constituents show fluorescence in the visible range in day light. The ultra violet light produces fluorescence in many natural products (e.g. alkaloids like berberine), which do not visibly fluoresce in day light. If the substances themselves are not fluorescent, they may often be converted into fluorescent derivatives by applying different reagents hence some crude drugs are often assessed qualitatively in this way and it is an important parameter of pharmacognostical evaluation.

Preliminary phytochemical screening of powdered drug of flowers indicated high concentration of glycosides, alkaloids along with other constituents like flavonoids, essential oils and steroids. These secondary metabolites are known to possess various pharmacological effects and may be responsible for various pharmacological effects of *Thevetia peruviana* flowers. (Table 5)

### ACKNOWLEDGEMENTS:

Authors are heartily thankful to Principal Dr. Jagdish Rathi and Assit. prof. Mr. kehar dhaker for providing necessary facilities at NRI Institute of Pharmaceutical Sciences, R.G.P.V. University, Bhopal, Madhya Pradesh, India.

### REFERENCES:

- Shannon D. Langford & Paul J. Boor "Oleander toxicity: an examination of human and animal toxic exposures". *Toxicology*, 1996; 109(1):1–13.
- Bose TK, Basu RK, Biswas B, De JN, Majumdar BC, Datta S. "Cardiovascular effects of yellow oleander ingestion". *J Indian Med Assoc*, 1996; 97(10):407–410.
- Kohls S, Scholz-Böttcher BM, Teske J, Zark P, Rullkötter J. "Cardiac glycosides from Yellow Oleander (*Thevetia peruviana*) seeds". *Phytochemistry*, 2012; 75:114–27.
- Rajapakse S. "Management of yellow oleander poisoning". *Clinical Toxicology*, 2009; 47:12-18.
- Roberts D.M.; Southcott E.; Potter J.M.; Roberts M.S.; Eddleston M.; Buckley N. "Pharmacokinetics of digoxin cross-reacting substances in patients with acute yellow oleander (*Thevetia peruviana*) poisoning, including the effect of activated charcoal". *Therapeutic Drug Monitoring*, 2009; 28(6):784–792.
- Bandara V.; Weinstein S.A.; White J.; Eddleston M. "A review of the natural history, toxinology, diagnosis and clinical management of Nerium oleander (common oleander) and *Thevetia peruviana* (yellow oleander) poisoning". *Toxicon*, 2010; 56(3):273–281.
- M. Eddleston; S. Rajapakse; Rajakanthan, S. Jayalath; L. Sjöström; W. Santharaj; et al.). "Anti-digoxin Fab fragments in cardiotoxicity induced by ingestion of yellow oleander: a randomised controlled trial". *Lancet*, 2000; 355:967–972.
- Kannan R. "Koels feeding on the yellow oleander". *Blackbuck*, 1991; 7(2): 48.
- Krishnan M. "Koels (*Eudynamis scolopacea*) eating the poisonous fruit of the Yellow Oleander." *J. Bombay Nat. Hist. Soc.*, 1952; 50(4):943–945.
- Raj, P J Sanjeeva . "Additions to the list of birds eating the fruit of Yellow Oleander (*Thevetia neriifolia*).". *J. Bombay Nat. Hist. Soc.*, 1963; 60(2):457–458.
- Raj, P J Sanjeeva . "Birds eating poisonous fruit of Yellow Oleander (*Thevetia neriifolia*).". *J. Bombay Nat. Hist. Soc.*, 1959; 56 (3): 639.
- Neelakantan, KK . "Common Grey Hornbill (*Tockus birostris*) eating fruits of the Yellow Oleander (*Thevetia neriifolia*).". *J. Bombay Nat. Hist. Soc.*, 1953; 51(3):738.
- Rajasingh, Simon G; Rajasingh, Irene V . "Birds and mammals eating the fruits of Yellow Oleander (*Thevetia peruviana*).". *J. Bombay Nat. Hist. Soc.*, 1970; 67(3): 572–573.
- Gupta R, Kachhawa JB, Gupta RS, Sharma AK, Sharma MC, Dobhal MP . "Phytochemical evaluation and antispermatogenic activity of *Thevetia peruviana* methanol extract in male albino rats". *Hum Fertil (Camb)*, 2011; 14(1):53–59. [Doi:10.3109/14647273.2010.542230](https://doi.org/10.3109/14647273.2010.542230).
- Kareru P.G.; Keriko J.M.; Kenji G.M.; Gachanja A.N. "Anti-termite and antimicrobial properties of paint made from *Thevetia peruviana* (Pers.) Schum. Oil extract". *African Journal of Pharmacy and Pharmacology*, 2010; 4(2):87–89.
- Islam Mazhar-Ul, Iqbal A, Usmanghani K, Shahab-Ud-Din, Ahmed A, *Pakistan Journal of Pharmacology*, 2001, 1-9.
- Gehlot NK, Sharma VN, Vyas DS, *Indian Journal of Pharmacology*, 1976; 8:92-949.