

Available online on 15.09.2018 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Antibacterial, antifungal and antioxidant activity of *Dichapetalum gelonioides* (Roxb.) Engl. (Dichapetalaceae)

Priyanka G.S, Nitish Bharadwaj A, Sachin M.B, Akhilesha S, Prashith Kekuda T.R*

Department of Microbiology, S.R.N.M.N College of Applied Sciences, N.E.S Campus, Balraj Urs road, Shivamogga-577201, Karnataka, India

ABSTRACT

Objectives: *Dichapetalum gelonioides* (Roxb.) Engl. belongs to the family Dichapetalaceae. In the present study, we investigated antibacterial, antifungal and antioxidant activity of methanolic extract of leaf and fruit of *D. gelonioides*.

Methods: Maceration process was carried out for extraction of leaf and fruit of *D. gelonioides*. Agar well diffusion method was employed to evaluate antibacterial activity of extracts against gram positive and gram negative bacteria. Poisoned food technique was performed to investigate antifungal activity of extracts against two seed-borne fungi. Antioxidant activity was evaluated by DPPH radical scavenging and ferric reducing assays.

Results: Both leaf and fruit extracts were effective in causing inhibition of all test bacteria. Highest and least inhibitory activity was observed against *Bacillus cereus* and *Escherichia coli* respectively. Both *Aspergillus niger* and *Bipolaris* sp. were inhibited to >50% by leaf and fruit extracts. Extent of inhibition of *Bipolaris* sp. was slightly higher when compared to *A. niger*. Both leaf and fruit extracts showed a dose dependent scavenging of DPPH radicals with high activity being showed by leaf extract. Leaf extract was shown to exhibit marked reducing potential than fruit extract.

Conclusions: Overall, leaf extract was shown to be more effective in displaying antioxidant activity and causing inhibition of bacteria and fungi when compared to fruit extract. The results indicate that the plant possess active principles which are to be purified, characterized and subjected for antimicrobial and antioxidant assays in further studies.

Keywords: *Dichapetalum gelonioides*, Maceration, Agar well diffusion, Poisoned food technique, DPPH, Ferric reducing

Article Info: Received 29 June, 2018; Review Completed 11 Aug 2018; Accepted 1 8 Aug 2018; Available online 15 Sep 2018



Cite this article as:

Priyanka GS, Bharadwaj NA, Sachin MB, Akhilesha S, Prashith Kekuda TR, Antibacterial, antifungal and antioxidant activity of *Dichapetalum gelonioides* (Roxb.) Engl. (Dichapetalaceae), Journal of Drug Delivery and Therapeutics. 2018; 8(5):263-267 DOI: <http://dx.doi.org/10.22270/jddt.v8i5.1836>

*Address for Correspondence:

Dr. Prashith Kekuda T.R, Department of Microbiology, S.R.N.M.N College of Applied Sciences, N.E.S Campus, Balraj Urs road, Shivamogga-577201, Karnataka, India

INTRODUCTION

Plants are widely used as indispensable sources of food, medicine, dyes, flavoring agents and timber worldwide. Plants have been considered as an integral part of ethnomedicine and people from several parts of the world rely on plant based medicines (in the form of single or polyherbal formulations, decoctions, paste, infusions and others) for primary healthcare needs. Plants are key ingredients in some medicinal systems such as Ayurveda, Traditional Chinese Medicine, Siddha and Unani.¹⁻⁷ The genus *Dichapetalum*

belonging to the family Dichapetalaceae includes shrubs or small trees. Species of *Dichapetalum* are rich source of triterpenoids. *Dichapetalum* species are used traditionally as medicine to treat certain illnesses.⁸⁻¹⁰ Bioactive dichapetalins and several other compounds have been isolated from *Dichapetalum* species.¹¹⁻¹⁶

Dichapetalum gelonioides (Roxb.) Engl (Figure 1) is a large shrub or small tree with pubescent branchlets and found distributed in evergreen and semi-evergreen forests as forest undergrowth. Leaves are elliptic or elliptic-lanceolate, 15x5cm, acuminate or caudate at

apex and glabrous. Flowering occurs more or less throughout year. Flowers are small, polygamo-monoecious, found in short-peduncled axillary clusters. Fruit is a 2-celled drupe, 1-2 seeded, densely tomentose, stone hard and enclosed in a scarlet mesocarp.¹⁷ The plant is known to be a hyperaccumulator of nickel.¹⁸ The fruits and leaves of *D. gelonioides* is used traditionally to treat amenorrhea, mouth ulcers, swelling, pit viper biting, polydipsia and wounds.¹⁰ Isolated constituents, in particular dichapetalins, exhibit some biological activities.^{11,14} The bark of the plant is reported to exhibit weaker muscarinic receptor activity.¹⁹ The present study was conducted to evaluate antibacterial, antifungal and antioxidant activity of leaf and fruit of *D. gelonioides*.



Figure 1: *D. gelonioides* (Photograph by Prashith Kekuda)

MATERIALS AND METHODS

Collection and extraction of plant material

Plant was collected during December 2017 at Haniya, Hosanagara Taluk, Shivamogga district, Karnataka. The plant was identified by Dr. Vinayaka K.S, Principal, KFGC, Shikaripura, Karnataka. Leaves and fruits were separated, washed to remove adhering matter, dried under shade and powdered mechanically. The powdered leaf and fruit materials were extracted separately by maceration process using methanol as solvent.⁶ The crude leaf and fruit extracts were screened for phytochemicals by standard tests.^{20,21}

Antibacterial activity of leaf and fruit extracts

Agar well diffusion method was used to investigate antibacterial activity of leaf and fruit extracts of *D. gelonioides* (20mg/ml of DMSO) against 24 hours old broth cultures of test bacteria viz. *Bacillus cereus*, *Escherichia coli*, *Shigella flexneri* and *Salmonella typhimurium*. Streptomycin (1mg/ml of sterile distilled water) was used as standard antibiotic. Diameter of zones of inhibition formed around wells was measured.⁶

Antifungal activity of leaf and fruit extracts

Poisoned food technique was employed to evaluate antifungal potential of leaf and fruit extracts of *D. gelonioides* against two seed-borne fungi viz. *Aspergillus niger* and *Bipolaris* sp. Antifungal activity of leaf and fruit extracts, in terms of inhibition of mycelial growth of test fungi (%), was determined using the formula:

Inhibition of fungal growth (%) = $[(D_c - D_t) / D_c] \times 100$, where 'D_c' and 'D_t' denotes the diameter of fungal colonies in control (without extract) and poisoned plates (1mg extract/ml of medium) respectively.⁶

DPPH free radical scavenging activity of leaf and fruit extracts

Various concentrations of ascorbic acid (reference antioxidant) leaf and fruit extracts of *D. gelonioides* (12.5-200µg/ml) were screened for antiradical potential by performing DPPH radical scavenging assay.⁶ The absorbance was measured spectrophotometrically at 517nm. Radical scavenging activity (%) was determined using the formula:

Radical scavenging activity (%) = $(A_c - A_t) / A_c \times 100$, where 'A_c' and 'A_t' represents absorbance of DPPH control and absorbance of DPPH in the presence of extract/ascorbic acid.

Ferric reducing activity of leaf and fruit extracts

Various concentrations of ascorbic acid (reference standard) and leaf and fruit extracts of *D. gelonioides* (12.5-200µg/ml) were screened for ferric reducing activity. The absorbance of reaction mixture in each tube was measured spectrophotometrically at 700nm. An increase in the absorbance with increase in concentration indicates ferric reducing potential.⁶

Statistical analysis

Antimicrobial studies were carried out in triplicates (n=3). Results are presented as Mean±Standard deviation (S.D).

RESULTS AND DISCUSSION

Plants produce a number of secondary metabolites by means of metabolic pathways such as mevalonic acid pathway, malonic acid pathway and shikimic acid pathway. These metabolites are distributed in various parts of the plant such as leaf, root, seed and flower. Secondary metabolites protect the plants from herbivores and pathogens and contribute to flavor and aroma of plants. Many of plant secondary metabolites exert definite physiological roles in human body and exhibit a range of biological activities including anticancer, antimicrobial and antioxidant activities.²²⁻²⁶ Saponins, tannins, alkaloids, flavonoids, triterpenoids and glycosides were detected in leaf extract of *D. gelonioides*. Sterols were not detected in leaf extract. In case of fruit extract, phytochemicals viz. alkaloids, flavonoids, tannins and glycosides were detected.

Plants are considered as one of the promising alternates for modern drugs as the use of antibiotics is accompanied with some drawbacks such as high cost, emergence of resistant strains of pathogenic bacteria and possible side effects. Plants have been widely used worldwide to cure diseases caused by pathogenic bacteria. It is evident from several studies that crude extracts and isolated metabolites from plants exhibit antibacterial activity against several pathogenic bacteria including antibiotic resistant bacteria.^{6,23,27-34} In this study, both leaf and fruit extracts of *D. gelonioides* were effective in causing inhibition of test bacteria as

evidenced by the zones of inhibition around wells. Among bacteria, marked susceptibility was shown by *B. cereus* while *E. coli* was inhibited to least extent. Leaf extract was found to inhibit test bacteria to greater extent than fruit extract. Susceptibility of gram negative

bacteria to extracts was in the order: *S. flexneri* > *S. typhimurium* > *E. coli*. Inhibition of bacteria by reference antibiotic was high when compared to leaf and fruit extracts. DMSO was shown to display no inhibitory activity (Table 1).

Table 1: Inhibition of test bacteria by leaf and fruit extracts of *D. gelonioides*

Treatment	Zone of inhibition in cm (Mean \pm S.D; n=3)			
	<i>B. cereus</i>	<i>E. coli</i>	<i>S. flexneri</i>	<i>S. typhimurium</i>
DMSO	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00	0.00 \pm 0.00
Antibiotic	3.23 \pm 0.05	2.93 \pm 0.05	3.10 \pm 0.00	3.13 \pm 0.05
Leaf extract	2.40 \pm 0.00	1.66 \pm 0.05	1.86 \pm 0.05	1.76 \pm 0.05
Fruit extract	2.13 \pm 0.05	1.40 \pm 0.00	1.60 \pm 0.00	1.50 \pm 0.00

Interest in botanicals with antifungal activity against phytopathogens is expanded due to several drawbacks of synthetic fungicides such as high cost, deleterious effects on soil flora and fauna, adverse effects on human health and onset of resistance in fungal pathogens. A number of studies have revealed the potential of botanical extracts to inhibit a range of phytopathogenic fungi including seed-borne fungi.^{6,35-40} In the present study, both leaf and fruit extracts of *D. gelonioides* were effective in suppressing mycelial growth of both test fungi. A considerable reduction in the mycelial growth of test fungi was observed in plates poisoned with leaf and fruit extracts. Among extracts, marked antifungal activity was shown by leaf extract when compared to fruit extract. Both leaf and fruit extracts caused >50% inhibition of test fungi (Figure 2; Table 2). Inhibitory activity of extracts was slightly higher against *Bipolaris* sp. when compared to their inhibitory potential against *A. niger*. In an earlier study by Nagabhushan and Raveesha¹⁰, the aqueous extract of *D. gelonioides* did not show antifungal activity against yeast, molds and dermatophytes.

Table 2: Colony diameter of test fungi in control and poisoned plates

Extract	Colony diameter of fungi in cm	
	<i>A. niger</i>	<i>Bipolaris</i> sp.
Leaf extract	2.26 \pm 0.05	1.70 \pm 0.00
Fruit extract	2.50 \pm 0.00	2.03 \pm 0.05
Control	5.43 \pm 0.05	4.56 \pm 0.05




Figure 2: Extent of inhibition of test fungi by leaf and fruit extracts

Scavenging of stable, organic, nitrogen centred DPPH radical is one of the extensively used in vitro radical scavenging assays for evaluating antiradical nature of plant extracts.^{6,41-45} In the present study, both leaf and fruit extracts of *D. gelonioides* showed a concentration dependent scavenging activity against DPPH free radicals. A scavenging activity of >50% was observed at extraction concentration of 25 μ g/ml and higher. At 200 μ g/ml concentration, only leaf extract displayed >90% scavenging of radicals. The scavenging activity observed was in the order: Ascorbic acid > leaf extract > fruit extract (Figure 3). Although the scavenging potential of ascorbic acid was higher, it is evident from the results obtained that both leaf and fruit extracts of *D. gelonioides* possess radical scavenging potential, hence, the extracts can act primary antioxidants.

The ferric reducing capability of extracts/compounds is considered as a significant indicator of antioxidant activity and the reducing ability is mainly due to the presence of reductones. Ferric reducing assay is widely used to evaluate antioxidant activity of plant extracts.^{6,46-51} In the present study, we evaluated the reducing capacity of leaf and fruit extracts of *D. gelonioides* by the measurement of the absorbance due to the formation of Perl's Prussian blue complex on addition of excess of ferric ions. An increase in the absorbance of reaction mixture with an increase in the concentration of extracts was observed and indicated the reducing ability of leaf and fruit extracts (Figure 4). The reducing potential observed was in the order: Ascorbic acid > leaf extract > fruit extract.




Figure 3: DPPH radical scavenging activity of leaf and fruit extracts




Figure 4: Ferric reducing activity of leaf and fruit extracts

CONCLUSIONS

Leaf extract was effective in displaying antibacterial, antifungal and antioxidant activity to higher extent than fruit extract. Based on the results obtained, it can be concluded that the plant *D. gelonioides* can be used in the treatment of infectious diseases, to manage seed-borne fungal pathogens and oxidative damage. The observed bioactivities could be attributed to the phytochemicals that have been detected in the plant. Further studies are to be undertaken to isolate and characterize bioactive principles from the plant.

ACKNOWLEDGEMENTS

Authors thank H.O.D, Dept. of Microbiology, and Principal, S.R.N.M.N College of Applied Sciences, Shivamogga for providing facilities to conduct and complete the work. Authors are thankful to N.E.S, Shivamogga for moral encouragement. Authors extend sincere gratitude for immense support given by Dr. Vinayaka K.S in collection and identification of the plant.

REFERENCES

- Vedavathy S, Scope and importance of traditional medicine, *Indian J Tradit Know*, 2003; 2(3):236-239.
- Cheikhoussef A, Shapi M, Matengu K, Ashekele HM, Ethnobotanical study of indigenous knowledge on medicinal plant use by traditional healers in Oshikoto region, Namibia, *J Ethnobiol Ethnomed*, 2011; 7:10.
- Borokini TI, Lawal IO, Traditional medicine practices among the Yoruba people of Nigeria: A historical perspective, *Journal of Medicinal Plants Studies*, 2014; 2(6):20-33.
- Bhat RB, Medicinal plants and traditional practices of Xhosa people in the Transkei region of Eastern Cape, South Africa, *Indian J Tradit Know*, 2014; 13(2):292-298.
- Kumar N, Medicinal plants used in Unani system of medicine for various disorders tehsil Joginder Nagar (H.P), *Int J Res Biol Sci*, 2014; 4(2) 41-45.
- Raghavendra HL, Kekuda PTR, Akarsh S, Ranjitha MC, Ashwini HS, Phytochemical analysis, antimicrobial and antioxidant activities of different parts of *Pleocaulus sessilis* (Nees) Bremek (Acanthaceae), *Int J Green Pharm*, 2017; 11(2):98-107.
- https://www.nhp.gov.in/introduction-and-importance-of-medicinal-plants-and-herbs_mtl
- Noumi E, Treating fibromyoma with herbal medicines in South Cameroon, *Indian J Tradit Know*, 2010; 9(4):736-741.
- Yetein MH, Houessou LG, Lougbegnon TO, Teko O, Tente B, Ethnobotanical study of medicinal plants used for the treatment of malaria in plateau of Allada, Benin (West Africa), *J Ethnopharmacol*, 2013; 146:154-163.
- Nagabhushan, Raveesha KA, Ethnobotanical survey and scientific validation of medicinal plants used in the treatment of fungal infections in Agumbe region of Western Ghats, India, *Int J Pharm Pharm Sci*, 2015; 7(4):273-277.
- Fang L, Ito A, Chai H, Mi Q, Jones WP, Madulid DR, Oliveros MB, Gao Q, Orjala J, Farnsworth NR, Soejarto DD, Cordell GA, Swanson SM, Pezzuto JM, Kinghorn DA, Cytotoxic constituents from the stem bark of *Dichapetalum gelonioides* collected in the Philippines, *J Nat Prod*, 2006; 69(3):332-337.
- Long C, Aussagues Y, Molinier N, Marcourt L, Vendier L, Samson A, Poughon V, Mutiso PBC, Ausseil F, Sautel F, Arimondo PB, Massiot G, Dichapetalins from *Dichapetalum* species and their cytotoxic properties, *Phytochemistry*, 2013; 94:184-191.
- Appiah AM, Phytochemical constituents and biological activity studies of the stem of *Dichapetalum madagascariense*, M.Phil thesis, University of Ghana, Legon, Ghana, 2013.
- Jing S, Luo S, Li C, Hua J, Wang Y, Niu X, Li X, Liu Y, Huang C, Wang Y, Li S, Biologically active dichapetalins from *Dichapetalum gelonioides*, *J Nat Prod*, 2014; 77(4):882-893.
- Chama MA, Dziwornu GA, Waibel R, Osei-Safo D, Addae-Mensah I, Otchere J, Wilson M, Isolation, characterization, and anthelmintic activities of a novel dichapetalin and other constituents of *Dichapetalum filicaule*, *Pharm Biol*, 2016; 54(7):1179-1188.
- Osei-Safo D, Dziwornu GA, Salgado A, Sunassee SN, Chama MA, Bi- and bisbibenzyls from the roots of *Dichapetalum heudelotii* and their antiproliferative activities, *Fitoterapia*, 2017; 122:95-100.
- Bhat GK, Flora of South Kanara, Akriti Prints, Mangalore, India, 2014b, Pp 465.
- Homer FA, Reeves RD, Brooks RR, Baker AJM, Characterization of the nickel-rich extract from the nickel hyperaccumulator *Dichapetalum gelonioides*, *Phytochemistry*, 1991; 30(7):2141-2145.
- Chung LY, Yap KF, Mustafa MR, Goh SH, Imiyabir Z, Muscarinic receptor activity of some Malaysian plant species, *Pharm Biol*, 2005; 43(8):672-682.
- Bhandary SK, Kumari S, Bhat VS, Sharmila KP, Bekal MP, Preliminary phytochemical screening of various extracts of *Punica granatum* peel, whole fruit and seeds, *Nitte Univ J Health Sci*, 2012; 2:34-38.
- Pandey A, Tripathi S, Concept of standardization, extraction and pre phytochemical screening strategies for herbal drug, *J Pharmacogn Phytochem*, 2014; 2(5) 115-119.

22. Cowan MM, Plant products as antimicrobial agents, Clin Microbiol Rev, 1999; 12(4):564-582.
23. Nascimento GGF, Locatelli J, Freitas PC, Silva GL Antibacterial activity of plant extracts and phytochemicals on antibiotic resistant bacteria, Braz J Microbiol, 2000; 31 247-256.
24. Kennedy DO, Wightman EL, Herbal extracts and phytochemicals: Plant secondary metabolites and the enhancement of human brain function, Adv Nutr, 2011; 2:32-50.
25. Li J, Zhao W, Li J, Xia W, Lei L, Zhao S, Research methods to study plant secondary metabolic pathways and their applications in the analysis of the biosynthetic pathway of stilbenes from *Polygonum multiflorum* – A review, Plant Omics Journal, 2014; 7(3):158-165.
26. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA, Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts, Plants, 2017; 6:42.
27. Srinivasan D, Nathan S, Suresh T, Perumalsamy LP, Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine, J Ethnopharmacol, 2001; 74(3): 217-220.
28. Ghosh A, Das BK, Roy A, Mandal B, Chandra G, Antibacterial activity of some medicinal plant extracts, J Nat Med, 2008; 62(2):259-262.
29. Aiyegoro OA, Okoh AI, Use of bioactive plant products in combination with standard antibiotics: Implications in antimicrobial chemotherapy, J Med Plants Res, 2009; 3(13):1147-1152.
30. Abdullah EM, Plants: An alternative source for antimicrobials, J Appl Pharm Sci, 2011; 1(6):16-20.
31. Djeussi D, Noumedem JAK, Seukey JA, Fankam AG, Voukeng IK, Tankeo SB, Nkuete AHL, Kuete V, Antibacterial activities of selected edible plants extracts against multidrug-resistant Gram-negative bacteria, BMC Complementary Altern Med, 2013; 13:164.
32. Shaik G, Sujatha N, Mehar SK, Medicinal plants as source of antibacterial agents to counter *Klebsiella pneumoniae*, J Appl Pharm Sci, 2014; 4(1):135-147.
33. Saranraj P, Sivasakthi S, Medicinal plants and its antimicrobial properties: A review, Global J Pharmacol, 2014; 8(3):316-327.
34. Islam R, Rahman MS, Hossain R, Nahar N, Hossain B, Ahad A, Rahman SM, Antibacterial activity of combined medicinal plants extract against multiple drug resistant strains, Asian Pac J Trop Dis, 2015; 5(S1):S151-S154.
35. Satish S, Mohana DC, Ranhavendra MP, Raveesha KA, Antifungal activity of some plant extracts against important seed borne pathogens of *Aspergillus* sp, J Agric Technol, 2007; 3(1):109-119.
36. Dellavalle PD, Cabrera A, Alem D, Larranaga P, Ferreira F, Rizza MD, Antifungal activity of medicinal plant extracts against phytopathogenic fungus *Alternaria* spp, Chilean J Agric Res, 2011; 71(2):231-239.
37. Wulff EG, Zida E, Torp J, Lund OS, *Yucca schidigera* extract: a potential biofungicide against seedborne pathogens of sorghum, Plant Pathol, 2012; 61(2):331-338.
38. Lakshmeesha TR, Sateesh MK, Vedashree S, Sofi MS, Antifungal activity of some medicinal plants on Soybean seed-borne *Macrophomina phaseolina*, J Appl Pharm Sci, 2013; 3(2):84-87.
39. Ahmad L, Pathak N, Zaidi RK, Antifungal potential of plant extracts against seed-borne fungi isolated from barley seeds (*Hordeum vulgare* L.), J Plant Pathol Microbiol, 2016; 7:350.
40. Kekuda PTR, Raghavendra HL, Antifungal activity of *Helichrysum buddleioides* DC. against seed borne fungi, EC Microbiology, 2017; 6(2):54-59.
41. Katalinic V, Milos M, Kulisic T, Jukic M, Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols, Food Chem, 2006; 94(4):550-557.
42. Dudonne S, Vitrac X, Coutiere P, Woillez M, Merillon J, Comparative study of antioxidant properties and total phenolic content of 30 plant extracts of industrial interest using DPPH, ABTS, FRAP, SOD, and ORAC assays, J Agric Food Chem, 2009; 57(5):1768-1774.
43. Khan MSS, Syeed SH, Uddin HF, Akter L, Ullah MA, Jahan S, Rashid MH, Screening and evaluation of antioxidant, antimicrobial, cytotoxic, thrombolytic and membrane stabilizing properties of the methanolic extract and solvent-solvent partitioning effect of *Vitex negundo* bark, Asian Pac J Trop Dis, 2013; 3(5):393-400.
44. Tel G, Ozturk M, Duru ME, Turkoglu A, Antioxidant and anticholinesterase activities of five wild mushroom species with total bioactive contents, Pharm Biol, 2015; 53(6):824-830.
45. Driss D, Kaoubaa M, Mansour RB, Kallel F, Abdelmalek BE, Chaabouni SE, Antioxidant, antimutagenic and cytotoxic properties of essential oil from *Corchorus olitorius* L. flowers and leaf, Free Rad Antiox, 2016; 6(1):34-43.
46. Jayanthi P, Lalitha P, Reducing power of the solvent extracts of *Eichhornia crassipes* (Mart.) Solms, Int J Pharm Pharm Sci, 2011; 3(3):126-128.
47. Shanta MA, Ahmed T, Uddin MN, Majumder S, Hossain MS, Rana MS, Phytochemical screening and in vitro determination of antioxidant potential of methanolic extract of *Streospermum chelonoides*, J Appl Pharm Sci, 2013; 3(3):117-121.
48. Pavithra GM, Siddiqua S, Naik AS, Kekuda PTR, Vinayaka KS, Antioxidant and antimicrobial activity of flowers of *Wendlandia thyrsoides*, *Olea dioica*, *Lagerstroemia speciosa* and *Bombax malabaricum*, J Appl Pharm Sci, 2013; 3(6):114-120.
49. Krishnamoorthy K, Krishnaswamy T, Subramaniam P, Sellamuthu M, Quantification of phytochemicals and in vitro antioxidant potential of various solvent extracts of certain species of Acanthaceae, Int J Green Pharm, 2014; 8(1):58-64.
50. Phatak RS, Hendre AS, Total antioxidant capacity (TAC) of fresh leaves of *Kalanchoe pinnata*, J Pharmacogn Phytochem, 2014; 2(5):32-35.
51. Ahmed D, Khan MM, Saeed R, Comparative analysis of phenolics, flavonoids, and antioxidant and antibacterial potential of methanolic, hexanic and aqueous extracts from *Adiantum caudatum* leaves, Antioxidants, 2015; 4:394-409.