

Available online on 15.09.2018 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Review Article

ETHNOPHARMACOLOGICAL STUDIES OF *Polypodium vulgare* Linn.: A COMPREHENSIVE REVIEW

Afshan Khan¹, Aisha Siddiqui^{2*}, M.A Jafri³, Mohd Asif²

¹ PG Scholar, Department of Ilmul Advia, School of Unani Medical Education and Research; Jamia Hamdard, New Delhi. India

² Assistant Professor, Department of Ilmul Advia, School of Unani Medical Education and Research; Jamia Hamdard, New Delhi. India

³ Professor, Department of Ilmul Advia, School of Unani Medical Education and Research; Jamia Hamdard, New Delhi. India

ABSTRACT

Polypodium vulgare Linn. commonly called as *Bisfaij* is a medicinal plant which is used in various ailments. It is one of the most archaic and essential plant, belonging to the family Polypodiaceae. It contains organic substances such as resin, tannins, steroids, flavonoids, alkaloids, glycosides, protein, reducing sugar and inorganic substances like calcium, magnesium, potassium, sulphur, iron and chloride. It is used in Unani system of medicine for the treatment of sorethroat, stomach-ache, leprosy, melancholia, spleenomegaly, haemorrhoids, rheumatic swelling of the joints etc. Besides this, it has other pharmacological activities too such as antiepileptic, neuro-psycho-pharmacological, antipyretic, antibiotic, antiviral etc. This review paper discusses the medicinal values of *Polypodium vulgare* Linn. in Unani medicine as well as its modern pharmacology and gives a new impetus to utilize *Bisfaij* in various disorders.

Keywords: *Bisfaij*, *Polypodium vulgare* Linn, Phytopharmacology, Unani Medicine.

Article Info: Received 23 June, 2018; Review Completed 24 July 2018; Accepted 26 July 2018; Available online 15 Sep 2018



Cite this article as:

Khan A, Siddiqui A, Jafri MA, Asif M, Ethnopharmacological studies of *Polypodium vulgare* linn. : A comprehensive review, Journal of Drug Delivery and Therapeutics. 2018; 8(5):73-76

DOI: <http://dx.doi.org/10.22270/jddt.v8i5.1848>

*Address Correspondence:

Aisha Siddiqui, Assistant Professor, Department of Ilmul Advia, School of Unani Medical Education and Research; Jamia Hamdard, New Delhi. India

INTRODUCTION

Unani system of medicine is one of the venerable traditional systems of medicine. The fortitude of the system is in its comprehensive and individualistic accession to health promotion, disease prevention and treatment. *Polypodium vulgare* Linn. (*Bisfaij*), is in consideration of ancient Greeco Arab physicians like Dioscorides, Galen (131-210 A.D), Zakaria Razi (850-925 A.D). In *Zakhirae-e- Kharazmshahi*, Jurjani cited that the rhizome of this plant is a Cardioactive. Avicenna postulated that polypody is a vigorous expeller of black bile from heart and other body organs. *Polypodium vulgare* Linn. is a small, winter green perennial fern attaining a height up to 30cm, extensively dispersed throughout the world. The name is derived from Ancient

Greek poly means “many” and podion means “little foot”, hence the name polypodium arises from the peculiar shape of its rhizomes branching like a foot. The Persian name derived from Bist (twenty) & Paya (foot), which specify foot like an arthropod which has numerous legs called Arba Arbain Milipede, as the rhizome has multiple of shoots, so due to resemblance it is named as Bistpaya. The Arabic name of bisfaij is Azras-ul-Kalb which means dog’s tooth, in delusion to the toothed appearance of the leaves^{1,2,3,4}. The traditional use of polypody rhizome has been archived in several books. The polypody rhizome is used for disease of air passages such as cough, cold, adenoids. Tea made from rhizome is used for pleurisy, sorethroat, stomach ache and poultice of root for inflammation¹¹. It is also an

important drug for lung and liver diseases. The drug also have ample of pharmacological actions like antiviral, antiepileptic, antipyretic, analgesic, hypotensive,

antibiotic and sometimes is used as an insecticide also^{6, 22, 23, 24}.



Images of *Polypodium vulgare* Linn.

TAXONOMICAL CLASSIFICATION

Kingdom	:	Plantae
Division	:	Pteridophyta
Class	:	Pteridospida
Order	:	Polypodiales
Family	:	Polypodiaceae
Genus	:	<i>Polypodium</i>
Species	:	<i>vulgare</i>
Botanical name	:	<i>Polypodium vulgare</i> Linn.

VERNICULAR NAMES

Arabic: *Azrasul kalb, Saquibal Hajer, Kasirul arjil, Tashmeez, Barzia*. Urdu: *Bisfaij*. English: *Common polypod, Wall fern, Adder's fern*. Hindi: *Bisfaija, Khankali, Kala bichwa*. Persian: *Bispaik*. Latin: *Bazbodia*. Barbary: *Pishnen*. Egyptian: *Ashtiwani*^{17,18}.

DISTRIBUTION

Polypodium vulgare Linn. is a herbaceous perennial fern found all through the year. It is a native to Europe and also found in eastern Asia, Africa predominantly in northern areas. In India, it has been brought from other countries^{25, 27}. It is common in France, ascending up to an altitude of 2000m. Also, quite common in Scandinavia, mostly in southern part of the area, and some species of *Polypodium vulgare* are commencing in America and eastern Asia. It is virtually found in every country of Europe¹⁰.

MORPHOLOGICAL DESCRIPTION OF PLANT IN UNANI (MAHIYAT)

Polypodium vulgare Linn (*Bisfaij*) is a perennial, small fern growing up to a height of about 30cm with an underground stem called rhizome. They are epiphytic ferns, with a creeping, hairy, scaly or irregular rhizome

bearing fronds at interval along its length. The rhizome is flattened, yellowish-brown in colour externally and green internally. The drug is characterized by an astringent, sweet and nauseous taste and brittleness in fracture. It has a long, dull green, pinnatifid leaves, present in two rows on the upper side of the stem alternately^{34,35}.

MORPHOLOGY

Macroscopic: *Polypodium vulgare* Linn. is a winter green fern, growing in large colonies. They are epiphytic or terrestrial ferns. Its rhizome is hairy, creeping and scaly bearing fronds at interval along its length. The stem scales are triangular in shape, irregular in size, red-brown in colour, up to 4 mm long. Leaves are long, dull green, glabrous and pinnatisect to pinnatifid, alternately arranged in two rows on the upper side of the stem. The texture of leaves is somewhat leathery and herbaceous³. The crude drug consists of dried rhizomes pieces of varying lengths and to the thickness of a squill. In cross-section, they are flattened to round and yellowish brown to dirty brown externally. The upper surface is attached with tubercles and some portion of the base of fronds still adheres. The under surface is more or less spinous from the remains of broken radicles. The drug has a characteristic ferry odour and a sweet, astringent, nauseous taste, moderately hard and brittle in fracture³⁶.

Microscopic: The transverse section of the rhizome is slightly oval to round in shape. The whole ground tissue of cortex is of thick-walled arranged parallel to the epidermis. Each vascular bundle is surrounded by a thin-walled barrel shaped, single layered endodermis, followed by a single layer of starch containing cells known as pericycle. The vascular bundles are amphicribal i.e phloem surrounds xylem. The xylem cells are differentiated into proto and metaxylems, both

are lignified and thick-walled. Phloem cells are impregnated with dark cells known as conducting parenchyma. The outer most layer is single layered epidermis surrounded by the whole mass cortex and vascular bundles. A dark brown substance is present in various cells of the cortex which is probably tannins³⁶.

CHEMICAL COMPOSITION

Polypodium vulgare Linn. contains organic substances such as resins, tannins, steroids, flavonoids, alkaloids, glycosides, protein, reducing sugar and inorganic substances like calcium, magnesium, potassium, sulphur, iron and chloride^{25,36}. Polypodin A (ecdysterone), polypodin B (5 β-hydroxy ecdysterone), glucocaffeic and polypodin have been isolated²⁶. The saponin osladins, polypodosaponin, have been isolated from the rhizome of *polypodium vulgare* Linn²². Its rhizome also contains a lauric acid, butyric acid, succinic acids, hexoic acid, methyl salicylate, isovaleric and α-methylbutyric esters, fatty oil acting as an energetic purgative^{11,19}. A resin containing benzylic alcohol and its esters which is strongly anthelmintic (a glucoside samambain)^{6,7,20,21} and saponins cyclolanostanic triterpenes-cyclolaudenol have been isolated from rhizomes.

Temperament (*Mizaj*):

Hot in 2nd degree and dry in 3rd degree².

Hot in 2nd degree and dry in 1st degree^{1, 25, 36, 37}.

Hot in 3rd degree and dry in 2nd degree¹⁸.

Parts Used: Root and Rhizome³⁶.

Dose (*Miqdar Khurak*): 5-10g¹

7-14g¹⁸

3-11g³⁷

10-15g^{25, 36}

Toxicity or adverse effect (*Muzir asrat*): Harmful for lungs and kidney, also produces nausea^{32, 33}.

Correctives (*Musleh*): Gul-e-surkh (*Rosa damascena* Mill.) and Halela zard (*Terminalia chebula* Retz) are used as corrective to avoid its toxicity and adverse effects¹.

Substitute (*Badal*): Aftimoon (*Cuscuta reflexa* Linn) and Ayarij fiqra can be used as its substitute^{1,18}.

Coumpound Formulations (*Murakkab*): Itrifal Aftimoon, Itrifal Kishnizi, Itrifal Ustukhuddus, Majoon Chobchini, Majoon Seer Alvikhan, Majoon Ushba, Majoon Najah, Safoof Chobchini^{2, 25, 36}.

PHARMACOLOGICAL ACTIONS (UNANI)

Mushil-e-Sauda wa Balgham (Purgative of Black bile and Phlegm)^{25, 36}

Mufarreh (Exhilarant)¹⁸

Muqawwi Qalb (Cardio tonic)^{1, 18}

Muhallil (Anti-inflammatory)^{18, 37}

Munaffis (Expectorant)³⁸

Mudir-e-Baul (Diuretic)¹

Mulaiyan (Laxative)¹

Hazim (Digestive)¹

Mane Sara (Antiepileptic)¹

Dafe Tashannuj (Antispasmodic)³⁸

Dafe Humma (Antipyretic)³⁸

Dafe Alam (Analgesic)¹

THERAPEUTIC USES IN UNANI

Amraz-e-Balghami wa Saudawi (Diseases of Phlegm and black bile)^{1,18,38}

Bawaseer (Haemorrhoids)^{1,18}

Damaa (Asthma)³⁷

Eruptions¹

Juzam (Leprosy)^{1,18,37}

Mafasil ka dard (Joint pain)³⁷

Malikhuliya (Melancholia)^{1,18,37}

Nafkh-e-shikam (Flatulence)^{1, 38}

Qulanj (Colitis)^{1,18,25,36}

Rheumatic disorders (*Amraze hudar*)^{1,18}

Sara (Epilepsy)^{1,37}

Warm-e-Tihal (Splenomegaly)³⁷

PHARMACOLOGICAL STUDIES

Analgesic activity³¹

Antibiotic activity²²

Antiepileptic activity³¹

Antipyretic activity³¹

Antiviral activity²⁸

Insecticidal activity⁵

Hypotensive activity³¹

Neuro-psychopharmacological activity³¹

CONCLUSION

Bisfail is one of the most important medicinal rhizome retrieve from the fern *Polypodium vulgare* Linn. of family Polypodiaceae, which is used by the Physicians of Unani system of medicine, for the treatment of various diseases. The modern investigations also confirmed the antiviral, antipyretic, antiepileptic, analgesic, hypotensive and neuro-psychopharmacological activity of *Polypodium vulgare* L. From the above review it can be concluded that new researches endorse the use of rhizome in so many diseases as mentioned in Unani classical literature.

REFERENCES

1. Ghani MN, YNM. *Khazainul Advia* (Musawwer Edition). Idara Kitab-us-Shifa, New Delhi. P. 370- 371.
2. Kabeeruddin HM. *Ilmul Advia Nafisi*. Aijaz Publishing House, New Delhi, 2007. P. 94- 95.
3. Pervaiz A Dar, G Sofi, Shabir A Parray, MA Jafri. Effect of rhizome extract of Bisfayej (*Polypodium vulgare* L) on chemically induced catalepsy in mice. *International Journal of Institutional Pharmacy and Life Sciences*. 2012; 2(4):94-108.
4. Rastogi RP, Mehrotra BN. *Compendium of Indian Medicinal Plants*, CDRI, Lucknow & National Institute of Science Communication, New Delhi. 1960-1969; I: 324-325.
5. Krishnakumaran A, Schneiderman HA. Chemical control of moulting in Arthropods. *Nature* 1968, 220:601-603.
6. Mannan A, Khan RA, Asif M. Pharmacodynamic studies on *Polypodium vulgare* Linn. *Indian Journal of Experimental Biology* 1989, 27:556-560.
7. Chevallier A. *The Encyclopedia of Medicinal Plants*. Dorling Kindersley, London, 1996, 252.
8. Dymogk W, Warden C.J.H, Hooper D. *A History of the principal Drugs of Vegetable Origin, Met with in British India .Part -III*. M/S Bishen singh Mahendra Pal Singh 23, Dehra Dun 621-622.
9. Rastogi R.P, Mehrotra B.N. *Compendium of Indian Medicinal Plants Vol-I*, CDRI, Lucknow & National Institute of Science Communication, New Delhi. 1960-1969, 324-325.
10. Ollgaard B, Tind K. *Scandinavian Ferns*. Rhodos, Aarhus 1993, 88-91.
11. Foster S. *A Field Guide to Medicinal Plants*. Houghton Mifflin, Boston 1990, 306.
12. Bown D. *New Encyclopedia of Herbs & Their Uses*. London: DK 2002, 327.
13. Moran RC. *A Natural History of Ferns*. Timber Press, Portland/Cambridge 2004, 90-96.
14. Ryvarden L. *Norske Planter* 1.J.W. Cappelen's Forlag A.S., Oslo 1993, 43-44; lid And lid 2005.
15. Shivas MG. Contribution to the cytology and taxonomy of species of *Polypodium* In Europe and America. *Journal of the Linnean Society of botany*, 1961; 58(370):13-25, 27-38.
16. http://en.wikipedia.org/wiki/common_polypod (Retrieved 15. November 2010.)
17. Anonymous, standardization of single Drugs of Unani Medicine, New Delhi, CCRUM, 1992: Vol-I, (Part I. P. 267-71, Part III. P. 86-90), Vol.II, (Part II. P. 282- 89), Vol III (Part III. P.20-25, 79-84).
18. Ibn Baitar Ziauddin Abdullah bin Ahmad; *Al Jamae ul Mufradat lil Advia wal Aghziya* (Urdu Translation). New Delhi, CCRUM ; (YNM). P.54-56, 94-97, 129-131, 231-33, 276-77, 339-41.
19. Chopra RN, Nayar SL, Chopra IC *Glossary of Indian Medicinal Plants*. Council of Scientific & Industrial Reserach, New Delhi 1956, 201.
20. Duke JA, Ayensu ES. *Medicinal Plants of China 4*, Vol. 2. Reference Publications; Michigan 1985, 517-518.
21. Chiej R. *The Macdonald Encyclopedia of Medicinal Plants*. Macdonald & Co, Verona 1988, 244.
22. Jizba J, Herout V, Šorm F. Isolation of Ecdysterone (Crustecdysone) from *Polypodium vulgare* L. Rhizomes. *Tetrahedron Letters* 1967, 18: 1689-1691.
23. Husson GP, Vilagines R, Delaveau P. Recherche de Propriétés Antivirales de quelques Extraits D'orgine Naturelle. *Ann. Pharmaceutiques Françaises* 1986, 44: 41-48.
24. Krishnakumaran A, Schneiderman HA. Chemical control of moulting in Arthropods. *Nature* 1968, 220: 601-603.
25. Anonymous. Bisfayej-Rhizome. *Unani Pharmacopoeia of India*. Department of AYUSH Government of India. Ministry of Health and Family Education. New Delhi, 2007; II (I):29-30.
26. Rastogi RP, Mehrotra BN. *Compendium of Indian Medicinal Plants*, CDRI, Lucknow & National Institutional of Science Communication, New Delhi. 1960-1969; I: 324-325.
27. Moran RC. *A Natural History of Ferns*. Portland: Cambridge Timber Press, 2004, 90-96.
28. Husson G., Vilagines R., & Delaveau P. Antiviral activity of extract of *polypodium vulgare*. *Pharmaceutical Francaises*. 1986; 44:41-48.
29. Jizba J., Herout V. & Sorm, F. Isolation of Ecdysterone from *Polypodium vulgar* L. Rhizomes. *Tetrahedron letters*. 1967; 18: 1689-1691.
30. Mannan, A., Khan, R., & Asif, M. Pharmacodynamic studies on *Polypodium vulgare* L. *Indian Journal of Experimental Biology*. 1989; 27:556-560.
31. Ibn-e-Sina. *Al-Qanun Fil-Tibb*. (English translation) Deptt. of Islamic Studies Jamia Hamdard, Hamdard University. New Delhi. 1998. P. 51, 54, 55, 83, 96.
32. Azam MK. *Mohit-e-Azam* (Persion). Nizami Press Lucknow, 1898. P. 306-307.
33. Kabirruddin, M. H., 2000. *Makhzanul Mufradat Al Maroof Khawasul Advia*, Deoband and, Faisal Publication; P.129- 130.
34. Ghani, M. N., YNM. *Khazainul Advia*. Musawwer Edition ed. Delhi: Idara Kitabus- Shifa; P. 792- 793.
35. Anonymous. *Standardisation of single drugs of Unani medicine*, Part III. New Delhi:CCRUM, Ministry of H & FW, Govt. of India: 1992. P.79-80.
36. Hakeem, M. A., 2002. *Bustan Al Mufradat*. Musawwar Edition ed. New Delhi: Idara Kitabus-Shifa; P. 92.
37. Sina I, *Al Qanoon Fit Tibb*. New Delhi: Ejaz Publishing House, Daryaganj; 1927. P.39, 40, 61, 212.