

Available online on 15.11.2018 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Antimicrobial activity of *Ocimum sanctum* leaves extracts and oil

Mittal R^{1*}, Kumar R², Chahal HS³¹ University college of Pharmacy, Guru Kashi University, Talwandi Sabo, Bathinda, India² Shivalik College of Pharmacy, Nangal, India³ Govt Polytechnic College for girls, Patiala, India

ABSTRACT

The aim of this study was to evaluate the antimicrobial effects of ethanol, Hexane and Chloroform leaf extracts of *O. sanctum* against pathogenic bacteria to determine their potentials as antibacterial agent. *Ocimum sanctum* Linn known as Holy Basil is an aromatic plant native to the tropics of Asia and Africa being medicinally important plant in the family Lamiaceae. The family Lamiaceae is one of the most exploited medicinal plant family worldwide not only as a source of medicinal plants but also with its valuable essential oils being used as spices and flavours for various food products. Leaves have petioles, and are ovate, up to 5 cm long, usually slightly toothed. Leaves were separated from the stem, washed in clear water and dried until they were adequately dry to be ground (dried for 7 days). Dried leaves were powdered separately in an electric grinder until a homogenous powder was obtained. Ethanolic, Hexane, Chloroform extract was prepared from the powder obtained using "cold extraction method. The activity of Tulsi extract against *K. pneumonia* & *Staphylococcus aureus* was found to be higher at a concentration of 100% followed by 75% & 50% respectively. The maximum zone of inhibition was found to be 21&11mm against *Klebsiella pneumonia* and 18&15mm against *Staphylococcus aureus*. The antimicrobial efficacy of *Ocimum sanctum* leaves indicates that the plant possesses potent antimicrobial properties as well as *Ocimum* is widespread in India, it can be recommended as an easily available and renewal source of antimicrobial agent instead of synthetic chemicals.

Keywords: *Ocimum santum*, *Staphylococcus aureus*, *Klebsiella pneumonia*, Disc Diffusion

Article Info:



Cite this article as:

Mittal R, Kumar R, Chahal HS, Antimicrobial activity of *Ocimum sanctum* leaves extracts and oil, Journal of Drug Delivery and Therapeutics. 2018; 8(6):201-204 DOI: <http://dx.doi.org/10.22270/jddt.v8i6.2166>

*Address for Correspondence:

Rohit Mittal, University College of Pharmacy, Guru Kashi University, Talwandi sabo, Bathinda, India

INTRODUCTION

Ocimum sanctum Linn known as Holy Basil is an aromatic plant native to the tropics of Asia and Africa being medicinally important plant in the family Lamiaceae. The family Lamiaceae is one of the most exploited medicinal plant family worldwide not only as a source of medicinal plants but also with its valuable essential oils being used as spices and flavours for various food products. Morphologically *O. sanctum* is an erect about 75 cm tall, much branched with hairy stems and simple opposite green leaves that are strongly scented. Leaves have petioles, and are ovate, up to 5 cm long, usually slightly toothed ^{8,9,10}.

The medical world is on an immense requirement to discover novel antibiotics due to wide spread emergence of resistance among microbial pathogens against currently available antibiotics. However, traditional plants have been proved to be better source for novel antimicrobial drugs. Most of Indian plants accounts for the richest resources of

natural drugs ⁴. Historically medicinal plants have been placed at top among the source of novel drugs with antimicrobial activity. These traditional medicinal herbs have made considerable contributions to human health. In addition, plants are considered as one of the most important sources of secondary metabolites and essential oils ⁵. On one hand the use of medicinal plants proved to be economical and effective and on the other hand they are easily available and safe to use ⁶. Indian traditional medicinal system includes hundreds of medicinal plants related to multiple effects ⁷. Furthermore, studies have revealed *O. sanctum* to possess anti-inflammatory, analgesic, antipyretic, antidiabetic, hepatoprotective, hypolipidemic and antistress activities ¹¹. Extracted essential oils have also been shown to contain biologically active constituents that are insecticidal, nematicidal and fungistatic. The disc diffusion method is the commonly used technique to check the antimicrobial activity. The antimicrobial activity was tested by the method based on the principle of Kirby Bauer test. An antimicrobial is a

substance that kills or inhibits the growth of microorganisms such as bacteria, fungi, all protozoans (Merriam-Webster Online Dictionary) ¹². The aim of this study was to evaluate the antimicrobial effects of ethanol, Hexane and Chloroform leaf extracts of *O. sanctum* against pathogenic bacteria to determine their potentials as antibacterial agent.

MATERIAL AND METHODS

The study employed an *in vitro* experimental design. Tulsi leaves were obtained from courtyards and local market from Shivalik hills of Himachal area. Authenticity of plant by Dr GS Chatha, Shivalik College, Department of Botany, Nangal. Leaves were separated from the stem, washed in clear water and dried until they were adequately dry to be ground (dried for 7 days). Dried leaves were powdered separately in an electric grinder until a homogenous powder was obtained. Ethanolic, Hexane, Chloroform extract was prepared from the powder obtained using "cold extraction method" ¹. Three hundred grams of finely powdered *Ocimum sanctum* (Linn.) was then macerated

with *Tulsi (Ocimum sanctum) extract preparation*; (a) *Ocimum sanctum* plant; (b) leaves separated and dried; (c) leaves ground to powder; (d) extract obtained 100% ethanol and other solvent. It was then subjected to filtration with Whatman filter paper to obtain a clear filtrate. The filtrate so obtained was reduced at a low temperature of less than 60 °C to obtain a solid residue of *Ocimum sanctum* (Linn.) extract². From 300 grams of *Ocimum sanctum* (Linn.) powder dissolved in 1 liter of ethanol and other solvent, 18 gram of extract (residue) was obtained and thus the yield was 6% w/v. Accurately weigh 1gm of each extract was reconstituted in 10 ml of respective solvents to obtain stock solution in eppendorf tube. Further the dilutions were made with respective solvents. Accurately weigh of 10 mg of Standard Gentamycin was dissolved in 10 ml of distilled water to get 1mg/ml. The different dilutions and standard pipetted out on into the marked plates. These were left for incubation at 37°C for 24 hrs and 25°C for 36 hrs. After incubation zones of inhibition were measured (from antibiotic zone measurement scale) in mm and compared with standard.



Collection of test organism

Collection of test organism and preparation of stock culture: The following strains were obtained for the antimicrobial tests. Gram negative bacteria were *Klebsiella pneumoniae* and Gram positive bacteria were *Staphylococcus aureus*. All the microbial strains were obtained from Microbial Type Culture Collection (MTCC), Chandigarh. Nutrient broth medium was prepared and autoclaved. After the bacterial cultures were inoculated to separate flasks and incubated in shaker for 24 hours

Microbiological Assay

The test organisms included for study were gram positive *Staphylococcus aureus* and gram negative *Klebsiella pneumoniae* ³.

Preparation of Media

For 100 ml of media, 40 gm of muller-hinton agar is dissolved in 100 ml distilled water. 250 ml media prepared and autoclaved at 121°C to 15-20 min at 15 lbs/inch².

Preparation of Disk

Freshly prepared and sterilized molten media was poured onto Petri plates inside Laminar and after pouring UV light turned on to avoid contamination on plates while media

solidifying. It was left for half an hour for proper solidification.

After media gets solidified than UV light is turned off and 10µl of bacterial suspension pipetted into plates and swabbed. Sterile discs were put (with the help of forceps) on plates along with one disc of standard (6 disk was put on a plate).

RESULT



The result of antibacterial activity of *Ocimum sanctum* against *Staphylococcus aureus*, and *Klebsiella pneumoniae* with different concentration. The least zones of inhibition were displayed by the negative control and Gentamycin exhibited the widest zones of inhibition against all the bacteria. Tulsi leaves' extract showed increasing zones of inhibition. Results of the study were shown in the table 1. We observed that *O. sanctum* have a potent antibacterial activity. The ethanolic extract showed maximum antibacterial activity followed by hexane and pure oil extracts against *S. aureus* and *K. pneumoniae*. etc. (Table 1 and 2). The activity of Tulsi extract against *K. pneumoniae* & *Staphylococcus aureus* was found to be higher at a concentration of 100% followed by 75% & 50% respectively. The maximum zone of inhibition was found to be 21&11mm against *Klebsiella pneumoniae* and 18&15mm against *Staphylococcus aureus*.

Table 1: Tulsi leaves serial dilution showing the Zone of inhibition (in mm) against *Klebsiella pneumonia* (Gram negative bacteria)

	Dilution 1	Dilution 2	Dilution 3	Dilution 4	Dilution 5	Dilution 6	Gentamycin
Chloroform extract	15	12	11	9	7	7	17
Haxane Extract	6	4	4	3	3	NZ	17
Alcoholic Extract	18	16	14	14	13	10	17
Pure oil	7	7	6	5	5	4	17

Table 2: Tulsi leaves serial dilution showing the Zone of inhibition (in mm) against *Staphylococcus aureus* (Gram positive bacteria)

	Dilution 1	Dilution 2	Dilution 3	Dilution 4	Dilution 5	Dilution 6	Gentamycin
Chloroform extract	11	11	10	9	5	4	17
Haxen Extract	9	7	7	4	3	2	17
Alcoholic Extract	21	19	18	16	14	10	17
Pure oil	8	7	7	3	2	NZ	17

**Figure 1: Tulsi leaves serial dilution showing the Zone of inhibition (in mm) against *Klebsiella pneumonia* (Gram negative bacteria).****Figure 2: Tulsi leaves serial dilution showing the Zone of inhibition (in mm) against *Staphylococcus aureus* (Gram positive bacteria).**

DISCUSSION

Antibacterial activity of different *Ocimum sanctum* extracts against *Klebsiella pneumonia* (Gram positive bacteria) and *Staphylococcus aureus* (Gram negative bacteria) were studied. According to the results, all different types of extracts obtained from *Ocimum sanctum* leaves shown to be with antibacterial activity against tested microbial pathogens. *Ocimum* extract has found to be with antimicrobial properties against *Staphylococcus aureus* and *Klebsiella pneumonia*. Highest antibacterial activity was shown by ethanol and chloroform extracts against both gram-negative bacteria (*Staphylococcus aureus*) and gram-positive bacteria (*Klebsiella pneumonia*). Chloroform extract shown higher antibacterial activity against gram negative bacteria compared to gram positive bacteria and ethanolic extract shows higher activity against gram

positive bacteria compared to gram negative bacteria in the present study. The activity of Tulsi extract against *Klebsiella pneumonia* & *Staphylococcus aureus* was found to be higher at a concentration of 100% followed by 75% & 50% respectively. As the concentration of extract increases, the activity also increases and thus the zone of inhibition too increases. Significant results were shown by *Klebsiella pneumonia* and *Staphylococcus aureus*.

CONCLUSION

The present study clearly indicates that *Ocimum sanctum* is a rich source of phyto-chemical constituents. The antimicrobial efficacy of *Ocimum sanctum* leaves indicates that the plant possesses potent antimicrobial properties as well as *Ocimum* is widespread in India, it can be recommended as an easily available and renewal source of antimicrobial agent instead of synthetic chemicals.

REFERENCES

1. Mallikarjun S, Rao A, Rajesh G, Shenoy R, Pai M. Antimicrobial efficacy of Tulsi leaf (*Ocimum sanctum*) extract on periodontal pathogens: An *in vitro* study. J Indian Soc Periodontol; 2016; 20:145-50.
2. Eswar P, Devaraj CG, Agarwal P; Anti-microbial Activity of Tulsi (*Ocimum Sanctum* (Linn.)) Extract on a Periodontal Pathogen in Human Dental Plaque: An *In vitro* Study; Journal of Clinical and Diagnostic Research. 2016; 10(3):ZC53-ZC56.
3. Ali H, Dixit S; In vitro antimicrobial activity of flavanoids of *Ocimum sanctum* with synergistic effect of their combined form; Asian Pacific Journal of Tropical Disease 2012; S396-S398.
4. Hammer KA, Carson CF, Riley TV, Antimicrobial activity of essential oils and other Plant extracts. J Appl Microbiol. 1999; 86:985-990.
5. Singh V, Amdekar S, Verma O, *Ocimum Sanctum* (tulsi): Biopharmacological activities. Web Med Cent. Pharmacolo. 2010; 1:1-7.
6. Govind P, Madhuri S, Autochthonous herbal products in the treatment of cancer. Phytomedica. 2006; 7:99-104.
7. Rahal A, Singh V, Mehra D, Rajesh S, Ahmad AH, Prophylactic efficacy of *Podophyllum hexandrum* in alleviation of immobilization stress induced oxidative damage in rats. J Nat Prod 2009; 4:110-115.
8. Jirovetz, L., Buchbauer, G., Shafi M.P. and, Kaniampady M. M. Chemotaxonomical analysis of the essential oil aroma compounds of four different *Ocimum* species from southern India. European Food Research and Technology, 2003; 217(2):120-124.
9. Malima, S. M., Massaga, J. J., Malecela, M. N. and Andrew, Y. Repellence effectiveness of essential oils from some Tanzanian *Ocimum* and *Hyptis* plant species against afro-tropical vectors of malaria and lymphatic filariasis. Journal of Medicinal Plants Research, 2013; 7(11):653-660.
10. Kaya, I., Yiğit N and Benli, M. Antimicrobial activity of various extracts of *Ocimum basilicum* L. and observation of the inhibition effect on bacterial cells by use of scanning electron microscopy. African Journal of Traditional, Complementary and Alternative Medicines, 2008; 5(4):363-369.
11. Baliga, M. S., Jimmy, R., Thilakchand, K. R., Sunitha, V., Bhat, N. R., Saldanha, E. and Palatty, P. L. *Ocimum sanctum* L (Holy Basil or Tulsi) and its phytochemicals in the prevention and treatment of cancer. Nutrition and cancer, 2013; 65(sup1):26-35.
12. Gadiyar A, Ankola AV, Rajpurohit L; Evaluation of the Antimicrobial Activity of *Ocimum Sanctum* L. (Tulsi) Extract against *Streptococcus Mutans* and *Lactobacillus Acidophilus* - An *In-Vitro* Study; Int J Health Sci Res. 2017; 7(4):224-228.