

Available online on 15.05.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Synthesis and Evaluation of Cr (III), Mn (III) and Fe (III) Schiff Base of Metals Complexes

Kalpataru Halder*, Dr. Utpal Saha

Department of Biochemistry, Sri Satya Sai University of Technology & Medical Sciences, Sehore-466001, Madhya Pradesh

ABSTRACT

A novel synthesized Schiff base metal complexes Cr (III), Mn (III) and Fe (III) of 2-hydroxy-5-chloro acetophenone 2-imino-4-phenyl thiazole were prepared by condensed from 2-hydroxy-5-chloro acetophenone and 2-amino-4-phenyl thiazole have been synthesized and characterized on the basis of elemental analysis, Infrared, ¹H NMR, molar conductance and magnetic susceptibilities analysis. The Schiff base acts as a monobasic bidentate ligand commonly coordinates through the oxygen atom of phenolic OH group and the nitrogen atom of azomethine group, which is confirmed by IR spectral data. All the metal complexes have studies thermal properties and their thermal parameter.

Keyword: Schiff base, Magnetic, Thermal studies.

Article Info: Received 18 March 2019; Review Completed 22 April 2019; Accepted 25 April 2019; Available online 15 May 2019



Cite this article as:

Halder K, Saha U, Synthesis and Evaluation of Cr (III), Mn (III) and Fe (III) Schiff Base of Metals Complexes, Journal of Drug Delivery and Therapeutics. 2019; 9(3):114-116 <http://dx.doi.org/10.22270/jddt.v9i3.2609>

*Address for Correspondence:

Kalpataru Halder, Department of Biochemistry, Sri Satya Sai University of Technology & Medical Sciences, Sehore-466001, Madhya Pradesh

INTRODUCTION

The Schiff bases are widely used ligands due to their facile synthesis, significant versatility and good solubility in common solvents. The structure of these coordination complexes arises due to the interesting ligand systems containing different donor sites in heterocyclic rings. Schiff base have received much great interest, mainly because of their wide application in the field of synthesis and catalysis ¹. Thus, they have played an important applicable role in research and development of coordination chemistry as they readily form stable metal complexes in different oxidation states ² Thiazole Schiff base ligands and their metal complexes are biologically active ³. Due to its biological potency, pharmacological properties and synthetic flexibility of Schiff base derived from isonicotinic acid hydrazide ⁴. Hydrazones, heteroaroyl hydrazones ligands and their metal complexes are biologically active. Heteroaroyl hydrazones forms stable metal complexes with transition metal ions and inner transition metal ions due to complexing ability of ligand through keto-enol tautomerism and availability of other donor sites in the ligand *i.e.* isonicotinoyl hydrazide is one of the drug in chemotherapy of tuberculosis ⁵.

The aim of present investigation is to synthesize various transition metal complexes of Schiff base condensed from 2-hydroxy-5-chloroacetophenone and 2-amino-4-phenyl thiazole.

MATERIAL AND METHODS

All the chemicals were of A.R. grade and used as received. 2-hydroxy-5-chloro-acetophenone (HCA) and 2-amino-4-phenylthiazole was prepared by known methods ⁶⁻⁹. The solvents were purified by standard methods ¹⁰.

Synthesis of 2-amino-4-phenylthiazole

The synthesis of 2-amino-4-phenylthiazole prepared by known method ⁷⁻⁹. The product was filtered and crystallized from 70% ethanol, after several minutes the golden coloured product of 2-amino-4-phenylthiazole was separated out.

Yield: 75%; m.p.: 148-150°C.

Synthesis of 2-hydroxy-5-chloro acetophenone 2-imino-4-phenyl thiazole [HCAIPT]

A solution of 2-hydroxy-5-chloro acetophenone (0.02 M) in 25 ml of ethanol was added to an ethanolic solution (25 ml) of 2-amino-4-phenylthiazole (0.02 M) and the reaction mixture was refluxed on a water bath for 4 h. After cooling a pale yellow coloured crystalline solid was separated out. It was filtered and washed with ethanol, crystallized from DMF and dried under reduced pressure at ambient temperature. The purity of ligand was checked by elemental analysis and m.p. It was also characterized by IR and ¹H NMR spectral studies.

Yield: 35%; m.p. 290°C

Preparation of complexes

All the metal complexes were prepared in a similar way by following method. To a hot solution of ligand HCAIPT (0.02 M) in 25 ml of ethanol a suspension of respective metal salts was added drop wise with constant stirring. The reaction mixture was refluxed on a water bath for 4-6 h. The precipitated complexes were filtered, washed with ethanol followed by ether and dried over fused calcium chloride. Yield: 45-50%

The complexes are soluble in DMSO and DMF but insoluble in water and common organic solvents. The metal chloride content of complexes were analyzed (Table 2) by standard methods ¹¹. The ¹H NMR spectra of ligand was recorded and obtained from RSIC Chandigarh. IR spectra of the compounds were recorded on Perkin Elmer 842 spectrophotometer in the region 400-4000 cm⁻¹, Carbon, Hydrogen and Nitrogen analysis were carried out at RSIC, Punjab University, Chandigarh. The molar conductance of the complexes at 10⁻³ M dilution in DMF were determined using equipotronic digital conductivity meter EQ-660 with a cell constant 1.00 cm⁻¹ at room temperature. The magnetic moment measurement were made on a Gouy balance at room temperature using [HgCo(SCN)₄] as the calibrant. The molecular weights of the complexes were determined by Rast method. The thermo-gravimetric analysis was performed on laboratory set up apparatus in air atmosphere at 10°C min⁻¹ heating rate. The molecular weights of the complexes were determined by Rast method.

RESULT AND DISCUSSION

The Schiff base ligand HCAIPT and its complexes have been characterized on the basis of ¹H-NMR, IR spectral data,

elemental analysis, molar conductance, magnetic susceptibility measurements and thermo-gravimetric analysis data. All these values and analytical data are consistent with proposed molecular formula of ligand. All the compounds are coloured solid and stable in air. They are insoluble in water but soluble in coordinating solvents like DMF and DMSO. The molar conductance values in DMF (10⁻³M) solution at room temperature (Table 2) shows all the complexes are non-electrolytes. The ¹H NMR spectra of ligand HCAIPT shows signals at δ 11.26, (1H, s phenolic OH), 7.41, 7.40, 7.39 and 7.38 (4H, m, phenyl) δ 6.51, 6.50, and 6.58 (3H, s Phenyl), 6.62 (¹H s thiophene), and 2.16 (3H, s, methyl) ¹²⁻¹⁵. IR spectra of ligand and metal complexes summarized in table 3. As per observation ν (C=N) peaks at 1618 cm⁻¹ and absence of C=O peak at around 1700-1730 cm⁻¹ indicates the Schiff base formation ¹⁶⁻¹⁹.

Thermo-gravimetric Parameter

Thermo-gravimetric study indicates all the complexes are stable up to 60-70°C. All the complexes show half decomposition temperature (Table 4). The Thermal activation energy, Frequency factor, Entropy change and Free Energy change was calculated by Freeman- Carroll, ¹⁸ Horowitz-metzger ¹⁹ and Broido ²⁰ method.

CONCLUSIONS

The analysis of magnetic moment and electronic spectral data shows characterization and structural changes in metal complexes. Thermal studies concluded conformed structure of metal complexes from the study of water loss in metal complexes and other parameter.

Table 1: Analytical data of the Ligands

S. No	Ligand	Molecular Formula	Formula Weight	Color and nature	Elemental Analysis				
					C % found (Cal.)	H % Found (Cal.)	N % Found (Cal.)	Cl % Found (Cal.)	S % Found (Cal.)
1.	HCAIPT	C ₁₇ H ₁₂ N ₂ O ₂ OSCl	327.6	Yellow Crystalline	61.38 (62.27)	03.10 (03.66)	08.24 (08.54)	10.21 (10.83)	09.13 (09.79)

Table 2: Analytical Data and Molar Conductance of the Compounds.


Compounds	Colour	Mol. wt.	Analysis % Found (calc.)					μ_{eff} B.M.	$\Omega^{-1}\text{cm}^2\text{mol}^{-1}$
			M	C	H	N	Cl		
[CrL ₂ (H ₂ O)Cl] H ₂ O	Green	776.7	6.32 (6.69)	52.25 (52.52)	3.36 (3.60)	6.81 (7.20)	9.08 (9.14)	3.90	18.2
[MnL ₂ (OAc)] H ₂ O	Brown	804.1	6.20 (6.82)	53.21 (53.72)	3.10 (3.48)	6.31 (6.96)	9.32 (4.41)	4.2	18.3
[FeL ₂ (H ₂ OCl)] H ₂ O	Black	780.6	6.82 (7.14)	52.01 (52.26)	3.32 (3.58)	6.73 (7.17)	9.01 (9.09)	5.2	22.1

Table 3: IR Spectra of Ligand and Metal Complexes.


Compound	$\nu(\text{O-H})$ hydrogen bonded	$\nu(\text{C=N})$ Imine	$\nu(\text{C-O})$ phenolic	$\nu(\text{C-S})$	$\nu(\text{M-O})$	$\nu(\text{M-N})$
HCAIPT	3109	1618	1514	1122	-	-
$[\text{CrL}_2(\text{H}_2\text{O})\text{Cl}]\text{H}_2\text{O}$		1590	1506	475	409	1115
$[\text{MnL}_2(\text{OAc})]\text{H}_2\text{O}$		1562	1462	498	420	1090
$[\text{FeL}_2(\text{H}_2\text{O})\text{Cl}]\text{H}_2\text{O}$		1602	1504	512	440	1080

Table 4: Thermal decomposition data of HCAIPT and its complexes.

Compound	Half Decomposition Temperature (°C)	Activation Energy (kJ mole ⁻¹)			Frequency Factor Z (sec ⁻¹)	Entropy Change -ΔS (J mol ⁻¹ K ⁻¹)	Free Energy Change ΔF (kJ mol ⁻¹)
		B*	H-M**	F-C***			
HCAIPT	266.43	2.88	5.76	4.82	98.79	213.62	128.84
$[\text{CrL}_2(\text{H}_2\text{O})\text{Cl}]\text{H}_2\text{O}$	592.66	11.52	14.39	11.56	228.46	207.66	199.68
$[\text{MnL}_2(\text{OAc})]\text{H}_2\text{O}$	443.85	4.95	9.91	9.84	196.27	208.36	168.49
$[\text{MnL}_2(\text{OAc})]\text{H}_2\text{O}$	443.85	4.95	9.91	9.84	196.27	208.36	168.49



SCHEME 1: SYNTHESIS OF THIAZOLE



SCHEME 2 SYNTHESIS OF HCAIPT

REFERENCES

- Canali L, Sherrington DC. Chem Soc Rev 1999; 28:52.
- Prasad KS, ShivaKumar L, Chandan S, Jayalakshmi B, Revanasiddappa HD. Spectrochim. Acta, Part A, 2011; 81:276.
- Kumar S, Dhar D, Saxena P, J Sci and Indu Research 2009; 68:181.
- Hawlder MB, Begum MS. Indian J Chem 2004; 43A: 2352.
- Sharma RP, Kothari AK, Sharma NK. Indian J Derm Vener Lepr 1995; 61:26.
- Ballhausen CJ, Gray HB. Inorg Chem 1962; 1:11.
- Furniss BS, Hannaford AJ, Smith PWG, Tatchell AR. Vogel's practical organic chem. Edn 5, Logman Scientific Technical, John Wiley and Sons 1989.
- Vogel AI. A Text book of quantitative inorganic chemistry, Edn 3, ELBS, London, 1961.
- Maurya MR, Gopinathan C. Indian J Chem 1996; 35A: 701.
- Jayaramadu M, Reddy KH. Indian J Chem 1999; 38A: 1173.
- Sastry PS, Rao TR. Proc Indian Acad Sci 1995; 107(1):25.
- Mahajan RK, Patial VP. J Indian Council Chem 2001; 18(1):4.
- Maurya MR, Antony DC, Gopinathan S, Gopinathan C. Bull Chem Soc Jpn 1995; 68:554.
- Viroopakshapp J, Vithalrao D. J Indian Chem Soc 1996; 73:531.
- Figgis BN. "Introduction to ligand field", weley Estern Ltd., 1976.
- Sacconi L. J Ame Chem Soc 1954; 76:3400.
- Panda AK, Dash DC, Mishra P, Mohnaty H. J Indian Chem Soc 1996; 35A:324.
- Mallikarjun K. E J Chem 2004; 1(2):105.
- Horowitz H, Metzger G. Anal Chem 1963; 35:1464.
- Broido A. J Polym Sci 1964; A2:1761.