

Available online on 15.05.2019 at <http://jddtonline.info>

# Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

## Evaluation of Anxiolytic Activity of Pitavastatin in Male Albino Mice

Premkumar Chiluveru\*<sup>1</sup>, G. Thirupathi<sup>2</sup>, T. Ravi Chander<sup>2</sup>

1. Nottingham Trent University, Clifton campus, Nottingham, NG11 8NS
2. Vaagdevi Pharmacy College, Bollikunta, Warangal, India

### ABSTRACT

To assess the anxiolytic activity of Pitavastatin in male albino mice. To compare the anxiolytic activity of Pitavastatin with standard drug as diazepam. Anxiety is an emotional state, unpleasant in nature, associated with uneasiness, discomfort and fear about something. This Anxiety higher in developed countries than developing countries, moreover women are more prone to this. At present we find number of marketed drugs all that are producing somewhat tolerated compounds. As by with that condition we conducted this study comparison with pitavastatin with diazepam. Pitavastatin and Diazepam were dissolved in saline. Control group mice were injected with saline. All drugs and saline were injected intraperitoneally, placed all in plus maze method and collected the data. Elevated plus maze Pitavastatin at dose 30 mg/kg and 50mg/kg increased number of open arm and total arm entries, time spent in open arm and decreased time spent in closed arm. The results suggest that the significant dose-dependent antianxiety activity of Pitavastatin is similar to the behavioral effects of Diazepam.

**Keywords:** Anxiolytic activity, Pitavastatin, Diazepam.

**Article Info:** Received 25 March 2019; Review Completed 02 May 2019; Accepted 06 May 2019; Available online 15 May 2019



### Cite this article as:

Chiluveru P, Thirupathi G, Ravi Chander T, Evaluation of Anxiolytic Activity of Pitavastatin in Male Albino Mice, Journal of Drug Delivery and Therapeutics. 2019; 9(3):269-271 <http://dx.doi.org/10.22270/jddt.v9i3.2648>

### \*Address for Correspondence:

Prem Kumar Chiluveru, Nottingham Trent University, Clifton campus, Nottingham, NG11 8NS

### INTRODUCTION

Anxiety is an emotional state, unpleasant in nature, associated with uneasiness, discomfort and fear about something.<sup>1</sup> Different types of Anxiety disorders includes panic disorder, generalized anxiety disorder, post-traumatic stress disorder, phobias, and separation anxiety disorder. Prevalence estimates of anxiety disorders are generally higher in developed countries than in developing countries.<sup>2</sup> Women are more prone for anxiety disorder than in men.<sup>3</sup> \$42.3 billion was the annual cost of anxiety disorder in US in 1990s, and most of it was due to non-psychiatric medical treatment costs.<sup>4</sup> The primary treatments for anxiety-related disorders include the SSRIs, SNRIs, benzodiazepines, non-benzodiazepines and beta adrenergic antagonists. But side effects such as day time sedation and dependence are generally associated with the use of benzodiazepines.<sup>5</sup> There is a need of new anxiolytic drug with lesser side effects and immediate onset of action. Clinical studies shows direct effects of statins in improvement of endothelial dysfunction, reduction in atherogenesis, Alzheimer's disease, dementia, and anti-inflammatory effects which are unrelated to their cholesterol reducing effects.<sup>6,7,8</sup> The aim of this study is to assess the anxiolytic effects of rosuvastatin.

### MATERIALS AND METHODS

This study was conducted at the Department of Pharmacology. Drugs: Pitavastatin and Diazepam were dissolved in saline. Control group mice were injected with saline intraperitoneally

**Animals:** Twenty four male Albino mice weighing 18- 25g (2 - 2.5months old) were used in this study. All animals were bred in standard laboratory conditions comprising 12-hour light/dark cycle and were allowed to access for food and water ad libitum. Select six mice in each were grouped into 4 groups as follows.

Group 1.(control group): Saline (i.p route)

Group 2.1 mg/kg Diazepam (i.p route)

Group 3.30 mg/kg Pitavastatin (i.p route)

Group 4.50 mg/kg Pitavastatin (i.p route)

**Experimental procedure:** Drugs and/or saline were injected i.p. route in to the Mice an hour before the experiment. The plus maze tests were used to assess the anxiolytic effects of Pitavastatin. It consists of two open arms and two closed arms connected with the central platform. The apparatus is elevated at a height of 25 cm above the ground level. The animals are placed at center of the maze facing towards open arm.

Parameters that are recorded are for five minutes. (1) mouse preference towards open or closed arm. (2) Frequency of entries in open arm. (3) Average duration of animal in open arm. Statistical Analysis: Within the group comparison was done using analysis of variance with multiple comparisons by post hoc Dunnett's t-tests. Between the two groups was done by Dunnett's two-sided t-tests. Statistical analysis was done using SPSS. P < 0.05 level were considered as statistically significant.

**RESULTS**

Administration of Diazepam has increased the number of open arm and total arm entries, fraction ratio of open/total arm entries, period spent in open arm but reduced period spent in closed arm. Pitavastatin at the dose of 30mg/kg and 50mg/kg has shown similar result by increasing number of open arm and total arm entries, percentage ratio of open/total arm entries, time spent in open arm, analogous results to that of standard drug Diazepam.

**Table 1: The number of entries (open and total) of mice in EPM (n=6)**

Groups	Mean±SEM		
	Number of entries in open arm	Total entries	% ratios of open/total arm entry
Control	5.83±0.54	28.00±0.63	20.70±1.51
Diazepam	33.83±0.54*	47.16±0.54*	71.74±0.93*
Pitavastatin30mg/kg	25.01±0.37*	41.16±0.48*	60.74±0.89*
Pitavastatin50mg/kg	29.17±0.75*	44.33±0.96*	65.77±0.49*

\*p<0.001when compared to control

**Table 2: The time spent by mice on EPM in open and closed arms (n=6)**

Group	Mean±SEM	
	Time spent in open arm in seconds	Time spent in closed arm in seconds
Control	4.33±0.62	265.67±0.63
Diazepam	132.330±.96*	167.67±0.96*
Pitavastatin30mg/kg	93.0±0.86*	207.0±0.86*
Pitavastatin50mg/kg	118.67±0.80*	181.33±0.80*

\*p<0.001when compared to control




Figure 1: Effect of Pitavastatin (RS) shows antianxiety effect in open arm entries




Figure 2: Effect of Pitavastatin shows antianxiety effect in total entries

## DISCUSSION

The elevated plus maze model works on the basis that, inhibition of normal behaviour will occur in the presence of brightly lit, unfamiliar, unprotective environment. It is one of the most widely used and easy animal anxiety model. Hence it was chosen to investigate the anxiolytic activity of Pitavastatin. Site of action of drugs is gamma-aminobutyric acid receptor complex. Standard drug Diazepam increased the number of entries in open arm and the total time spent in the open arms, hence confirming its anxiolytic effects.<sup>11,12</sup> In this study Pitavastatin at the doses 30,50 mg/kg increases the number of entries in open arm in comparison to control and significantly decreased the number of entries in closed arm compared to that of control. Diazepam showed decreased fear and increase in exploratory behaviour. The BZDs are comparatively safe and most commonly used anxiolytic agents. Site of action is BZD-GABA receptors; the role of GABA in anxiety is well established. The behavioural changes by Pitavastatin was comparable to that produced by diazepam and suggestive of anxiolytic effect of the statins. The results of our study are consistent with earlier studies. However, several human studies have shown contrary results demonstrating that there were no association between statins and anxiety and depression. The contrary outcomes may mainly result from their different methodology and follow-up.

## CONCLUSION

The results suggest that the behavioural disinhibitory effects of Pitavastatin is similar to the behavioural effects of Diazepam

## REFERENCES

1. Rao EV. Modern approaches to herbal medicine. East Pharmacist, 2000; 5:35-38.
2. Sagrawat H, Mann AS, Kharya MD. Pharmacological potential of *Eugenia jambolana*, A review. Pharmacog. Mag. 2006; 2(6):96 -105.
3. Martin KP, Benna MR, Joseph D. High frequency axillary bud multiplication and ex - vitro rooting of *Wedelia chinensis* (Osbeck) Merr, A medicinal plant. Ind. J Exp. Biol. 2003; 41(3):262 -266.
4. Kirtikar KR, Basu BD. Indian Medicinal Plants. Dehradun, International book distributors, 2006, 1364 -1345.
5. Anonymous. A Dictionary Indian Raw Materials and Industrial Products, The Wealth of India, Raw Materials, Council of Scientific and Industrial Research, 2005, 567 -568.
6. Anonymous. The Wealth of India, Raw Materials, Publication and Information Directorate, New Delhi, 1948, 687 -688.
7. Suresh V, Kumar RM, Suresh A, Kumar NS, Arunachalam G, Umasankar KI, CNS activity of ethanol extract of *Wedelia chinensis* in experimental animals. Int J Pharm. Sci. Nanotechnol. 2010; 3(1):881 - 886.
8. Mathew KM. Flora of Tamilnadu -carnatic. Trichirapalli, St. Josephs College, 1983, 392.
9. Linn SC, Linn CC, Linn YH, Shyuu SJ. Hepatoprotective effect against liver injuries. Ame. J Chin. Med. 1994; 22(3):155 -168.
10. Manjamalai A, Shakur AS, Haridas A, Berlin G. Evaluation of antifungal and anti - inflammatory effects on methanolic extract of *Wedelia chinensis*, Int J Pharm. Biol. Res. 2011; 2(1):30 -37.
11. Yang LL, Yen KY, Kiso Y, Hikino H. Antihepatotoxic actions of Formosan plant drugs. J Ethanopharmacol. 1987; 19 :103 -10.
12. Pate RM, Patel NJ. In vitro antioxidant activity of coumarin compounds by DPPH, super oxide and nitric oxide free radical scavenging methods. J Adv. Pharm. Edu. Res. 2011; 1:52 -68.

