

Available online on 30.08.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Research Article

Evaluation of Methanolic Extract of *Clitoria ternatea* Hepatoprotective & Nephroprotective Activity in Rats

Srikanta Chandra^{1*}, Avik Das², Tathagata Roy³, Preeta Bose⁴, Lucky Mukherjee⁵, Jyotirmay Samanta⁶, Rumpa Banerjee⁷, Rakhi Bakuli⁸, Mayukh Jana⁹, Debraj Mukhopadhyay¹⁰

1. Assistant Professor of Jakir Hossain Institute of Pharmacy, Department of Pharmacology, Raghunathganj, Murshidabad, West Bengal, 742225
2. Assistant Professor of Gupta College of Technological Sciences, Department of Pharmacology, Asansol, West Burdwan, West Bengal, 713301
3. Department of Pharmaceutical Technology, JIS University, Kolkata, West Bengal, 700109
4. Department of Pharmaceutical Technology, JIS University, Kolkata, West Bengal, 700109
5. Bengal College of Pharmaceutical Science & Research, Durgapur, West Burdwan, West Bengal, 713212
6. Assistant Professor of Genex College of Pharmaceutical Science & Technology, Chinsurah, Hooghly, West Bengal, 712102
7. Assistant Professor of Eminent College of Pharmacy, Barasat, 700124
8. Assistant Professor of PGMS, Department of Pharmaceutics, Medinipur, 721253
9. Assistant Professor of Bharat Technology, Uluberia, Howrah, 711315
10. Delhi Institute of Pharmaceutical Sciences & Research, New Delhi

ABSTRACT

Objective: The aim of the study was to investigate the nephroprotective & hepatoprotective activity of methanolic extract of *Clitoria ternatea* in Cisplatin & CCl₄ induced in rats

Methods: Methanolic extract of aerial part of *Clitoria ternatea* plant was studied for its Nephroprotective & Hepatoprotective activity in animal experiment models. Nephrotoxicity was induced by Cystone 16 mg/kg b.w. Standard drug was taken Silymarin. Test drug were given methanolic extract *Clitoria ternatea* 500 mg/kg, 1000 mg/kg. Hepatotoxicity was induced by CCl₄. Standard drug was taken cisplatin 100 mg/kg. Test drugs were given extract of *Clitoria ternatea* 500 mg/kg & 1000 mg/kg as per b.w

Results: In Hepatoprotective activity positive control group was provided with CCl₄ and increased SGPT, SGOT, ALP level compare to negative control group whereas Test(2) group was provided with methanolic extract of *Clitoria ternatea* 1000 mg/kg decreased SGPT, SGOT, ALP level compare to standard group. In nephroprotective activity positive control group was provided with CCl₄ increased Urea and creatinine level where as Test(2) group are provided with methanolic extract of *Clitoria ternatea* 1000 mg/kg decreased urea and creatinine level

Conclusion: On evaluating biochemical parameters it was found that methanolic extract of *Clitoria ternatea* 1000 mg/kg showed hepatoprotective and nephroprotective activity in rats.

Keywords: SGPT, SGOT, ALP, Nephroprotective, Hepatoprotective

Article Info: Received 02 July 2019; Review Completed 15 Aug 2019; Accepted 20 Aug 2019; Available online 30 Aug 2019



Cite this article as:

Chandra S, Das A, Roy T, Bose P, Mukherjee L, Samanta J, Banerjee R, Bakuli R, Jana M, Mukhopadhyay D, Evaluation of Methanolic Extract of *Clitoria ternatea* Hepatoprotective & Nephroprotective Activity in Rats, Journal of Drug Delivery and Therapeutics. 2019; 9(4-A):313-319 <http://dx.doi.org/10.22270/jddt.v9i4-A.3478>

*Address for Correspondence:

Srikanta Chandra, Assistant Professor of Jakir Hossain Institute of Pharmacy, Department of Pharmacology, Raghunathganj, Murshidabad, West Bengal, 742225

Introduction:

The Liver is among the most complex and pivotal organs in the human body it lays below the diaphragm in the abdominal pelvic region of the abdomen. The Liver is a reddish brown organ with four lobes of unequal size and shape. It is both largest internal organ & largest gland in the human body. It is connected to two large blood vessels one

is called hepatic artery and another one is called portal vein. Its constituents about 2.5% of an adult's body weight. It produced bile, an alkaline compound which aids in digestion via the emulsification of lipids. Two major types of cells populate the liver lobes one is parenchyma and another one is non parenchyma cells. Sinusoidal endothelial cells, kuffer cells & hepatic stellate cells are some of the

non parenchyma cells that line hepatic sinusoid. Liver plays a pivotal role in regulating various physiological processes. It is also involved in several vital functions, such as metabolism, secretion and storage. It has great capacity to detoxicate toxic substances and synthesize useful principle. It helps in the maintenance, performance and regulating homeostasis of the body. It involved with almost all the biochemical pathways to growth, fight against disease, nutrient supply, energy provision and reproduction. It aids metabolism of carbohydrate, protein and fat detoxification, secretion of bile and storage of vitamins. Chemical that cause liver injury are called hepatotoxins. [1] Certain medicinal agents, when taken in overdose & sometimes even when introduced within therapeutic ranges, may injure the organ. Other chemical agents, such as those used in laboratories (eg. CCl₄, Paracetamol) and industries (e.g.: Lead, arsenic), natural chemicals (e.g.: microcystins, aflatoxins) and IFN, herbal remedies (*Cascara sagrada*, ephedra) can also induce for hepatotoxicity. Chemicals these agents can convert in chemically reactive metabolites in Liver, which have the ability to interconnect with cellular macromolecules namely protein, lipids & nucleic acids, leading to protein dysfunction, lipid peroxidation, DNA damage and oxidative stress. This damage of cellular function can dismiss in cell death & likely liver failure [2]. In the modern treatment strategy having some limitation. Silymarin that is associated with nausea, vomiting, headache. Hence we found a good rationale beyond probing for the Hepatoprotective and Nephroprotective activity in our pipeline drug that is *Clitoria ternatea*. [3]

Materials & Method:

Animal Husbandry & Statutory Approval

Healthy male & female rats (Wistar albino) age of 4-8 weeks were selected after physical and behavioral veterinary examination from Institutional Animal Ethics Committee of Gupta College of Technological Sciences, Asansol. The weight range was fall within $\pm 20\%$ of the mean body for each sex at the time of initiation of treatment. All experiments involving animals complies with the ethical standards of animal handling and approved by IAEC. All the selected animals were kept under acclimatization on the same day. The animal will be acclimatized for minimum 5 days before initiation of dosing. The rats were housed in standard polypropylene cages with stainless steel top grill in group of 6 rats per cage. Clean autoclaved paddy husk was used as bedding. The paddy husk was changed at least thrice in a week. The animals were kept in a clean environment with 12 hour light & 12 hour dark cycles. The air was conditioned at $22 \pm 3^\circ \text{C}$ & the relative humidity was maintained between 55-65% with 100% exhaust. Standard rat pellet feed was provided *ad libitum* throughout the study, except over night fasting prior to blood collection & was offered the feed immediately after completion of blood collection of all the animals. Drinking water was provided *ad libitum* in polypropylene bottles with a stainless steel sipper tube throughout study period. [4]

Collection of plant:

Leaves of *Clitoria ternatea* were collected from Rampurhat (W.B) India in August 2018 and were authenticated by the Head of Botany Department of Government College, Rampurhat, Birbhum, West Bengal

Authentication :

A herbarium sheet was prepared and it was send to head of Botany Department of Government College, Rampurhat,

Birbhum, West Bengal, for authentication no of study plant is Ref No: Rph/BOT/2018/42

Extraction :

Extraction may be defined as the treatment of the plant or animals tissues with solvent, where by the medicinally active constituents are dissolve (menstrum) and most of the inert matter remains undissolved (marc). The effective extraction of leaves depends largely on solubility and functional group consideration. [5]

Soxhlet Extraction:

Soxhlet extraction is used where small volume of hot menstrum is passed over the drug time and again to dissolve out the active constituents until the drug is exhausted. This process is known as soxhlation. The Soxhlet apparatus required for the hot percolation is made from a very high grade of glass and consists three parts (a) a flask in which the menstrum is boiled (b) an extracting chamber in which drug is filled, is fitted with the side tube and a siphon (c) a condenser. The drug to be extracted, in suitably comminuted form was unusual packed in a thimble made of filter paper which was then placed into the wider part of extractor. Thimble was used to prevent chocking of the lower part of extraction by drug particles. Menstrum was placed into the flask and boiled, the vapor was allowed to pass through the side tube to the condenser where they are condensed and fall on the packed drug, through which it extract out the active constituents. As the volume of menstrum in the extractor increase, the level of liquid in the siphoned out into flask. On further heating the menstrum vaporizes while the dissolve active constituents remains behind in the flask. The alternate filling the emptying of the body of the extractor goes on continuously till the drug was exhausted. Thus the same quantity of menstrum was made to Soxhlet repeatedly, about 14-15 times through the drug and the active constituents were collected in flask. This process is not suitable for the drugs containing thermo labile active constituents. [6]

Successive Solvent Extraction :

After the selection, collection and drying of leaf of *Clitoria ternate* Linn extraction was done. In pharmacy the solvent used for extraction purpose is known as menstrum and residue left after extracting the desired constituent is known as marc. The effective extraction of plant materials depends largely on solubility, and functional group consideration. The powder rhizomes were subjected to cold maceration and successive Soxhlet extraction using various solvent of increasing polarity namely petroleum ether, chloroform, acetone and methanol. Each time before extracting with next solvent powdered was dried in an air oven below 50°C . The extract was concentrated by distilling off the solvent and then evaporating to dryness on water-bath. [7]

Extractive Value:

The method determines the amount of active constituent in a given amount of medicinal plant material when extracted with solvents. It is employed for those plant materials for which no chemical or biological assay method exist. The extraction of any crude drug with a particular solvent yields a solution containing different phytoconstituents. The composition of these phytoconstituents depends on the nature of drug and solvent used. The use of a single solvent can be the means of providing preliminary information of the quality of a particular drug sample. [8]

Table 1: Extractive Values of leaves of *Clitoria ternatea* in different solvents

Solvent Used	Extractive value(%w/w)
Distilled water	12.46
Petroleum ether(60-80)	2.46
Chloroform	6.88
Ethyl acetate	9.84
Methanol	14.42

Preliminary Phytochemical Screening:

Preliminary tests were carried out for the presence or absence of phytoconstituents like Alkaloids, Carbohydrates, Flavonoids, Glycosides, Saponins, Sterols, Terpenes and Tannins in all the extracts by using the above four solvents individually. A description of methods adopted for performing the tests are summarized below.

Test for Alkaloids:

A portion of the extract was made acidic with dilute sulphuric acid. This portion was divided into two parts and was tested with the following precipitating reagents:

Mayer's Reagent:

1.36 gm of mercuric chloride dissolved in 60 ml of water, was added in a solution of 5 gm of potassium iodide in 20 ml of distilled water. They are mixed properly and volume was made upto 100 ml with distilled water. Buff colored precipitate was considered to be a positive test [9]

Dragendorff Reagent :

1 gm of bismuth sub nitrate in 20 ml of acetic acid was added to 20 gram of potassium iodide in 100 ml of water. Orange or brown colored precipitate was considered to be positive test[10]

Test for Carbohydrates:**Molish's Test:**

It was performed for the conformation of carbohydrates. 1 ml of 10% of acidic solution of α -naphthol was added to the extracts & mixed. Then 1 ml of concentrated sulphuric acid was carefully poured along the sides of the test tubes. The violet ring at the juncture of the two layers was considered to be positive test [11]

Test for Glycosides**Kedde Test (For Aglycone):**

The extract was evaporated to dryness and one drop of 90% alcohol and two drop of 2% 3,5 dinitrobenzoic acid in 90 % alcohol was added and the above mixtures was made alkaline with 20 % Sodium hydroxide to get purple color. Appearance of purple color showed the presence of free aglycone moiety [12]

Keller -Killani Test(For Sugar) :

To the dried extract 0.4 ml of glacial acetic acid containing a trace of Ferric chloride was added. To the mixture, 0.5 ml of concentrated H_2SO_4 was added. Presence of green blue colour in the upper acetic acid layer indicates the presence of sugar moiety[13]

Test for Flavonoids:

1.The extract was treated with magnesium (dust) and concentrated HCl. The appearance of pink tomato colour was indicative the presence of Flavonoids

Shinoda's Test:

2. 5 – 10 drops of dilute hydrochloric acid were added to 0.5 ml of extract. A small piece of magnesium was added to it. The appearance of pink, reddish pink or brown coloration was considered positive test. The appearance of yellow, orange red or brick color precipitate with lead acetate indicates the presence of Flavonoids[14]

3. 10 ml solution of the extract was hydrolyzed with dilute sulphuric acid. This was extracted with ether and divided into two portions. 1 ml dilute ammonia solution was added to one portion, a greenish yellow color conform presence of Flavonoids. To the other portion 1ml of dilute sodium bicarbonate solution was added, a pale yellow color conform the presence of Flavonoids

Test for Sterols and Terpene :**Benedict's Test:**

To 5 ml of Benedict's reagent few drops of extract was added and boiled in water bath for 5 mins. Appearance of green, yellow or orange – red precipitate indicates the presence of reducing sugar. [15]

Fehling's Test for Reducing Sugars:

To the equal volume of Fehling A and Fehling B mixture, 2 ml of extract was added and boiled for 5 minutes in water bath. Appearance of red precipitate indicates the presence of reducing sugar

Test for Saponins:**Foam Test :**

A small amount of dry extract was boiled with water and allowed to cool. It was then shaken vigorously for a minute. The formation of persistent honeycomb like froth was taken as positive results for Saponins [16]

Test for Tannins:

A small portion of extract was treated with 5% ferric chloride solution appearance of green to blue color is positive test for tannins

A creamy precipitate with lead acetate was considered positive test for tannins.

Table 2: Result of Preliminary Phytochemical Screening of Extracts for Various Phytoconstituents

Extracts	Alkaloid	Carbohydrates	Flavonoid	Glycosides	Red sugar	Saponins	Sterols	Terpenes	Tannins
Petroleum ether	--	--	--	++	--	--	++	++	--
Choloroform	+	--	--	+	++	-	+	--	+
Ethyl acetate	--	+	+++	+	--	+	--	--	--
Methanol	+++	+	+++	++	++	+	+	+++	--
Water	+++	+++	+++	++	++	--	--	--	++

+++ Prominently present ; ++ Moderately present , +Slightly present , -- Absent

Acute Toxicity & Gross Behavioral Studies:

Acute toxicity studies were carried out for methanolic extract using Acute Toxic Method as described in OECD (Organization of Economic Co-operation & Development) Guide Line No: 423. Animals were given increasing doses of 30, 100, 300, 600, & 1000 mg/kg p.o of the methanolic extract suspended in 2% tween -80 solution. The animals were observed continuously for 2 hours gross behavioral changes and intermittently once every 2 hours and finally at the end of 24 and 72 hours to note any toxic sign. [17]

Experimental Design:

Group I: Negative control (Normal Saline)

Group II: Positive control (CCl₄ 1ml/kg/day) 9 days

Group III : Standard (Silymarin 100 mg/kg/day + CCl₄ 1ml/kg/day) 9 days

Group IV: Test(1) (Methanolic extract of *Clitoria ternatea* 500 mg/kg/day + CCl₄ 1ml/kg/day) 9 days

Group V: Test(2) (Methanolic extract of *Clitoria ternatea* 1000 mg/kg/day + CCl₄ 1ml/kg/day) 9 days

Assessment of Liver Function :

The Hepatoprotective effect of extract was evaluated by the assay of Liver function biochemical parameters namely Serum glutamate pyruvate transaminase (SGPT), Serum glutamate oxaloacetate transaminase (SGOT), alkaline phosphates (ALP) & Total serum bilirubin (SB) according to the standard methods.

Histopathological Studies :

Formalin fixed tissues were subjected to graded dehydration in ascending strength of alcohol 70%, 80%, 90% and 100% and a subsequent wash with xylene. Following these the tissues were embedded in liquid paraffin to facilitate preparation of histopathological blocks which are essential for imparting strength to the tissue so that it can withstand the abrasive force of the blade while being sectioned. The 5 µm sections which were obtained by trimming with manual rotary microtome are subjected to staining by Harris' haematoxylin and counter stained by Eosin. The sections were viewed under trinocular microscope of different magnifications which were photographed by Motif software inbuilt in the systems. [22]

Statistical Analysis :

Results have been expressed as mean ± SEM. One way ANOVA have been employed for comparing majority of parameters. Post hoc tests were used for identification of groups having significant differences for one way ANOVA. Turkey's Multiple Range Test were used for comparisons. Whereas for two way anova Bonferroni test was used for the post hoc analysis. The significant groups were identified on the figure by designated alphabets. [23]

Nephroprotective Activity:

Nephrotoxicity can be defined as renal disease or dysfunction that arises as a direct or indirect result of exposure to medicines, and industrial or environment chemicals. Drug nephrotoxicity is therefore any renal dysfunction attributable to drugs. Drug nephropathies are not restricted to a single type of renal injury. Drugs target one or more discrete anatomical regions of the kidney and may affect only one cell type. The resulting insult to the kidney may result in a spectrum of nephropathies that are indistinguishable from those that do not have chemical

etiology. The nephron is the functional unit of the kidney and consists of a continuous tube of highly specialized heterogeneous cells, which show sub-specialization along the length of nephron and between them. It is the major organ of excretion and homeostasis for water soluble molecules because it is a metabolically active organ, it can concentrate certain substances actively. In addition, its cells have the potential to bioconvert chemicals and metabolically activate a variety of compounds. Since the kidney excretes many drugs, it is routinely exposed to high concentrations of these drugs or their metabolites or both. [24] Furthermore, the kidney has several features that allow nephrotoxins to accumulate. It is highly vascular, receiving about 25% of the resting cardiac output. The proximal renal tubule presents a large area for nephrotoxin binding and transport into the renal epithelium. Reabsorption of the glomerular filtrate progressively increases intraluminal nephrotoxin concentrations, while specific transport pathways in the kidney may engender site specific toxicity. Free radicals are highly reactive substances formed in the body as a result of metabolic processes. [25] Many of these molecular species are oxygen (and sometimes nitrogen) centered free radical and its non-radical products. The term reactive oxygen species (ROS) collectively denotes oxygen centered radicals (super oxide and hydroxyl radicals) as well as non-radical species derived from oxygen such as hydrogen 1 peroxide, singlet oxygen (O). The increased production of ROS seems to accompany most forms of tissue injury. Free radical can also react with DNA, proteins or lipids in the cell membrane and cause damage. The involvement of ROS in aging and in many chronic diseases has been considered. The defense provided by antioxidant system is crucial for the survival of organisms. Detoxification of ROS in the cell is provided by both enzymatic and non-enzymatic systems which constitute the antioxidant defense system. Many plants contain antioxidant compounds and these compounds protect cells against the damaging effects of reactive oxygen species (ROS) such as singlet oxygen, superoxide, peroxy radicals, hydroxyl radicals and peroxy nitrite. These compounds or antioxidant that can scavenge free radicals have vital role in improvement of diseased conditions. [26, 27]

Experimental Design :

Group(1) : Negative control (Normal Saline)

Group(2) : Positive control Cisplatin (16mg/kg b.w single dose i.p)

Group(3) : Standard Cystone (5ml/kg) + Cisplatin (16 mg/kg b.w single dose i.p)

Group (4): Test(1) Methanolic extract of *Clitoria ternatea* (500 mg/kg/day, po) + Cisplatin (16 mg/kg b.w single dose i.p)

Group (5) : Test(2) Methanolic extract of *Clitoria ternatea* (1000 mg/kg /day, po) + Cisplatin (16 mg/kg b.w single dose i.p)

Assessment of Renal Function:

The nephroprotective effect of extract was evaluated by the assay of kidney function biochemical parameters such as kidney weight as % of the total body weight, Blood urea level, Serum creatinine level and Histopathological evaluation of the kidney [28]

Statistical Analysis:

Results have been expressed as mean ± SEM. One way ANOVA have been employed for comparing majority of

parameters . Post hoc tests were used for identification of groups having significant differences for one way ANOVA . Turkey's Multiple Range Test were used for comparisons.

Whereas for two way anova Bonferroni's test was used for the post hoc analysis. The significant groups were identified on the figure by designated alphabets. [29 , 30, 31]

Table 3: Serum Biochemical Analysis Result of Hepatoprotective Activity

Group	Treatment region	SGPT(IU/L)	SGOT(IU/L)	ALP(IU/L)	Serum bilirubin(IU/L)
1	Normal saline	62.58±4.12	155.45±6.32	264.35±8.22	0.42±0.04
2	CCl ₄	240.58±13.54 ^a	490.68±15.58 ^a	534.82±19.43 ^a	1.42±0.32 ^a
3	Silymarin+ CCl ₄	73.42±5.45 ^{***}	172.43±6.75 ^{***}	283.76±10.41 ^{***}	0.45±0.04 ^{***}
4	MCT 500 mg/kg+ CCl ₄	115.45±7.56 ^{***}	226.43±8.68 ^{***}	324.96±12.47 ^{***}	0.62±0.05 ^{***}
5	MCT 1000 mg/kg+ CCl ₄	92.65±7.42 ^{***}	205.54±8.62 ^{***}	301.82±10.32 ^{***}	0.52±0.05 ^{***}

Values are expressed as mean ± SEM (n=6) and analyzed by using ANOVA followed by Bonferroni's multiple comparison test. Percentage change of protective effect compared with the CCl₄ treated control is expressed within brackets . P : ^a< 0.001 vs . vehicle control ,^{ns}> 0.05 , < 0.05 , < 0.01 , < 0.001 vs CCl₄ treated control




Fig 1: Histopathological evaluation of Hepatoprotective Activity

- (A) Group 1: Negative control (Normal saline)
 (B) Group 2: Positive control (CCl₄)
 (C) Group 3: Standard(Silymarin + CCl₄)
 (D) Group 4 : Test(1) (Methanolic extract *Clitoria ternatea* 500 mg/kg + CCl₄)
 (E) Group 5 : Test(2) (Methanolic extract of *Clitoria ternatea* 1000 mg/kg + CCl₄)

Table3: Serum biochemical analysis result for Nephroprotective activity

Group	Treatment Regimen	Urea(mg/dL)	Creatinine(mg/dL)
1	Normal Saline	38.76±3.76	1.06±0.40
2	Cisplatin	75.71±7.50 ^a	2.51±0.88 ^a
3	Cystone 5ml/kg + Cisplatin 16 mg/kg b.w	52.57± 4.56 ^{***}	1.36±0.62 [*]
4	<i>Clitoria ternatea</i> 500mg/kg +Cisplatin 16mg/kg i.p	68.74 ± 5.46 ^{***}	1.82±0.83 ^{ns}
5	<i>Clitoria ternatea</i> 1000 mg/kg/day + Cisplatin16mg/kg b.w .single dose i.p	57.43±7.34 ^{***}	1.55±0.42 [*]

Values are expressed as mean ± SEM (n = 6) and analyzed by using ANOVA followed by Bonferroni's multiple comparison test. Percentage change of protective effect compared with the Cisplatin treated control is expressed within brackets . p: ^a< 0.001 vs vehicle control ,^{ns}> 0.05 , * < 0.05 , ** < 0.01 , *** < 0.001 vs. Cisplatin treated control .




Fig 12 :Histopathological evaluation of Nephroprotective Activity

- (A) Group1 : Negative control(Normal saline)
 (B) Group2: Positive control (Cisplatin 5 ml)
 (C) Group3: Standard(Cystone 5 ml + Cisplatin 16 mg/kg)
 (D) Group4: Test(1) (Methanolic extract of *Clitoria ternatea* 500 mg/kg+ Cisplatin16 mg/kg)
 (E) Group 5: Test(2) (Methanolic extract of *Clitoria ternatea* 1000 mg/kg+ Cisplatin16 mg/kg)

DISCUSSION:

In the present study it was found that the methanolic extract *Clitoria ternatea* can modulate the nephrotoxicity induced by CCl_4 . The hepatotoxicity of CCl_4 has been reported to be due to the formation of the highly reactive trichloro(CCl_3) free radical, which alters function of endoplasmic reticulum and causes peroxidative degradation of lipid membrane of the adipose tissue leads to loss of metabolic enzyme located in the intracellular structures. Further it has been evident that several phytoconstituents have the ability to induce microsomal enzymes either by accelerating the excretion of CCl_4 or by inhibition of lipid peroxidation induced by CCl_4 . Phytoconstituents namely Flavonoids, Triterpenoids, Saponins and Alkaloids are known to possess Hepatoprotective activity.

In this study, the CCl_4 induced Liver damage was characterized by increased level of SGPT, SGOT, ALT and SB. Pre-treatment of animal with silymarin 100 mg/kg p.o. could reduce the level of SGPT, SGOT, ALP and SB. Similar results were obtained by pre-treatment of animal with methanolic extract of *Clitoria ternatea* (500 mg/kg & 1000 mg/kg p.o.) compared with the CCl_4 treated group. However the effectiveness of the extract the dose levels tested, was less compared to the standard Hepatoprotective drug used in the study, silymarin.

Silymarin which has very low toxicity and has been shown to possess a good safety profile. At high doses, a laxative effect is observed due to increased bile secretion and bile flow. Adverse of patients in a clinical trial. Serious adverse

effects, which are rare, include gastroenteritis associated with collapse and allergy. Thus combination therapy may up to some extent reduce these adverse effects of silymarin.

It is evident from the studies carried out by earlier researchers that the tuber of the plant contained higher phenolic and Flavonoids content and scavenging activities. The qualitative phytochemicals investigations carried out on the methanolic extract of *Clitoria ternatea* has significant Hepatoprotective activity. This may be probably due to the higher content of Flavonoids.

In the present study it was found that the tuber extract of *Clitoria ternatea* can modulate the nephrotoxicity induced by cisplatin. Cisplatin is a potent drug used in the management of wide range of cancer. However, the severe toxic side effects are the major limitation in its usage although the mechanism of cisplatin induced nephrotoxicity is exactly unknown. Several studies have suggested that the nephrotoxicity is induced by lipid peroxidation and free radicals. On the other hand many antioxidants have been shown to be protective against cisplatin induced nephrotoxicity. Also various free radical scavengers have been shown to be effective in protection against cisplatin induced nephrotoxicity & treatment with such agents provides significant protection against cisplatin induced acute renal failure. In this study, the cisplatin induced kidney damage was characterized by significant increase in serum creatinine ($p < 0.01$) and urea ($p < 0.001$) in comparison with the control group. Pre treatment of animal with cystone (5ml/kg p.o) for 6

consecutive days & a single dose of cisplatin (16mg/kg i.p) prevented the elevation of urea & serum creatinine as compared with cisplatin induced nephrotoxicity group. However, pre treatment of animal with methanolic extract of *Clitoria ternatea* (500mg/kg) for 6 consecutive days and a single dose of cisplatin did not cause a significant reduction of serum creatinine as compared with cisplatin induced nephrotoxicity group. The histopathological evaluation of the kidney preparations in treatment group also revealed a decreased induced tubular congestion, Tubular cast, Epithelial Desquamation, Glomerular congestion, blood vessel congestion & inflammatory cells.

CONCLUSION:

The methanolic extract of leaves of *Clitoria ternatea* 1000 mg/kg was found to possess Hepatoprotective & Nephroprotective activity. The activity was found to be less compared to the standard drugs used in the study. The studies were done using the crude lyophilized extract.

ACKNOWLEDGEMENT: Avik Das, Tathagata Roy, Preeta Bose they have a plethora of contribution on this scientific research.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial or not for profit sectors.

REFERENCES:

- Brown AC, Reitzenstein JE, Liu J, Jadus MR, Hepatoprotective activity of *Colocasia esculenta* in mice; International Journal of Pharmaceutical Science & Drug Research 2005; 19:767-71
- B.P. Kale, M.A. Kothekar, H.P. Tayade, J.B. Jaju, M. Mateenuddin; Effect of aqueous extract of *Azadirachta indica* leaves on Hepatotoxicity induced by antitubercular drugs in rats; Indian Journal of Pharmacology, 2003;35:177-180
- Marcinek K, Krejpcio Z; *Stevia rebaudiana* Bertonia chemical composition & functional properties; Acta Sci Pol Aliment, 2015 April 14(2):145-152
- Devmurari V.P, Jivani N.P; Hepatoprotective activity of methanolic & aqueous extract of *Azadirachata indica* leaves; International Journal of Pharm Tech Research, Volume:2, No:2; 2010,1037-1040
- Jorge Carlos Ruiz-Ruiz, Yolanda Beatriz Moguel-Ordonez, Maira Rubi Segura-Campos; Biological activity of *Stevia rebaudiana* Bertoni & their relationship to health; Critical Reviews in Food Science & Nutrition, Volume:57, No:12, 2017,2680-2690
- Robin S, Sunil K, Rana A.C, Nidhi S; Different models of hepatotoxicity and related liver diseases: a review; International research journal of pharmacy 2012. 3(7):86-95
- Ursula Wolwer-Rieck; The leaves of *Stevia rebaudiana*, their constituents & the analyses thereof: A review; Journal of Agricultural & Food Chemistry, 2012, 60(4):886-895
- Yadav P N, Pal A, Shanker K, Bawankule U D, Gupta K A, Darokar P M Khanuja S P S; Synergistic effect of silymarin and standardized extract of *phyllanthus amarus* against CCl4 induced hepatotoxicity of *rattus norvegicus*; Phytomedicine 2008;15:1053-1061
- Sonia Goswami, Rashmi Singhai; Activity of *Moringa oleifera* using successive solvent extraction technique; International Journal of Theoretical & Applied Sciences; 8(1):2016,23-27
- Prem Kumar Govindappa, Vidhi Gautam, Syamantak Mani Tripathi, Yash Pal Sahni, Hallur Lakshmana Shetty Raghavendra; Effect of *Withania somnifera* on gentamicin induced renal lesions in rats; Revista Brasileira de Farmacognosia, Volume:29, No:2; 2019
- Pal S, Bhattacharjee A, Mukherjee S, Bhattacharya K, Khowala S; Antioxidant and Hepatoprotective activity of ethanolic extract of *Alocasia indica* tuber; AJPCT 2014;2(2):191-208
- Abraham Z, Bhakuni DS, Garg HS, Goel AK, Mehrota BN, Patnaik GK; Screening of Indian plants for biological activity; Exp Biol Indian J 1986;24(1):48-68
- Chattopadhyay RR, Sarkar SK, Ganguly S, Banerjee RN, Basu TK, Mukherjee A; Hepatoprotective activity of *Azadirachta indica* leaves on paracetamol induced hepatic damage in rats; Indian Journal of Experimental Biology; 1992,30(8):738-40
- Harborne JB, Chapman and Hall; Phytochemical Methods. London 1983;2:277-9
- Iwashina T, Konishi T, Takayama A, Fukada M, Ootani S; Isolation and identification flavonoids in the leaves of taro: Ann Tsukuba Botanical Garden 1999; 18:71-4
- Hung AS, Tanudjaja LS; Application of anion exchange high performance liquid chromatography in determining oxalates in taro (*Colocasia esculenta*) corms; J Agric Food Chem 1992; 40:2123-6
- Zhang Q, Yang H, Li Y, Liu H, Jia X; Toxicological evaluation of ethanolic extract from *Stevia rebaudiana* Bertonia leaves: Genotoxicity & Subchronic oral toxicity; Regulatory Toxicology & Pharmacology, Volume:86, June 2017, Pages:253-259
- N.S. Baligar, R.H. Aladakatti, Mukhtar Ahmed, M.B. Hiremath; Hepatoprotective activity of the neem-based constituent azadirachtin-A in carbon tetrachloride intoxicated Wistar rats; Canadian Journal of Physiology & Pharmacology, Volume:92, 2014, 267-277
- Ismet Ara Jahan, Khondoker Shahin Ahmed, Zakia Sultana, M. Hemayet Hossain, Pijush Kanti Biswas, Katrun Nada; Antioxidant activity of methanolic extract of *Withania somnifera* in Mice Oriental Pharmacy & Experimental Medicine; Volume:18, December 2018, 299-307
- A.B. Falowo, V. Muchenje, A. Hugo, O.A. Aiyegoro, P.O. Fayemi; Antioxidant activities of *Moringa oleifera* & *Bidens pilosa* L. leaf extracts & their effects on oxidative stability of ground raw beef during refrigeration storage; CyTA -Journal of Food 15:2, 2016, 249-256
- Shah BN, Nayak BS, Bhatt SP, Jalalpure SS, Sheth AK; The antioxidant activity of leaves of *Colocasia esculenta*; Pharmacognosy Journal 2007;15:3-4
- Vusumzi Pakade, Ewa Cukrowska, Luke Chimuka; Comparison of antioxidant activity of *Moringa oleifera* & selected vegetables in South Africa; South African Journal of Science, Volume:109, 2013, 95-99
- Jeyanthi T, Subramanian P; Nephroprotective effect of *Withania somnifera*: a dose-dependent study, Renal Failure 31:9, 2009; 814-821
- Mugdha Rajeeva Padhya, Sangita Devrao Jogdand, Jagruti Bhattacharjee; Evaluation & comparison of nephroprotective effect of *Hemidesmus indicus* Linn. & *Withania somnifera* Linn. on gentamicin induced nephrotoxicity in rats; International Journal of Basic & Clinical Pharmacology, April 2018, Volume:7, 691-695
- Vimlesh Kaushwaha, Monica Sharma, Pinki Vishwakarma, Manish Saini, Kuldeep Saxena; Biochemical assessment of nephroprotective & nephrocurative activity of *withania somnifera* on gentamicin induced nephrotoxicity in experimental rats; International Journal of Research in Medical Sciences, 2016 jan;4(1):298-302
- Shimeda Y, Hirota Y, Akimoto Y, Shindou K, Yoshio I, Nishihori T, Tanaka K; Protective effects of Capsaicin against Cisplatin-Induced Nephrotoxicity in Rats; Biol Pharm Sci 2005;28(9):1635-8
- Preethi CK, Kuttan R; Hepato and Reno Protective action of *Calendula officinalis* Linn. Flower extract; Ind J Exp Biol 2009;47:163-8
- Kalariya M, Parmar S, Sheth N; Neuropharmacological activity of hydroalcoholic extract of leaves of *Stevia rebaudiana*; Biol Pharm 2010;48:1207-12
- Syed Safiullah Ghori, Tahreen Shazia Siddiqua, Anees Fathima; Nephroprotective effect of *Ficus Dalhousiae* Miq leaf methanolic extract in Albino Wistar Rats; International Journal of Pharma Research & Health Sciences. 2016
- N. Kamkaen, J.M. Wilkinson; The antioxidant activity of *Clitoria ternatea* flower petal extracts & eye gel; Wiley Online Library, Volume:23, 23rd Oct, 2009
- Chong FC, Gwee XF; Ultrasonic extraction of anthocyanin from *Clitoria ternatea* flowers using response surface methodology; Nat Prod Res, 2015; 29(15):1485-7