


Available online on 15.02.2020 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

Antimicrobial and Antioxidant Activities of Flavonoids Extracted from *Pistacia lentiscus L.*, Leaves

Bakli Sabrina^{1,2*}, Harzallah Daoud¹, Zerroug Amina¹, Sadrati Nouari¹, Bouguerra Asma¹, Gaamoune Soufiane^{1,3} and Naili Oumaima¹

¹: Laboratory of Applied Microbiology, Faculty of Natural Sciences and Life, University Ferhat Abbas Setif1, Setif 19000, Algeria.

²: Department of Natural Sciences and Life, University of Abdel Hafid Boussouf, Mila 43000, Algeria.

³: National Institute of Agriculture Research, Setif 19000, Algeria.

ABSTRACT

Increasingly, the flavonoids becoming the subject of biomedical researches, and many groups have isolated and identified the structures of flavonoids possessing important biological activities. This assess investigates the antimicrobial fight and the antioxidant strengths of flavonoids extracted from *Pistacia lentiscus L.* leaves. The extract was first analyzed for its phenolic content, then the flavonoids were tested for their antimicrobial capacities by the method of diffusion against fourteen bacteria, two fungi and one yeast. The minimum inhibitory concentration (MIC) and the minimum bactericidal concentration (MBC) of flavonoids were determined using the dilution method on solid medium, however, the antioxidant activity was evaluated through the ability of the extract to scavenge DPPH (1,1-diphenyl-2-picrylhydrazyl) radicals and the reducing power assay. The results indicate that flavonoids extract contain other phenolics groups. On the other side, the Antibacterial tests exhibited different activities depending on the strains. Unlike, the MIC and the MBC showed that the flavonoids extract showed a best activity against *Candida albicans* and *Vibrio cholerae*. Concerning antioxidant activity, the results indicate a moderate capacity of the flavonoid extract. These preliminary results may justify the use of this plant in the treatment of some diseases and may be exploited for therapeutic purposes.

Keywords: Antimicrobial, Antioxidant, *Pistacia lentiscus*, flavonoids, MIC, MBC.

Article Info: Received 26 Nov 2019; Review Completed 18 Jan 2020; Accepted 27 Jan 2020; Available online 15 Feb 2020



Cite this article as:

Bakli S, Harzallah D, Zerroug A, Sadrati N, Bouguerra A, Gaamoune S, Naili O, Antimicrobial and Antioxidant Activities of Flavonoids Extracted from *Pistacia lentiscus L.*, Leaves, Journal of Drug Delivery and Therapeutics. 2020; 10(1-s):82-89 <http://dx.doi.org/10.22270/jddt.v10i1-s.3895>

*Address for Correspondence:

Bakli Sabrina, Laboratory of Applied Microbiology, Faculty of Nature and Life Sciences, University Ferhat Abbas Setif1, Setif 19000, Algeria.

INTRODUCTION

P. lentiscus leaves contain differing types of secondary metabolites. Among these, the most abundant compounds in *P. lentiscus* leaves are reportedly flavonoids that have displayed a powerful antioxidant capacity as well as hepatoprotective, anti-inflammatory, anticancer effects, antimicrobial, antiviral, anticholinesterase, anti-inflammatory, antinociceptive, antidiabetic, antitumor, antihyperlipidemic, antiatherosclerotic and hepatoprotective activities and also their beneficial effects in gastrointestinal disorders [1]. Thus, the characterization of bioactive compounds from *P. lentiscus* leaves extracts has relevance for developing nutraceuticals and dietary supplements [2].

Nowadays, the development of resistance of a pathogen to a large number of commonly used antibiotics encourages the search for new antimicrobial agents to fight infections and

overcome the resistance and side effects of antimicrobial agent currently available [3]. Due to the alarming increase in the incidence of new and re-emerging infectious diseases, there is a continuing and urgent need to find new antimicrobial compounds with different chemical structures and new mechanisms of action. Another major concern is the development of antibiotic resistance in current clinical use [4]. Flavonoids are increasingly the subject of anti-infectious research. Several flavonoids have antibacterial and antifungal activities [5].

The aim of this study is to evaluate the antioxidant and antimicrobial activities of flavonoids extract obtained from leaves of *P. lentiscus*, in order to find new molecules able to reduce the use of synthetic medicines.

MATERIAL AND METHODS

Plant material

The leaves of *P. lentiscus* were harvested in July 2014, in Bni Ourtilan – Setif– Algeria. The *P. lentiscus* leaves were air-dried in the shade at room temperature. After drying, the plant material was ground to a fine powder using an electric mill.

Flavonoids extraction

The powder was taken up three times with 70 % ethanol (raw material: solvent ratio was 1: 10) for 90 minutes at 100 °C. The extracts were combined and concentrated under vacuum to collect the aqueous residue (10 mL), which was extracted with hexane and chloroform, then acidified with 20 % H₂SO₄ (pH = 5) and finally extracted with ethyl acetate. The appearance of an interphase precipitate was observed during extraction with ethyl acetate. The ethyl acetate fraction was taken as a flavonoids fraction for our experiment [6].

Determination of total flavonoids contents

The flavonoids contents in the extract was estimated by the Aluminium chloride solution according to the method described by Mimica-Dukić (1992) in Maksimović *et al.* [7]. Briefly, 2 mL of the methanol solution of the extract was added to 1 mL of AlCl₃ (133 mg crystalline aluminium chloride and 400 mg crystalline sodium acetate were dissolved in 100 mL of distilled water) in methanol. After 10 minutes, the absorbance was determined at 430 nm. Results were expressed as mg equivalent rutin per gram of extract (mg ER/GE).

Antioxidant activities

DPPH radical scavenging activity

The DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging activity of the flavonoids extract was measured according to the procedure described by Sreenivasan *et al.* [8]. Briefly; 50 µL of the extract at different concentrations were added to 5 mL (0,04 %) of DPPH methanolic solution. The decrease in absorbance was determined at 517 nm, after incubation for 30 minutes at laboratory temperature in the dark.

The antiradical activity was expressed as IC₅₀ (micrograms per millilitre) and the antiradical dose required to cause a 50 % of inhibition. A lower IC₅₀ value corresponds to a better antioxidant activity. The ability to scavenge DPPH radical was calculated by employing the subsequent equation:

$$\text{DPPH scavenging effect (\%)} = [(A_0 - A_1) / A_0] \times 100$$

Where A₀ is the absorbance of the control at 30 minutes and A₁ is the absorbance of the sample at 30 minutes. BHT was used as a standard and samples were analyzed in triplicate.

Ferric reducing power (FRP)

The reducing power of the extract was measured with a method described by Kaur *et al.* [9]. One millilitre of the extract at different concentrations was mixed with 2.5 mL of phosphate buffer solution (0.2 M, pH 6.6) and 2.5 mL of potassium ferricyanide (1 %). The mixture was incubated for 20 minutes at 50 °C. After cooling, 2.5 mL of 10 % trichloroacetic acid was added and the mixture was centrifuged at 3000 rpm for 10 minutes. 2.5 mL of supernatant was mixed with 2.5 mL of distilled water and 0.5 mL of ferric chloride (0.1 %). The absorbance of the mixture thus obtained is measured at 700 nm against a blank.

Higher absorbance indicates higher reducing power. EC₅₀ value (µg extract/mL) is the effective concentration at which the absorbance was 0.5 for reducing power, obtained by interpolation from linear regression analysis. BHT was used as standards.

Antimicrobial activity test

Microorganisms

Evaluation of the antimicrobial activity of extract was performed against fourteen strains of pathogenic bacteria, obtained from the American Type Culture Collection, namely: *Staphylococcus aureus* ATCC 25923, *methicillin resistant Staphylococcus aureus (MRSA)* ATCC 43300, *Bacillus cereus* ATCC 10876, *Enterococcus faecalis* ATCC 49452, *Citrobacter freundii* ATCC 8090, *Escherichia coli* ATCC 25922, *Klebsiella pneumoniae* ATCC 700603, *Salmonella typhi* ATCC 14028, *Salmonella enterica* ATCC43972, *Salmonella typhimurium* ATCC 13311, *Pseudomonas aeruginosa* ATCC 27853, *Proteus mirabilis* ATCC 35659, *Vibrio cholerae* ATCC 14035 and *Listeria innocua* CLIP 74915 from Collection *Listeria* Pasteur Institute.

Two fungi: *Aspergillus niger* 2CA936, *Aspergillus flavus* NRRL 3357 and one yeast: *Candida albicans* ATCC1024. Sabouraud Dextrose agar and Mueller–Hinton agar were used for fungi and bacteria, respectively.

Screening for antibacterial activity

The screening of antibacterial activity of flavonoids extract of *P. lentiscus* was carried out with agar disc diffusion method [10].

The extract was dissolved in dimethylsulfoxide (DMSO) to a final concentration of 200 mg/mL and sterilized by filtration through a 0.22 µm membrane filter. The bacteria inoculum was prepared by suspending colonies during 24 hours culture. The cell density of each inoculum was adjusted at 10⁸ CFU/mL suspension with a spectrophotometer (DO=0.08-0.1 / λ = 600 nm) [11]. The discs Watt-man paper discs N°3 of 6 mm diameter were each impregnated with 10 µL and 20 µL of extract equal to 2mg and 4mg by disc, respectively. The plates were held for 3 hours at 4 °C for diffusion of extract into the agar [12] [13] and then incubated at 37 °C for 24 hours. Amoxicillin, Ciprofloxacin, Teicoplanin, Doxycyclin, Pefloxacin, Optochin were used as standards and dimethylsulfoxide as a control.

The activity is determined by the measurement of the inhibitory zone diameter in mm. The antibacterial activity is considered starting from a diameter of 6 mm or higher, and is classified as follows [14]:

- ✓ Very sensitive: diameter ≥ 20 mm;
- ✓ Sensitive enough: diameter between 15-19 mm;
- ✓ Sensitive: diameter between 09-14 mm;
- ✓ Resistant: diameter ≤ 08 mm.

Screening for Antifungal activity

The antifungal activity was tested by disc diffusion method [15]. The Sabouraud plates were inoculated with each fungal culture.

The spore suspension was prepared in saline solution (0.9 % NaCl), adjusted to a concentration of 10⁶ spores/mL [15] [16] corresponding to 0.15 to 0.17 absorption at 530 nm [15]. One hundred microliter of suspension was placed over agar in Petri dishes and dispersed using a sterile swab. Then, the sterile paper discs (6 mm diameter) were placed on agar to load 10 µL and 20 µL equal to 2 mg and 4mg by disc, respectively. And the plates were held for 3 h at 4 °C for diffusion of extracts into the agar and then incubated at 37 °C

for 48 hours for *C. albicans* and 72 hours at 28 °C for the fungi.

Nystatin, clotrimazon and amphotericin were used as standards and dimethylsulfoxide as a control.

Determination of Bactericidal and Fungicidal Activity

Minimum inhibition concentration (MIC)

The MIC of the extract was determined using agar dilution method that has been already described by [17]. One milliliter of different concentrations of extract was added to 14 mL of nutrient agar to make the final concentration ranging from 10 mg/mL to 0.05 mg/mL. Standardized inocula approximately containing 1×10^4 CFU/mL [18] were spotted on solidified plates containing various concentrations of the extract. The lowest concentration of extract inhibiting any visible bacterial or fungal growth after an incubation period of 24 h at 37 °C and 48 h at 28 °C, respectively, was taken as the minimum inhibitory concentration [17][19].

Minimum Bactericidal Concentration (MBC) and Minimal Fungicidal Concentration (MFC)

Samples were taken from the nutrient agar plates that showed no visible growth after 24 hours incubation and sub cultured into tubes containing nutrient broth. The least concentration that did not produce growth after 24 hours was regarded as the minimal bactericidal concentration [17].

Statistical analysis

Experiments were carried out in triplicate and expressed as the mean \pm standard deviation, data were analysed and compared using the one-way ANOVA and Tukey Multiple Comparison with 95 % confidence limits ($P < 0.05$), using Graphpad prism 5 Demo Software.

RESULTS AND DISCUSSION

Flavonoids extracted from the leaves of *P. lentiscus* reach the yield of 3.91 % with 278.507 ± 0.377 mg ER/GE of total flavonoids.

Antioxidant activity

The isolated compounds were screened for antioxidant activity. The results of DPPH test were showed in figure 1.




Figure 1: Scavenging activity on the DPPH radical of flavonoids extracted from leaves of *P. Lentiscus*

The IC_{50} of flavonoids extract was $455 \pm 0,048$ µg/mL*** against $4,31 \pm 0,79$ µg/mL for BHT.

Studies have recently shown that flavonoids have an antioxidant effect on human health. These compounds

scavenge free radicals and chelate metal thus avoiding the Fenton reaction to occur [20]. The role of antioxidants in the inhibition of oxidant processes occurring in living organisms consists of scavenging free radicals and quenching singlet oxygen, disconnection of radical reactions, chelate metals which catalyze the oxidation process, inhibition of certain enzymes (eg., oxidases). Flavonoids are active in these processes [21]. However the weakness of our extract due to the presence of other molecules which affect the ability of scavenging in terms of hydrogen donating ability.

In addition, it is known that only flavonoids having a certain structure and in particular a hydroxyl position in the molecule can act as proton donors and show a radical scavenging activity [22][23].

The reducing capacity of a compound can be considered as an important indicator of its antioxidant activity [24]. Their absorption values increased with increase in concentration, showing that as the concentration of the extract was increased their ability to reduce Fe^{3+} to Fe^{2+} was also increased. Figure 2 shows the dose-dependent curve for the reducing power of the extract:




Figure 2: Reducing power of flavonoids extracted from the leaves of *P. lentiscus*

An increase in absorbance corresponds to an increase in the reducing power of the extract tested. Flavonoids showed increased ferric reducing power with the increasing concentration with EC_{50} correspond to 15.0 ± 0.001 µg/mL against 16.0 ± 0.001 µg/mL of BHT. The extract expressed electron donating activity and its power was equal to BHT.

High reducing power of flavonoids extract suggested its remarkable potency to donate electrons to reactive free radicals, thus, converting them into more stable non-reactive species and finally terminate the free radical chain reaction [25]. It was confirmed that the hydroxyl groups at C-3' and C-4' of the B-ring to be more active in reducing iron concentration [26]; this may explain our results.

The study of correlation between radical scavenging power and the capacity of metal ions chelation seems interesting. Firstly because oxidative stress is the cause of formation of free radicals through metals transition (iron, copper) via Fenton reaction [20]. Secondly, the evaluation of the plant antioxidant effect is made by combining the results of at least two *in vitro* assays [27].

Antimicrobial activity

The results obtained from antimicrobial assay are presented in Table 1, 2 and 3:

Table 1: Antibacterial activities of flavonoids, pure phenolic compounds, standards and control

Bacterial strains	Zone of inhibition (mm)													
	<i>S. aureus</i>	<i>SARM</i>	<i>L. innocua</i>	<i>B. cereus</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>S. typhi</i>	<i>S. enterica</i>	<i>S. typhimurium</i>	<i>C. freundii</i>	<i>P. aeruginosa</i>	<i>P. mirabilis</i>	<i>V. cholerae</i>	<i>E. faecalis</i>
Flavonoids (200 mg/mL)														
10µL	12,7±0,47 ^a	9,0±0,0 ^a	-/-	8,0±0,00 ^a	-/-	-/-	9,0±0,0 ^a	-/-	-/-	-/-	9,0±0,0 ^a	7,3±0,47 ^a	10,3±0,47 ^a	-/-
20µL	18,8±0,24 ^c	13,0±0,41 ^b	8,3±0,47 ^a	14,7±0,47 ^c	8,8±0,24 ^a	-/-	11,5±0,41 ^b	8,7±0,47 ^a	-/-	-/-	14,0±0,00 ^e	13,0±0,0 ^b	12,7±0,47 ^b	8,3±0,47 ^a
Pure phenolic compounds														
Tannic acid														
10µL	13,8±0,3 ^a	14,0±0,0 ^b	-/-	10,0±0,6 ^b	-/-	-/-	18,0±0,0 ^c	12,0±0,0 ^b	-/-	-/-	12,0±0,0 ^c	14,0±0,0 ^b	15,5±0,7 ^c	8,0±0,0 ^a
20µL	17,3±0,6 ^b	18,0±0,0 ^c	11,3±0,6 ^b	16,5±0,5 ^d	12,0±0,0 ^b	-/-	20,3±0,6 ^d	15,7±0,6 ^e	-/-	-/-	13,0±0,0 ^d	18,2±0,8 ^c	22,0±0,0 ^d	12,3±0,6 ^b
Gallic acid														
10µL	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
20µL	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	8,0±0,0 ^a	11,0±0,0 ^a	-/-
Quercetin														
10µL	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
20µL	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
Standards														
Amoxicillin	30,0±0,0 ^d	9,0±0,0 ^a	23,0±0,0 ^d	-/-	12,0±0,0 ^b	-/-	22,0±0,0 ^e	23,0±0,0 ^f	-/-	-/-	-/-	39,67±0,47 ^f	22,0±0,0 ^d	22,0±0,0 ^d
Ciprofloxacin	34,7±0,47 ^{ef}	35,0±0,0 ^e	32,0±0,0 ^e	37,3±0,47 ^g	46,0±0,0 ^e	32,0±0,0	32,0±0,0 ^g	26,7±0,47 ^g	45,3±0,47	51,3±0,47	37,7±0,47 ^g	33,67±0,47 ^e	49,0±0,82 ^g	31,7±0,47 ^f
Teicoplanin	19,7±0,47 ^c	20,0±0,0 ^d	21,0±0,0 ^c	15,0±0,0 ^c	-/-	-/-	22,3±0,47 ^e	20,0±0,0 ^d	-/-	16,8±0,62	-/-	25,0±0,0 ^d	26,0±0,0 ^e	20,0±0,0 ^c
Doxycycline	36,0±0,0 ^f	39,0±1,0 ^f	34,0±0,0 ^f	26,0±0,0 ^e	30,0±0,0 ^c	10,0±0,0	50,0±0,0 ^h	30,0±0,0 ^h	25,0±0,0	28,0±0,0	10,0±0,0 ^b	44,0±0,0 ^g	44,0±0,0 ^f	37,0±0,0 ^g
Pefloxacin	33,7±0,6 ^e	35,5±0,5 ^e	-/-	30,5±0,7 ^f	44,0±0,0 ^d	26,0±0,0	25,0±0,0 ^f	22,0±0,0 ^e	40,0±0,0	46,0±0,0	22,0±0,0 ^f	40,0±0,0 ^f	44,0±0,0 ^f	30,0±0,0 ^e
Optochin	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
Control	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition	No inhibition

-/-: No zone of inhibition

In the same column, means followed by the same letters are not significantly different (p < 0.05)

Table 2: Antifungal activities of flavonoids, pure phenolic compounds, standards and control

Fungal strains	Zone of inhibition (mm)		
	<i>Aspergillus flavus</i> NRRL3357	<i>Aspergillus niger</i> 2CA936	<i>Candida albicans</i> ATCC1024
Flavonoids (200 mg/mL)			
10µL	-/-	-/-	12,67±0,47 ^b
20µL	-/-	-/-	14,0±0,0 ^b
Purphenolic compounds			
Tannic acid			
10µL	11,3± 0,6 ^a	-/-	18,8± 1,0 ^d
20µL	12,9± 0,3 ^b	-/-	25,0±0,0 ^e
Galic acid			
10µL	-/-	-/-	-/-
20µL	-/-	-/-	-/-
Quercetin			
10µL	-/-	-/-	-/-
20µL	-/-	-/-	-/-
Standards			
Amphotericin	16,67±0,29 ^d	17,17±0,29 ^c	15,67±0,29 ^c
Clotrimazon	23,50±,50 ^e	15,67±0,29 ^b	44,67±0,58 ^f
Nystatin	15,17±0,29 ^c	9,17±0,29 ^a	9,17±0,29 ^a
Control	No inhibition	No inhibition	No inhibition

-/-: No zone of inhibition;

In the same column, means followed by the same letters are not significantly different (p < 0.05)

Table 3: MIC and MBC of flavonoids and pure phenolic compounds against the tested microbial strains

Strains	MIC et MBC/MFC (mg/mL)															Fungal strains	
	Bacterial strains															<i>C. albicans</i>	<i>A. flavus</i>
	<i>S. aureus</i>	MRSA	<i>B. cereus</i>	<i>L. innocua</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>S. typhi</i>	<i>S. enterica</i>	<i>S. Typhimurium</i>	<i>C. freundii</i>	<i>P. aeruginosa</i>	<i>P. mirabilis</i>	<i>V. cholerae</i>	<i>E. faecalis</i>			
Flavonoids																	
MIC	2	0,5	1	5	4	NT	0,6	5	NT	NT	3	3	0,3	5	0,1	NT	
MBC/MFC	3	1	6	5	5	NT	3	6	NT	NT	5	3	0,5	5	5	NT	
Pure phenolic compounds																	
Tannic acid																	
MIC	0,4	0,3	0,3	1	> 10	NT	<50 ug	1	NT	NT	0,9	0,1	0,1	1	0,1	> 10	
MBC/MFC	0,4	0,3	0,6	2	> 10	NT	3	8	NT	NT	0,9	0,3	0,1	5	5	> 10	
Galic acid																	
MIC	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	10	8	NT	NT	NT	
MBC/MFC	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	10	8	NT	NT	NT	
Quercetin																	
MIC	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
MBC/MFC	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Antibiotic standards																	
Amoxicillin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Ciprofloxacin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Teicoplanin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Doxycyclin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Pefloxacin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Optochin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Antifongic standards																	
Amphotericin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Clotrimazon	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	
Nystatin	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT	

NT : Not Tested.

The first table (**table 1**) indicates that the flavonoids extract is weak to moderate inhibitory activities against the tested bacteria even zero against some strains. The flavonoids at 4 mg/disc showed variable inhibitory activity against all bacteria (Gram-positive and Gram-negative) with inhibition zone diameters ranging from 8.33–18.83 mm. An extract is considered active when it reveals a zone of inhibition greater than or equal to 9 mm [14]. Furthermore, the assayed extract indicates no activity against tested fungi *A. niger* and *A. flavus* (**Table 2**), but showed an important antifungal activity against *C. albicans* with inhibition zone of 14.0±0.0 mm. This inhibition was low compared to Amphotericin.

Antimicrobial activity of pure phenolic compounds showed that tannic acid has more impact on *V. cholerae* (22.0±0.0 mm) (**Table 1**). Galic acid has showed minimum zone of inhibition on *P. mirabilis* (8.0±0.0 mm) and *V. Cholera* (11.0±0.0 mm). On the other hand tannic acid showed significant activity against *C. albicans* (25.0±0.0 mm) compared to standards nystatin (9.17±0.29mm) and amphotericin (15.67±0.29mm) and low activity against *A. flavus* (12.9±0.3 mm) (**Table 2**) while quercetin shows no inhibitory effect.

Among the six different types of antibiotics used in the study, Ciprofloxacin has a broad spectrum of activity on all species of human pathogenic bacteria and the widest zone of inhibition was observed against *C. freundii* (51.33±0.47 mm) (**Table 1**). Whereas the minimum zone of inhibition was exhibited in Amoxicillin (9.0±0.0 mm) against MRSA. Optochin has no antibacterial activity on the selected test organisms.

The MICs results are not in good agreement with those of the antibiogram for all tested species (**Table 3**); since the smallest MIC was observed with *C. albicans* (0.1 mg/mL) and the latter has small zone of inhibition (14.0±0.0 mm), when *S. aureus* showed a high MIC (2mg/mL) and the largest zone of inhibition (18.83±0.24 mm).

These results show that the flavonoids extract has a lower value of MIC 0.1 mg/mL on the fungal strain *C. albicans*, also showed a remarkable activity on *V. Cholera* with value of MBC 0.3 mg/mL and the best activity were obtained with tannic acid with MIC value of <50ug /mL against *S. typhi* and MBC 0.1 mg/mL against *V. Cholera*.

In this study, we find out that the extract could give higher inhibition to Gram-positive bacteria compared to Gram-negative bacteria (**Table 1**). These results can be explain that Gram-negative microorganisms are typically more resistant to antimicrobial agents than Gram-positive bacteria. This has long been explained by the presence of an outer-membrane permeability barrier in Gram-negative bacteria, which limits access of the antimicrobial agents to their targets in the bacterial cells [28] [29]. In addition, the periplasm contains enzymes that destroy foreign molecules introduced from the outside [30].

Antimicrobial activity that is observed in this study is due to the presence of flavonoids compounds in this extract because flavonoids are known for their antimicrobial activity against a wide range of microorganisms [31] [32] and they are multiple cellular targets and may apply to different components and functions in the bacterial cell [33] [34]. Extracts of various medicinal plants containing phenolics and flavonoids have been previously reported to possess antimicrobial activity [35] [36]. The presence of these compounds might contribute to antimicrobial activity of *P. lentiscus leaves* since **Cushnie et Lamb** [37] reported that flavonoids (kaempferol, myricetin, naringin, quercetin and rutin) have antimicrobial activity against human pathogenic

microorganisms with some mechanisms of action such as inhibition of nucleic acid synthesis, cytoplasmic membrane function by perforation mechanism and energy metabolism [35] [37] [38]. The antimicrobial activity of the flavonoids extract of *P. lentiscus leaves* might be due to one of the mechanisms of action mentioned above.

The results of the present study are quite encouraging as the flavonoids extract exhibited antimicrobial activity against most of the pathogens. However, the antimicrobial activity varies widely, depending on the microorganism.

CONCLUSION

Recently there is a huge interest in the use of plant material as an alternative method to control pathogenic microorganism and antioxidant agents.

This study shows that flavonoids extract of *P. lentiscus leaves* was potent as antimicrobial substances and powerful chelating capacity. It is suggested that could be considered as source of antimicrobial and antioxidant agent which might be applied in pharmaceutical and cosmetic products.

Further studies are needed to purify and investigate the molecular composition of this natural product which constitutes an alternative of the synthetic medicines.

REFERENCES

- [1] Bozorgi M, Memariani Z, Mobli M, Salehi Surmaghi MH, Shams-Ardekani MR, Rahimi R, Five *Pistacia* species (*P. vera*, *P. atlantica*, *P. terebinthus*, *P. khinjuk*, and *P. lentiscus*): A Review of Their Traditional Uses, Phytochemistry, and Pharmacology, *The Scientific World Journal*, 2013; 1-32.
- [2] Rodríguez-Pérez C, Quirantes-Piné R, Amessis-Ouchemoukh N, Madani K, Segura-Carretero A, Fernández-Gutierrez A, A metabolite-profiling approach allows the identification of new compounds from *Pistacia lentiscus leaves*, *Journal of Pharmaceutical and Biomedical Analysis*, 2013; 77:167-174.
- [3] Ali-Shtayah MS, YaghmourReem M-R, Faidi YR, Salem k, Al-Nuri MA, Antimicrobial activity of 20 plants used in folkloric medicine in the Palestinian area, *Journal of Ethnopharmacology*, 1998; 60:265-271.
- [4] Rojas R, Bustamante B, Bauer J, Fernández I, Albán J, Lock O, Antimicrobial activity of selected Peruvian medicinal plants, *Journal of Ethnopharmacology*, 2003; 88:199-204.
- [5] Jayshree N, Narayanan N, Sriram L, Antibacterial, antifungal and antimycobacterial studies on some synthetic dimethoxy flavones, *Asian Journal of Pharmaceutical and Clinical Research*, 2012; 5 (1):101-103.
- [6] Chirikova N K, Olennikov DN, Tankhaeva LM, Quantitative Determination of Flavonoid Content in the Aerial Part of Baical Scullcap (*Scutellaria baicalensis* Georgi), *Russian Journal of Bioorganic Chemistry*, 2010; 36 (7):915-922.
- [7] Maksimović Z, Malencić ND, Kovacević N, Polyphenol contents and antioxidant activity of *Maydis stigma* extracts, *Bioresource Technology*, 2004; 96 (8):873-877.
- [8] Sreenivasan S, Ibrahim D, Kassim MJNM, Free radical Scavenging Activity and Total Phenolic Compounds of *Gracilaria Changii*, *International Journal of Natural and Engineering Sciences*, 2007; 1 (3):115-117.
- [9] Kaur G, Jabbar Z, Athar M, Alam MS, *Punica granatum* (pomegranate) flower extract possesses potent antioxidant activity and abrogate Fe-NTA induced hepatotoxicity in mice, *Food Chemical Toxicology*, 2006; 44:984-993.
- [10] Benhammou N, AlikBenkkara F, Panovska TK, Antioxidant and antimicrobial activities of the *Pistacia lentiscus* and *Pistacia atlantica* extracts, *African Journal of Pharmacy and Pharmacology*, 2008; 2 (2):022-028.

- [11] Haddouchi F, Lazouni HA, Meziane A, Benmansour A, Etude physicochimique et microbiologique de l'huile essentielle de *Thymus fontanesii* Boiss & Reut, Afrique SCIENCE, 2009; 05(2):246-259.
- [12] Ali NAA, Julich W-D, Kusnick C, Lindequist U, Screening of Yemeni medicinal plants for antibacterial and cytotoxic activities, Journal of Ethnopharmacology, 2001; 74:173-179.
- [13] Bansemir A, Blume M, Schröder S, Lindequist U, Screening of cultivated seaweeds for antibacterial activity against fish pathogenic bacteria, Aquaculture, 2006; 252:79-84.
- [14] Chaouche T, Atik Bekkara F, Haddouchi F, Boucherit Z, Antibacterial activity of different extract of *Echium pycnanthum pommel*, Journal of Chemical and Pharmaceutical Research, 2012; 4(1):216-220.
- [15] Yazdani D, Zainal Abidin M A, Tan YH, Kamaruzaman S, Jaganath IB, Screening of phytochemical from ethnomedicinal plants in Malaysia for use against toxigenic *Aspergillus flavus*, Journal of Medicinal Plants Research, 2012; 6(42):5464-5468.
- [16] Nenaah, Antimicrobial activity of *Calotropisprocera* Ait. (*Asclepiadaceae*) and isolation of four flavonoid glycosides as the active constituents, World Journal Microbiology Biotechnology, 2013; 29:1255-1262.
- [17] Oyeleke SB, Dauda BEN, Boye OA, Antibacterial activity of *Ficuscapensis*, African Journal of Biotechnology, 2008; 7(10):1414-1417.
- [18] Ahameethunisa, Hopper, *In vitro* antimicrobial activity on clinical microbial strains and antioxidant properties of *Artemisia parviflora*, Annals of Clinical Microbiology and Antimicrobials, 2012; 11:30.
- [19] Hamouda Ali I, Doumandji A, Comparative phytochemical analysis and *in vitro* antimicrobial activities of the cyanobacterium *Spirulina platensis* and the green alga *Chlorella pyrenoidosa*: potential application of bioactive components as an alternative to infectious diseases, Bulletin de l'Institut Scientifique, 2017; 39:41-49.
- [20] Ramdan B, Amakran A, Bakrim N, Vannier B, Greche H, Nhiri M, Anti-glycation and radical scavenging activities of hydro-alcohol and aqueous extracts of nine species from *Lamiaceae* family, Journal of Medicinal Plants Studies, 2017; 5(1):331-345.
- [21] Nijveldt RJ, van Nood E, van Hoorn DN, Boelens PG, van Norren K, van Leeuwen PAM, Flavonoids: a review of probable mechanisms of action and potential applications, Journal of Clinical Nutrition, 2001; 74:418-425.
- [22] Nickavar B, Kamalinejad M, Izadpanah H, *In vitro* free radical scavenging activity of five *Salvia* species, Pak. J. Pharm. Sci., 2007; 20(4):291-294.
- [23] Wojdyło A, Oszmiański J, Czemerys R, Antioxidant activity and phenolic compounds in 32 selected herbs, Food Chemistry, 2007; 105 (3):940-949.
- [24] Hsu B, Coupar IM, NgK, Antioxidant activity of hot water extract from the fruit of the Doum palm, *Hyphaenethebaica*, Food Chemistry, 2006; 98:317-328.
- [25] Zha XQ, Wang JH, Yang XF, Liang H, Zhao LL, Bao SH, Luo JP, Xu YY, Zhou BB, Antioxidant properties of polysaccharide fractions with different molecular mass extracted with hot-water from rice bran, Carbohydrate Polymers, 2009; 78:570-575.
- [26] Moran JF, Klucas RV, Grayer RJ, Abian J, Becana M, Complexes of iron with phenolic compounds from soybean nodules and other legume tissues: Prooxidant and antioxidant properties, Free Radical Biology and Medicine, 1997; 22:861-870.
- [27] Bekara A, Ait Hamadouche N, Kahloula K, Sadi N, Aoues A, Etude phytochimique et activité antioxydante de l'extrait aqueux de *Pimpinella anisum* L., Algerian Journal of Natural Products, 2016; 4 (3):299-307.
- [28] Lacombe A, Wu VCH, Tyler S, Edwards K, Antimicrobial action of the American cranberry constituents; phenolics, anthocyanins, and organic acids, against *Escherichia coli* O157:H7, International Journal of Food Microbiology, 2010; 139:102-107.
- [29] Fertout-Mouri N, Latreche A, Mehdadi Z, Bengherraz Z, Activité antibactérienne de quatre extraits de *Teucrium polium* L. du mont de Tessala (Algérie occidentale), Bulletin de la Société royale des Sciences de Liège, 2016; 85:253-262.
- [30] Klančnik A, Piskernik S, Jeršek B, Možina SS, Evaluation of diffusion and dilution methods to determine the antibacterial activity of plant extracts, Journal of Microbiological Methods, 2010; 81:121-126.
- [31] Orhan DD, Özçelik B, Özgen S, Ergun F, Antibacterial, antifungal and antiviral activities of some flavonoids, *Microbiological Research*, 2010; 165:496-504.
- [32] Yala J-F, Ntsameso-Mve-Mba V, Azzizet Issembe Y, Lepengue NA, Souza A, Évaluation *in vitro* de l'activité antimicrobienne de l'extrait aqueux d'*Eryngium foetidum* récolté dans la ville de France ville, Journal of Applied Biosciences, 2016; 103:9886-9893.
- [33] Rodriguez-Vaquero MJ, Alberto MR, Manca de Nadra MC, Influence of phenolic compounds from wines on the growth of *Listeria monocytogenes*, Food Control, 2007; 18 (5):587-593.
- [34] Boban N, Tonkic M, Modun D, Budimir D, Mudnic I, Davorica Sutlovic D, Punda-Polic V, Boban M, Thermally treated wine retains antibacterial effects to food-borne pathogens, Food Control, 2010; 21(8):1161-1165.
- [35] Cowan MM, Plant products as antimicrobial agents, Clinical Microbiology Reviews, 1999; 12 (4):564-582.
- [36] Benyagoub E, Boulanouar A, Souid Ahmed M, Nebbou N, Bouloufa A, Essai d'évaluation de l'activité antibactérienne de la gomme arabique d'*Acacia tortilis*(Forssk) contre quelques souches bactériennes pathogènes, Bulletin de la Société Royale des Sciences de Liège, 2016; 85:237-252.
- [37] Cushnie TPT, Lamb AJ, Antimicrobial activity of flavonoids, International Journal of Antimicrobial Agents, 2005; 26:343-356.
- [38] Ahmad A, Kaleem M, Ahmed Z, Shafiq H, Therapeutic potential of flavonoids and their mechanism of action against microbial and viral infections-A review, Food Research International, 2015; 77:221-235.