

Available online on 15.11.2018 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

Antibiotic evaluation of odontogenic microbiological spectrum of orofacial infection

Neha Vishnoi¹, Sapna Singh¹, Ram Singh Bishnoi², M K Gupta³¹ Peoples Dental Academy, People's University, Bhopal, Madhya Pradesh, India² School of Pharmaceutical Sciences, Rajiv Gandhi Proudyogiki Vishwavidyalaya, Bhopal, Madhya Pradesh, India.³ Oral Maxillofacial Surgeon, Nagpur, Maharashtra, India

ABSTRACT

The aim of this study was to investigate the microbial flora and simultaneously evaluate its antibiotic response in patients associated with odontogenic infection. Samples for the analysis were taken from the Maxillofacial Surgery Department of People's Dental Academy, People's University, Bhopal. Our results resemble with current knowledge of odontogenic microbial flora. In this study frequently isolated isolates were 40(51%) of *Staphylococcus aureus*, 65(83%) of *Streptococcus mutans*, 23(29%) of *Streptococcus salivarius*, 30(38%) of *Streptococcus sanguis*, 21(27%) of *Streptococcus mitis*, 17(22%) of *Pseudomonas aeruginosa* and 14(18%) *Klebsiella pneumoniae*. The average sensitivity of antimicrobials against all isolated organisms were studied and it was found that common sensitive antimicrobials were clindamycin (88%), Metronidazole (79%), cefotaxime (72%), linezolid (72%), erythromycin (72%), amoxycillin (71%), ornidazole (67%), ciprofloxacin (67%), vancomycin (65%), imipenem (64%), cefadroxil (59%), ceftazidime (59%), azithromycin (58%), cefoperazone sulbactam (56%), whereas as resistant antimicrobials were penicillin (83%), levofloxacin (79%), gentamycin (77%), penicillin G (72%), cefuroxime (72%), ceftriazone (65%), ampicillin (65%), amikacin (64%), norfloxacin (59%), piperacillin (56%), clarithromycin (55%), ofloxacin (55%), ampicillin sulbactam (51%), azithromycin (50%), ampicillin sulbactam (50%), ceftazidime (50%). To attain maximum antimicrobial effect and minimal risk, clinical analysis and efficacy of antibiotics with regards to isolated bacteria is of great importance. Thus it indicates that microbiological analyses have significance in clinical diagnosis whereas evaluation of antibiotic susceptibility helps clinician to opt appropriate antibiotic regime for suffered patient.

Keywords: Orofacial infection, Odontogenic infection, Dental abscess, Microbiology, Antibiotics.

Article Info: Received 01 Oct, 2018; Review Completed 02 Nov 2018; Accepted 06 Nov 2018; Available online 15 Nov 2018



Cite this article as:

Vishnoi N, Singh S, Bishnoi RS, Gupta MK, Antibiotic evaluation of odontogenic microbiological spectrum of orofacial infection, Journal of Drug Delivery and Therapeutics. 2018; 8(6):179-182

DOI: <http://dx.doi.org/10.22270/jddt.v8i6.2238>

*Address for Correspondence:

Ram Singh Bishnoi, School of Pharmaceutical Sciences, Rajiv Gandhi Proudyogiki Vishwavidyalaya, Bhopal, Madhya Pradesh, India-462033

INTRODUCTION

Odontogenic infections are those which frequently occur in the pulp, periodontium or periapex of the tooth and mainly attributed due to normal flora of the mouth, whereas non odontogenic infections occur due to penetration of organisms through skin or mucous membrane and depend on the type and site of infection¹. The acute dental abscess is very much underestimated in terms of its morbidity and mortality. One of the commonest and major problems encountered in dental practice is bacterial infections. The most varied and vast flora in entire human body harbors in oral cavity in which dental abscess is usually polymicrobial which majorly includes *alpha-haemolytic streptococci*, coagulase negative *Staphylococci*, *Staphylococcus epidermidis*, *Staphylococcus aureus*, *Streptococcus viridans*

group, *Streptococcus anginosus* group, and strict anaerobes like *Prevotella*, *anaerobic cocci*, *Fusobacterium* species. Pain, swelling, erythema, suppuration are characteristic features of dental abscess ²⁻³.

The pathologic progression of this disease is a necrotic inflammation of pulp which extends in form of dentoalveolar abscess into periapical area and may penetrate through cortical bone to involve possible spaces⁴. Odontogenic infections usually react well to dental care, surgical intervention and antimicrobial therapy is essential for the speedy resolution of infection⁵. This study explores bacterial isolates in patients with odontogenic infection and its antibiotic susceptibility to different antibiotics which may commonly prescribe as prophylaxis in such types of infection in odontology.

MATERIALS AND METHOD

This study was conducted in People's Dental Academy, People's University, Bhopal. All patients presenting with maxillofacial infection in outpatient of the department were thoroughly screened with complete case history and detailed examination. The cases diagnosed as having maxillofacial infection was carried out in study whereas patients presenting with wound sepsis, history of anti-cancer chemotherapy or any other additional co-morbidity such as chronic renal failure, severe anemia and administration of antibiotic within a week were excluded from the study.

Collection of pus sample

Pus sample was preferentially collected by closed aspiration using an 18 gauge needle and 10 mL disposable syringe. Site of aspiration was chosen after careful examination and before aspiration it was cleaned with isopropyl alcohol. Intra-oral site was cleaned using 0.2% chlorhexidine mouth rinse, subsequently sterile dry gauze was used to wipe the area. Maximum amount of sample was aspirated in a single attempt to avoid contamination. In cases where significant aspirate was not available after incision and drainage, the sample was collected by sterile culture swabs. Immediately upon aspiration residual air was evacuated from the syringe and the needle was capped with a sterile rubber cork. The samples were transported to laboratory without any delay.

Assessment of microbial profile

Pus samples were directly observed by gram staining as well as it is processed on appropriate culture media for isolation and identification of micro-organisms. All culture media used in the study was prepared by reconstituting the commercially available dehydrated media of HiMedia, India. Samples were inoculated on blood agar, MacConkey agar media at 37°C for 18-24 hrs of incubation. After growth appears on media smear was prepared and Gram

staining was performed to study morphologic characteristics followed by various biochemical tests for identification of micro-organisms.

Antibiotic susceptibility testing

Isolated and identified colonies were then inoculated on Muller Hinton Agar for growing organisms. Antibiotic sensitivity for isolates was done by standard Kirby-Bauer Disk Diffusion Technique. Interpretations were carried out based on the diameter of the zone of inhibition as sensitive, moderate sensitive or resistant using manual provided by Himedia Pvt. Ltd., India. Antibiotics used were Amoxicillin-Clavulanic acid, Ampicillin, Ampicillin-Sulbactam, Cefadroxil, Clarithromycin, Clindamycin, Linezolid, Norfloxacin, Azithromycin, Vancomycin, Amikacin, Cefoparazone-Sulbactam, Cefotaxime, Ceftazidine, Ceftriaxone, Cefuroxime, Ciprofloxacin, Erythromycin, Gentamycin, Imipenem, Levofloxacin, Penicillin G, Metronidazole, Ornidazole and Ofloxacin.

RESULTS AND DISCUSSION

In our study 78 patients with maxillofacial odontogenic infection (Table 1) were included which were fulfilling the inclusion and exclusion criteria's of the study and 30 samples were taken of control (Healthy individuals). In the gender distribution of the subjects (Table 1) it was observed that out of 78 patients the females were 37(47.4%) and males were 41(52.5%). In control group out of 30 individuals 15 females and 15 males were enrolled. These findings were also be compared with gender distribution in which out of 90 patients 57(63.33%) cases were of males and 33(36.67%) cases were of female, In some studies male patients were 54% and female patients were 46% and 56(70%) were males and 24(30%) were female⁶⁻⁸. In our study also male patients was higher than females patients, thus males are much prone to the infection rather than females.

Table 1: Gender wise distribution

S No	Gender	No. of Patients (%) (n= 78)	Control Group (n=30)
1	Male	41 (52.5%)	15
2	Female	37(47.4%)	15

In the abscess distribution by tooth type (Table 2) it has been found that out of 78 patients the total 190 teeth were affected in which 108(57%) teeth are of molars, 16(8%) are canines and 33(17%) are premolars and incisors tooth (Table 2). No teeth were involved in control group. The

most frequently implicated teeth were those located in the lower posterior segments (61.5%), followed by the lower molars (26.6%). In other study they found the lower molars to be the most causal teeth⁹.

Table 2: Abscess distribution by tooth type

S.No.	Tooth type	No. of cases	No. of tooth involving	Percentage (%)	Control group
1	Molars	62	108	57%	00
2	Premolars	20	33	17%	00
3	Canines	10	16	08%	00
4	Incisors	13	33	17%	00
	Total	105	190	100	00

In some studies also stated that most odontogenic infections arise as a sequel to pulp necrosis caused by caries, periodontal infections, gingivitis, pericoronitis, trauma and surgery are other sources responsible for orofacial infections. In our study also it was observed that the main origin/cause of odontogenic infection (Table 3)

was dental caries i.e 62 (79.4%), followed by gingivitis 44(56.4%), periodontitis 30(38.4%), periapical 5(6.4%) and pericoronitis 3(3.8%). Local odontogenic infections usually originate from apical periodontitis (66.7%-70.7%) or initially from dental caries (33.8%-80.6%). Other dental

diseases causing local infection complications are pericoronitis, alveolar osteitis, periodontitis and cysts⁸⁻¹⁵.

Table 3: Origin of infection

S.No	Origin of infection	No. of patients (%)	Control Group
1	Periodontal	30 (38.4%)	00
2	Pericoronitis	03 (03.8%)	00
3	Periapical	05 (06.4%)	00
4	Dental caries	62 (79.4%)	00
5	Gingivitis	44 (56.4%)	00

It was observed that in total 78 abscess patients (Table 4) there was 40(51%) of *Staphylococcus aureus* isolates, 65(83%) of *Streptococcus mutans*, 23(29%) of *Streptococcus salivarius*, 30(38%) of *Streptococcus sanguis*, 21(27%) of *Streptococcus mitis*, 17(22%) of *Pseudomonas aeruginosa* and 14(18%) are of *Klebsiella pneumoniae*. Total isolates were 210 in odontogenic patients in which multiple isolates are found in single patient. Studies by some scientists covering diverse populations and performed at different time periods show that maxillofacial infections are polymicrobial mixed micro flora¹⁶⁻²¹. The most common micro-organisms in dentoalveolar infection is *Streptococcus viridans*²². The results of our study are in

concurrence with those of Rega (2006) et al and Rao D et al (2010)^{7,23}. In many studies it was found that *Staphylococci* are now more frequent colonizer of oral tissues than previously thought. The frequent isolation of *Staphylococci* in pus samples from odontogenic infections have been reported in previous studies^{19,24}. The most common organisms isolated from aerobic bacteria were *Streptococcus viridans* 24(36.4%), followed by *Klebsiella* 18(27.3%), *Pseudomonas aeruginosa* 12(18.2%), coagulase negative *Staphylococci* 6(9.1%), *Nisseria* 3(4.54%) and *Enterobacter* spp. 3(4.54%)⁶. The isolation of *pseudomonas* in 13 cases (11.60%) is high when compared to other studies¹⁰.

Table 4: Isolates from patients (n=78)

S No.	Isolates	No. of isolates (%)	Control (%)
1	<i>Staphylococcus aureus</i>	40 (51%)	12
2	<i>Streptococcus mutans</i>	65 (83%)	28
3	<i>Streptococcus salivarius</i>	23 (29%)	25
4	<i>Streptococcus sanguis</i>	30 (38%)	07
5	<i>Streptococcus mitis</i>	21 (27%)	15
6	<i>Pseudomonas aeruginosa</i>	17 (22%)	04
7	<i>Klebsiella pneumonia</i>	14 (18%)	02
	Total Isolates	210	93

The average of antimicrobials activity against all different isolates in which sensitive antimicrobials (Figure 1) are clindamycin (88%), metronidazole (79%), cefotaxime (72%), linezolid (72%), erythromycin (72%), amoxclav (71%), ornidazole (67%), ciprofloxacin (67%), vancomycin (65%), imipenem (64%), cefadroxil (59%), ceftazidime (59%), azithromycin (58%), cefoperazone sulbactam (56%). A total of 27 antibiotics were tested in the present study. Over all excellent responses were evident for Clindamycin, Linezolid and Metronidazole, Cefotaxime, Erythromycin. Baumgartner et al. 2003, Chang et al. 2005 also reported the high sensitivity for clindamycin²⁵⁻²⁶, while Lewis M et al (1990) and Boyanova et al (2006) reported Clindamycin and Metronidazole to be highly

efficient against gram negative rods^{17,27}. Salins BM et al (2006) have also shown high susceptibility for Linezolid and Clindamycin²⁸. Gutierrez-Perez JL et al (2004) who recommended Amoxicillin-Clavulanic acid as the first line antibiotic, our study found an antibiotic sensitivity on 70.5% to this antibiotic²⁹. *In-vitro* susceptibility (88.46%) to Clindamycin and the clinical response to this antimicrobial in the present study appears to justify its choice as the first line drug for odontogenic sepsis. In the study of Flynn TR (2000) all the aerobic bacteria showed 3(14.3%) sensitive to Ampicillin and 10(47.6%) resistance to the same. Most of the organisms were sensitive to Ceftriaxone 20(95.2%), Levofloxacin 19(90.5%) and 17(81%) sensitive for Amoxicillin and Clavulanic acid⁵.




Figure 1 Antimicrobial profile (Average) of all isolates

CONCLUSION

Management of odontogenic infection should consider the therapeutics success which lies in control of infection whether by using mechanical surgical debridement or by antimicrobial therapy. Microbiological analysis is one of the reliable way to define therapeutic success by precisely characterize prognosis of recurrence and is enable to select most appropriate antibiotic therapy to increase therapeutic efficacy.

REFERENCES

1. Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE. Defining the normal bacterial flora of the oral cavity, *Journal of Clinical Microbiology*, 2005; 43(5):721-32.
2. Al-Qamachi LH, Aga H, MaMahon. Microbiology of odontogenic infections in deep neck spaces: a retrospective study, *British Journal of Oral and Maxillofacial Surgery*, 2010; 48:37-39.
3. Maeda Y, Goldsmith CE, Coulter WA, Mason C, Dooley JSG, Lowery CJ, Moore JE. The viridans group streptococci, *Reviews in Medical Microbiology*, 2010; 21:69-79.
4. Samaranayake L. Essential microbiology for dentistry. 3rd ed. Churchill Livingstone; 2006. P. 232-236.
5. Flynn TR. The swollen face: severe odontogenic infections, *Emergency Medicine Clinics of North America*, 2000; 18:481-519.
6. Santosh AN, Viresh AN, Sharmada BK. Microbiology and antibiotic sensitivity of odontogenic space infection, *International journal of Medical and Dental Sciences*, 2014; 3(1):303-313.
7. Rega AJ, Aziz SR, Ziccardi VB. Microbiology and antibiotic sensitivities of head and neck space infections of odontogenic origin, *Journal of Oral and Maxillofacial Surgery*, 2006; 64:1377-80.
8. Kohli M, Mathur A, Kohli M, Siddiqui S R. In vitro evaluation of microbiological flora of orofacial infections, *Journal of Oral and Maxillofacial Surgery*, 2009; 8(4):329-33.
9. Sanchez R, Mirada E, Arias J, Pano JR, Burgueno M. Severe odontogenic infections: Epidemiological, microbiological and therapeutic factors, *Medicina Oral Patología Oral y Cirugía Bucal*, 2011; 16(5):670- 6.
10. Gill Y, Scully C. Orofacial odontogenic infections : Review of microbiology and current treatment, *Oral Surgery, Oral Medicine, Oral Pathology*, 1999; 70(2):155-58.
11. Ylijoki S, Suuronen R, Jousimies Somer H. Differences between patients with or without the need for intensive care due to severe odontogenic infections, *Journal of Oral and Maxillofacial Surgery*, 2001; 59: 867-72.
12. Flynn TR, Shanti RM, Hayes C. Severe odontogenic infections, part 2:prospective outcomes study *Journal of Oral and Maxillofacial Surgery*, 2006; 64:1104-13.
13. Saito C, Gulinelli J, Marao H, Garcia I, Filho O, Sonoda C, Poi W, Panzarini S. Occurrence of odontogenic infections in patients treated in a postgraduation program on maxillofacial surgery and traumatology, *Journal of Craniofacial Surgery*, 2011; 22:1689-94.
14. Indresano AT, Haug RH, Hoffman MJ. The third molar as a cause of deep space infections, *Journal of Oral and Maxillofacial Surgery*, 1992; 50:33-5.
15. Smith JL, Kellman RM. Dentigerous cysts presenting as head and neck infections, *otolaryngology, head and neck surgery*, 2005; 133: 715-17.
16. Heimdahl A, Von Konow L, Satoh T. Clinical appearance of orofacial infections of odontogenic origin in relation to microbiological findings, *Journal of Clinical Microbiology*, 1985; 22:299-302.
17. Lewis MAO, MacFarlane TW, McGowan DA. A microbiological and clinical review of to alveolar abscess, *British Journal of Oral and Maxillofacial Surgery*, 1990; 28:359-66.
18. Kuriyama T, Nakagawa K, Karasawa T, Saiki Y, Yamamoto E, Nakamura S. Past administration of β -lactam antibiotics and increase in the emergence of β -lactamase- producing bacteria in patients with orofacial odontogenic infections, *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 2000; 89:186-92.
19. Storoe W, Haug RH, Lillich TT. The changing face of odontogenic infections, *Journal of Oral and Maxillofacial Surgery*, 2001; 59(7):739-48.
20. Huang TT, Liu TC, Chen PR, Tseng FY, Yeh TH, Chen YS. Deep neck infection: analysis of 185 cases, *Head & Neck*, 2004; 26:854-60.
21. Robertson D and Smith AJ. The microbiology of the acute dental abscess, *Journal of Medical Microbiology*, 2009; 58:155-62.
22. Ulubau IC, Jaunay T, Goss AN. Severe odontogenic infections, *Australian Dental Journal Medications Supplement*, 2005; 50(2): S74-S81.
23. Rao DD, Desai A, Kulkarni RD. Comparison of maxillofacial space infection in diabetic and nondiabetic patients, *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 2010; 110:e7-e12.
24. Kannangara DW, Thadepalli H, McQuirter JL. Bacteriology and treatment of dental infections, *Oral Surgery, Oral Medicine, Oral Pathology*, 1980; 50(2):103-09.
25. Baumgartner JC and Tian X. Antibiotic Susceptibility of Bacteria Associated with Endodontic Abscesses, *Journal of Endodontics*, 2003; 29(1):44-47.
26. Chang CM, Lub FH, Guod HR, Ko WC. *Klebsiella pneumoniae* fascial space infections of the head and neck in Taiwan: emphasis on diabetic patients and repetitive infections, *Journal of Infection*, 2005; 50:34-40.
27. Boyanova L, Kolarov R, Gergova G, Deliverska E, Madjarov J, Marinov M and Mitov I. Anaerobic bacteria in 118 patients with deep space head and neck infections from the University Hospital of Maxillofacial Surgery, Sofia, Bulgaria, *Journal of Medical Microbiology*, 2006; 55:1285-89.
28. Salinas MB, Noelia CR, Leonardo BA, Cosme GE. Antibiotic susceptibility of the bacteria causing odontogenic infections, *Medicina Oral Patología Oral y Cirugía Bucal*, 2006; 11: E70-5.
29. Gutierrez Perez JL, Perea Perez EJ, Romero Ruiz MM, Giron Gonzalez JA. Orofacial infections of odontogenic origin, *Oral Medicine and Pathology*, 2004; 9:280-7.