

Available online on 15.12.2018 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

The anti-arthritis and anti-inflammatory activities of aqueous extract powder bark of *Anthocleista schweinfurthii* Gilg (Loganiaceae)

Djova Steve Valdi ^{1*}, Nyegue Maximilienne Ascension ², Etoa François-Xavier ²

¹Department of Biochemistry, University of Yaounde I, PO Box 812 Yaounde, Cameroon.

²Department of Microbiology, University of Yaounde I, PO Box 812 Yaounde, Cameroon.

ABSTRACT

Aqueous extract bark of *Anthocleista schweinfurthii* (ASE) was investigated for anti-arthritis, anti-inflammatory activities and phytochemicals screening. Phytochemical screening was carried out according to LC-MS procedures, the anti-inflammatory effect of ASE was evaluated *in vitro* inhibition of 15-lipoxygenase (15-LOX) enzyme and proteinase inhibiting action assays. The anti-arthritis activity was performed according to the Bovine Serum Albumin (BSA) test. The chemical composition of ASE showed that it was mainly composed of isocyanate, biphenyltriol, methoxyflavanone, soscoparin and glucopyroside. In the anti-inflammatory assays, ASE showed selective inhibition of 15-lipoxygenase with IC_{50} value of $37.19 \pm 0.54 \mu\text{g/mL}$ and moderate activity against the inhibition of proteinase action with IC_{50} value of $244.33 \pm 2.08 \mu\text{g/mL}$. Both the methods showed that the ASE posses good anti-inflammatory activity. ASE had strong *in vitro* anti-arthritis activity with IC_{50} of $13.67 \pm 0.88 \mu\text{g/mL}$, the positive control Diclofenac had IC_{50} value of $4.66 \pm 0.33 \mu\text{g/mL}$. The result of this study justified the use of this plant in traditional Cameroonian medicine in the treatment of inflammation. ASE may offer a new source of potential therapeutic agents for the effective treatment of arthritis and inflammation.

Keywords: *Anthocleista schweinfurthii*, phytochemical screening, anti-arthritis activity, anti-inflammatory activity,

Article Info: Received 20 Oct 2018; Review Completed 03 Dec 2018; Accepted 04 Dec 2018; Available online 15 Dec 2018



Cite this article as:

Djova SV, Nyegue MA, Etoa FX, The anti-arthritis and anti-inflammatory activities of aqueous extract powder bark of *Anthocleista schweinfurthii* Gilg (Loganiaceae), Journal of Drug Delivery and Therapeutics. 2018; 8(6-s):174-181
DOI: <http://dx.doi.org/10.22270/jddt.v8i6-s.2212>

***Address for Correspondence:**

Djova Steve Valdi, Department of Biochemistry, University of Yaounde I, PO Box 812 Yaounde, Cameroon. Email: djova.stevevaldi@yahoo.fr

INTRODUCTION

Inflammation is a defense response of our body which involves a complex reaction of vascularised tissues to infection, allergens, toxin exposure and cell injury^{1,2}. Conversely, uncontrolled inflammatory response is the main cause of a vast and continuous disorders including cancer, septic shock, diabetes, atherosclerosis, obesity, heart disease, age-related muscular degeneration, chronic obstructive pulmonary disease, multiple sclerosis, arthritis, rheumatoid arthritis^{3,4,5}.

Rheumatoid arthritis is a chronic, systematic inflammatory disease predominantly affecting the joints and periarticular tissues⁶. Currently, steroid drugs, non steroidal anti-inflammatory drugs (NSAIDs) and immunosuppressants, usually used for the relief of inflammatory diseases, require long-term treatment and their use is often associated with serious side effects such as bleeding, gastrointestinal and peptic ulcers⁷. Hence, continuous search for the development of potent anti-inflammatory drugs with fewer side effects is necessary from medicinal plants.

A. schweinfurthii is a plant of the Loganiaceae family, it is a source of traditional recipe for the treatment of male and female fertility problems in Togo, Nigeria, Cameroon, Gabon and Equatorial Guinea⁸. The decoction of the bark of *A. schweinfurthii* is used for the treatment of sexually transmitted infections⁹. A decoction of barks of root and leaves of *A. schweinfurthii* in Tanzania, is used for the treatment of malaria. A decoction of bark and leaves of *A. schweinfurthii* is used in Tanzania for the treatment of pain, injury and inflammatory diseases^{9,10}. In Cameroon, Gabon, Equatorial Guinea, Congo, and Tanzania, a decoction of *A. schweinfurthii* bark is used for the treatment of microbial infections and bronchitis¹¹. In Congo, the decoction of bark of *A. schweinfurthii* is used to treat hernia¹². In Nigeria, *A. djalonensis* is used in traditional medicine to treat breast cancer¹³. In Cameroon, the decoction of bark of *A. schweinfurthii* commonly known as "Bopolopolo" in the Douala language is used for the treatment of gastrointestinal disorders, gastric ulcers, abdominal pain, *A. schweinfurthii* is used for its galactogenic properties^{9,14}.

The *in vitro* antioxidant activity, cytotoxicity and genotoxicity of extracts from the leaves and barks of *A.*

schweinfurthii have been already done¹⁵. Djeussi et al¹⁶, evaluated the antibacterial properties of the methanol extract of *A. schweinfurthii*; the vasoconstrictor and inotropic effects induced by the root bark extracts of *A. schweinfurthii* was investigated by Ngombe et al¹⁷. A new steroid, schweinfurthiin, two known compounds, bauerenone, bauerenol and xanthones were isolated in the dichloromethane/methanol extract of the roots of *A. schweinfurthii* by Mbouangouere et al¹⁸. The acute and subacute toxicity of the aqueous extract of *A. schweinfurthii* bark was investigated by Christophe et al⁹. However, as far as known, no work has been devoted to the study of the anti-arthritis and anti-inflammatory activities of aqueous extracts of *A. schweinfurthii*. Therefore, the aim of the present study was to evaluate the anti-arthritis and the anti-inflammatory activities of aqueous extract powder barks of *Anthocleista schweinfurthii* Gild (Loganiaceae).

MATERIALS AND METHODS

Plant material and extraction procedure

A. schweinfurthii plant was collected at Loum in the Littoral region of Cameroon in August 2016, it is commonly known in Cameroon as "Betbreu" in Medumba local language and "Bopolopolo" in Douala local language. The plant identification was done at the Cameroon National Herbarium by comparison with specimen number 52349/HNC of *A. schweinfurthii* Gild (Loganiaceae).

The barks of *A. schweinfurthii* were air-dried for one week at room temperature and weighed. The sample were then ground to fine powder in a mortar and 500 g of dried powder was soaked for 48 hours in water. The mixing was filtered through Whatman No. 1 filter paper and lyophilized using the lyophilizator Biobase. Then, the aqueous extracts of *A. schweinfurthii* was collected in Eppendorf tubes and preserved in a refrigerator at 4 °C for further use.

LC-MS procedures

LS-MS analysis of OSE was carried out following a modified method of Abay et al¹⁹. High resolution mass spectra were obtained with an OTOF Spectrometer (Bruker, Germany) equipped with a HESI source. The spectrometer was operated in positive mode (mass range: 100-1000, with a scan rate of 1.00 Hz) with automatic gain control to provide high-accuracy mass measurements within 2 ppm deviation using Na Formate as calibrant. The spectrometer was attached to an Ultimate 3000 (Thermo Fisher, USA) HPLC system consisting of LC-pump, Diode Array Detector (DAD) ($\lambda = 260$ nm), auto sampler (injection volume 5 μ L). MS analyses were performed in multiple reaction monitoring (MRM) mode by measuring the fragmentation products of the protonated pseudomolecular ions of COC, BE and EME and their internal standards in three different fragmentation windows. The ESI voltage was set at 4.5 kV. Parameters were optimised by injection of known amounts of the extract and evaluating retention times, analyte response, peak widths and shapes. C₁₈ reversed-phase column oven (30°C) was used. The isocratic mobile phase was made up of two solvents water/AcN at different concentration water/AcN 95:5 and water/AcN 90:10. The starting eluent was used for equilibration of the column. Nitrogen was used as nebulising gas at a pressure of 40 psi. Dry gas debit was set at 8 L/min and capillary temperature at 200 °C.

Anti-inflammatory assays

Inhibition of 15-lipoxygenase (15-LOX) enzyme assay

The assay was performed according to a previously described procedure by Pinto et al²⁰ and Delong et al²¹ with slight modifications. The assay measures the Fe³⁺/xylenol orange complex at 560 nm formed when the extract reacts with the assay reagents. Briefly, 15-LOX from Glycine max was incubated with compounds or standard inhibitor at 25 °C for 5 min. Then linoleic acid (final concentration, 140 mM) in Tris-HCl buffer (50 mM, pH 7.4) was added and the mixture was incubated at 25 °C for an additional 20 min in the dark. The reaction was terminated by the addition of 100 mL of FOX reagent consisting of sulphuric acid (30 mM), xylene orange (100 mM), iron (II) sulphate (100 mM) in methanol/water (9:1). For the control, only LOX solution and buffer were pipetted into the wells. Blanks contained the enzyme LOX during incubation, but the substrate (linoleic acid) was added after the FOX reagent. The LOX inhibitory activity was evaluated by calculating the percentage of the inhibition of hydroperoxide production from the changes in absorbance values at 560 nm after 30 min at 25 °C.

$$\% \text{ inhibition} = ((A_{\text{control}} - A_{\text{blank}}) - (A_{\text{sample}} - A_{\text{blank}})) / (A_{\text{control}} - A_{\text{blank}}) \times 100$$

Where, A_{control} is the absorbance of control well, A_{blank} is the absorbance of blank well and A_{sample} is the absorbance of sample well.

Proteinase Inhibitory Action

The anti-proteinase activity of ASE was determined using the method of Oyedepo et al²² with slight modifications. The reaction mixture (2 mL) was containing 0.06 mg trypsin, 1 mL of 20 mM Tris HCl buffer (pH 7.4) and 1 mL test sample of different concentrations. The reaction mixture was incubated at 37 °C for 5 min and then 1 mL of 5 % (W/V) casein was added. The mixture was incubated for an additional 20 min, 2 mL of 70 % perchloric acid was added to terminate the reaction. Cloudy suspension was centrifuged and the absorbance of the supernatant was read at 210 nm against buffer as blank. The experiment was performed in triplicate. The percentage of inhibition of proteinase inhibitory activity was calculated.

Anti-arthritis assay

Bovine Serum Albumin (BSA) denaturation assay

Protein denaturation was performed as described by Sakat et al²³ with slight modifications. The test solution consisting of 1 mL of different concentrations of extracts preparation ranging from 0.5 to 1.5 mg/mL or standard Diclofenac sodium 0.1 and 0.25 mg/mL was mixed with 1 mL of egg albumin solution (1 mM) and incubated at 27 ± 1 °C for 15 min. Denaturation was induced by keeping the reaction mixture at 70 °C in a water bath for 10 min. After cooling, the turbidity was measured using the Jenway 6305 spectrophotometer at 660 nm. Percentage inhibition of denaturation was calculated from control where no drug was added. Each experiment was done in triplicate.

Statistical analysis

The results are presented as means of three experiments. Statistical significance between groups was calculated by using a paired t-test with GraphPad Prism software (version 7). Values were expressed as mean ± SD and differences were considered significant statistically if $P < 0.05$.

RESULTS

Figure 1 resumes the number of phytoconstituents present in the extract by comparing with the blank.






Figure 1: LC-MS chromatogram obtained from ASE

The compounds constituting the ASE were identified through interpretation of their mass spectrum obtained by LC/MC in comparison with previously reported data from the literature. The compounds were identified from their protonated molecular ion $[M]^+$. The figure 1 shown that ASE exhibited presence of compound 2 to compound 25. The compounds with high amount were also identified in the case of this study. The compound 3 exhibited a pseudo-molecular ion peak at m/z 239,067 $[M+H^+]^+$ in the HRESIMS corresponding to the molecular formula of $C_{11}H_{16}N_6O$ and retention time of 1.3 min, compound 3 was identified as 3-Isocyanato-8-(4-pyridinyl)[1,2,4]triazolo[4,3-b] pyridazin. The compound 4 had a pseudo-molecular ion peak at m/z 203.06 $[M+H^+]^+$ in the HRESIMS corresponding to the molecular formula of $C_{12}H_{10}O_3$ with retention time of 1.8 min, identified as biphenyltriol. The compound 8 showed a pseudo-molecular ion peak at m/z 302.0874 $[M+Na^+]^+$ in the HRESIMS corresponding to the





molecular formula of $C_{16}H_{14}O_6$ and retention time of 3.3 min, the compound 8 was identified as 3',5,7-Trihydroxy-4'-methoxyflavanone. The compound 13 had a pseudo-molecular ion peak of m/z 362.3123 $[M+Na^+]^+$ in the HESIMS corresponding to the molecular formula of $C_{21}H_{38}N_4O$ with retention time of 5.1 min, was identified as 8-[1-(1-Azepanyl)-2-propanyl]-2-(3-methylbutyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one. The compound 17 showed a pseudo-molecular formula ion peak at m/z 302.0874 $[M+Na^+]^+$ in the HRESIMS corresponding to the molecular formula of $C_{21}H_{20}O_{10}$ and retention time of 5.9, the compound 17 was identified as 3,8-Dihydroxy-6-methyl-9,10-dioxo-9,10-dihydro-1-anthracenyl beta-D-glucopyranoside. The last one, the compound 20 exhibited a pseudo-molecular formula ion peak at m/z 463,12 $[M+Na^+]^+$ in the HRESIMS corresponding to the molecular formula of $C_{22}H_{22}O_{11}$ with retention time of 6.4 was identified as soscoparin.


Compound 3



3

3-Isocyanato-8-(4-pyridinyl)[1,2,4]triazolo[4,3-b]pyridazine

Compound 4**biphenyltriol****Compound 8****3',5,7-Trihydroxy-4'-methoxyflavanone****Compound 13****8-[1-(1-Azepanyl)-2-propanyl]-2-(3-methylbutyl)-1,3,8-triazaspiro[4.5]dec-1-en-4-one****Compound 17****3,8-Dihydroxy-6-methyl-9,10-dioxo-9,10-dihydro-1-anthracenyl beta-D-glucopyranoside**

Compound 20**Soscoparin****Anti-inflammatory activity****Ferrous oxidation-xylenol orange (Fox) assay**

Inhibition of 15-lipoxygenase by ASE using *in vitro* 15-Lox enzyme assay showed average inhibition of 15-Lox with IC_{50} value of $37.19 \pm 0.54 \mu\text{g/mL}$ while the standard Quercetin with IC_{50} value of $6.71 \pm 0.1931 \mu\text{g/mL}$ which is shown in figures 2 and 3




Figure 2: % of Inhibition of the 15-lipoxygenase activity of ASE and Quercetin at different concentrations from which IC_{50} values of OSE and Quercetin was obtained. Values are mean \pm SD of three experiments.




Figure 3: Comparison of IC_{50} of ASE with Quercetin. Data are expressed as mean \pm SD, Quercetin was used as a reference compound. Statistical differences between quercetin and ASE as analyzed by the paired t-test ($^{****}P < 0.0001$).

Proteinase Inhibitory Action

The ASE showed moderate inhibition of proteinase action with IC_{50} value of $244.33 \pm 2.08 \mu\text{g/mL}$ whereas for Aspirin, a standard drug had IC_{50} value of $53.79 \pm 2.61 \mu\text{g/mL}$ (figure's 4 and 5).




Figure 4: % of Inhibition of the proteinase action of ASE and Aspirin at different concentrations from which IC_{50} values of ASE and Aspirin was obtained. Values are mean \pm SD of three experiments.




Figure 5: Comparison of IC_{50} of ASE with Aspirin. Data are expressed as mean \pm SD, Aspirin was used as a standard drug. Statistical differences between Aspirin and ASE as analyzed by the paired t-test ($***P < 0.0001$).

Anti-arthritic activity

The anti-denaturation study for investigating anti-arthritis activity was ascertained in this study using Bovine Serum Albumin (BSA) assay. The ASE displayed a strong inhibition activity of Bovine Serum Albumine denaturation with IC_{50} value of $13.67 \pm 0.88 \mu\text{g/mL}$ as compared with that of the standard drug Diclofenac with IC_{50} value of $4.66 \pm 0.33 \mu\text{g/mL}$. The results were illustrated by the figures 6 and 7.




Figure 6: % of Inhibition of the protein denaturation activity of ASE and Diclofenac sodium at different concentrations from which IC_{50} value of ASE and Diclofenac sodium was obtained. The error bars represent the standard deviation of measurement of the absorbance, experiment was done in triplicate.




Figure 7: Comparison of IC₅₀ of ASE with diclofenac sodium. Data are expressed as mean \pm SD, Diclofenac sodium was used as a standard drug. Statistical differences between Diclofenac sodium and ASE as analyzed by the paired t-test ($^{****}P<0.0001$).

DISCUSSION

We reported the identification of isocyanate, Soskoparin, biphenyltriol, methoxy-flavone, glycoside and flavonoids in the ASE. This result confirms literature review which revealed that *Antocleista* genus is a rich source of biocompounds such as alkaloids, xanthone, scoiridoids, and terpene²⁴, contradicting in other hands to some author who reported that reducing sugar and glycosides were found to be absent in both the leaf and the stem-bark of the *Anthocleista* species^{25,26}. Inflammation is induced by several pathway, one of the most pathway, it is initiated by the action of lipoxygenase on arachidonic acid producing trioxilins and leukotriene²⁹. Agent which possess the ability of inhibition of the 15-Lipoxygenase could be used as a potent anti-inflammatory agent. The anti-inflammatory effect was ascertained in this study by the inhibition of 15-lipoxygenase and proteinase inhibition assays. Statistically ($^{***}P<0.0001$) ASE showed selective inhibition of 15-lipoxygenase compared with that of the standard Quercetin. Proteinases have been implicated in arthritic reactions, it was previously reported that proteinase plays an important role in the development of tissue damage during in inflammatory reactions and significant level of protection was provided by proteinase inhibitors³⁰. The present investigation has shown that, ASE had moderate proteinase inhibition action, statistically significant ($^{***}P<0.0001$) compared with that of the standard drug, Aspirin. There are several inflammatory study showing anti-inflammatory activity of the different parts of *Anthocleista* species. Analgesic properties of *A. vogeli* stem barks has been reported by Mbiantcha et al³¹, Baba and Usifoh³² showed that the ethanol-water extract of *A. djalonensis* root revealed significant anti-inflammatory activity. The anti-lipoxygenase activity of ASE may be linked to the presence of biomolecules such as flavonoids, alkaloids and terpenes³³. Flavonoids are known to interfere with the different stages of the arachidonate cascade via cyclooxygenase or lipoxygenase pathways to alleviate inflammatory responses³⁴. Protein denaturation is

one of the key features of inflammatory tissue and it undergoes denaturation and antigens are expressed which are associated with type-III hypersensitivity reaction, which in turn is related to diseases such as serum sickness, glomerulonephritis, rheumatoid arthritis and systemic lupus erythematosus^{4,27,35}. Mechanism of denaturation probably involves alterations in electrostatic, hydrogen, hydrophobic and disulphide bonding³⁴. It is believed that agent that can help in anti-protein denaturation could be used as a potent anti-inflammatory drug in future²⁸. The anti-denaturation study for investigating anti-arthritis activity was ascertained in this study using BSA assay. ASE had showed a strong *in vitro* anti-arthritis activity statistically significant ($^{***}P<0.0001$) as compared with that of the standard drug Diclofenac. These findings support the traditional use of *Anthocleista shweinfurthii* as anti-arthritis and in treatment of inflammatory disease in Cameroon.

CONCLUSION

Aqueous extracts of the bark of *A. schweinfurthii* was evaluated for their anti-arthritis and anti-inflammatory activities and their phytochemical study. The findings confirm the ethno medicinal use of *A. schweinfurthii* to manage the inflammation condition and arthritic. Isocyanate, biphenyltriol, methoxyflavanone, soskoparin and glucopyroside were identified in ASE. The isolation and characterization of these biomolecules and the understanding of the therapeutic targets of ASE will in the future allow the development of potential anti-inflammatory agents from *A. schweinfurthii* bark.

CONFLICT OF INTERESTS

The authors declared that there is no conflict of interest regarding the publication of this paper.

Acknowledgment

Authors are very grateful to Cameroon National Herbarium (Yaounde) for identification of the plant.

REFERENCES

- Vane JR and Botting RM, New insights into the mode of action of anti-inflammatory drugs, *Journal of Inflammation Research*, 1995; 44(1):1-10.
- Umapathy E, Ndebia EJ, Meeme B, Menzina P, Nkeh-Chungay BN, and Iputo JE, An experimental evaluation of *Albuca seroseae* aqueous extract on membrane stabilization, protein denaturation and white blood cell migration during acute inflammation, *Journal of Medicinal Plant Research*, 2010; 4(9):789-795.
- Harsh Mohan. Text book of pathology. 6th ed. Jaypee publications ; 2010. P. 851.
- Kishore G, Siva G, Sindhu ES, *in Vitro* Anti-Inflammatory and Anti-Arthritic Activity of Leaves of *Physalis Angulata* L, *international journal pharmaceutical ind research*, 2011; 1:211-213.
- Kumar V, Bhat ZA, Kumar D, Bohra P, and Sheela S, *in vitro* anti-inflammatory activity of leaf extracts of *Basella alba* Linn.Var Alba, *International Journal of Drug Development and Research*, 2011; 3(3):176-17.
- Deepika A, Mukesh M, Neha G and Vidya P, In vitro anti-inflammatory and anti-arthritic activity in methanolic extract of *coccus hirsutus*. Diels. *In vivo* and *in vitro*, *International journal of pharmaceutical Science and research*, 2014 ; 5(5):1957-1962.
- Corley DA, Kerlikowske K, Verma R, Buffler P, Protective association of aspirin/ NSAIDs and esophageal cancer, a systematic review and meta-analysis. *Gastroenterology*. 2003; 124(1):47-56.
- Ruijter A, *Anthocleista schweinfurthii* Gilg. In Schmelzer, G.H. & Gurib-Fakim, A. (Editors). PROTA (Plant Resources of Tropical Africa), Wageningen, Netherlands. 2007.
- Christophe M, Frida L, Cline N, Zacharie S, Emilie N, Vernyuy PT, Evaluation of acute and subacute toxicity of stem bark aqueous extract of *Anthocleista schweinfurthii* (Loganiaceae), *World journal of pharmacy and pharmaceutical sciences*, 2015; 4:197-208.
- Burkill HM, The useful plants of West Tropical Africa. 2nd Edition. Volume 3, Families J-L. Royal Botanic Gardens, Kew, Richmond, United Kingdom ; 1995. P.857.
- Fowler DG, Traditional Fever Remedies: A List of Zambian plants. http://www.Gift.org/ritam/news/Traditional_Fever_remedies1.Pdf (accessed 20.04.2016)
- Schmelzer GH, In Schmelzer GH, Gurib-Fakim A, (Eds.), Arroo R, (Associate Ed.) Lemmens RHMJ, Oyen LPA, (General Ed.), *Plant Resources of Tropical Africa: Medicinal plants*, 2008; 11:1.
- Gbadamosi IT and Erinoso SM, A review of twenty ethnobotanicals used in the management of breast cancer in Abeokuta, Ogun State, Nigeria. *African Journal of pharmacy and Pharmacology*, 2016; 10(27):546-564.
- Adongo SO, Morongo J, Anjou R, New Analysis of selected essential elements of medicinal plants used by Chuka community, Tharaka Nithi county, Kenya, *Science Journal Technology Special issue*, 2012; 87-94.
- Djova SV, Nyegue MA, Afagnigni AD, Etoa FX, *In vitro* antioxidant activity, cytotoxicity and genotoxicity of extracts from the leaves and barks of *Anthocleista schweinfurthii* gilg (Loganiaceae), *Asian Journal of Research in Biological and Pharmaceutical Sciences*, 2018 ; 6(2):58-69.
- Djeussi DE, Noumedem JAK, Kuete V, Ngadjui BT, Antibacterial and antibiotic-modulation activity of six Cameroonian medicinal plants against Gram-negative multi-drug resistant phenotypes, *BMC complementary and alternative medicine*, 2016 ; 16:124.
- Ngombe NK, Kalenda DT, Quetin-Leclercq J, Morel N, Vasoconstrictor and inotropic effects induced by the root bark extracts of *Anthocleista schweinfurthii*. *Natural product communications*, 2010; 5(3):369-72.
- Mbouangouere RN, Tane P, Ngamga D, Khan SN, Choudhary MI, Ngadjui BT, A new steroid and α -glucosidase inhibitors from *Anthocleista schweinfurthii*, *Research Journal of Medicinal Plant*, 2016; 1(3):106-111, DOI:10.3923/rjmp.2007.106.11.
- Abay G, Altun M, Koldas S, Tüfekci AR, Demirtas I, Determination of antiproliferative activities of volatile contents and HPLC profiles of *Dicranum scoparium* (Dicranaceae, Bryophyta), *Combinatorial Chemistry High Throughput Screening*, 2015; 18(5):453-63.
- Pinto MC, Tejeda A, Duque AL, Macias P, Determination of lipoxygenase activity in plant extracts using a modified ferrous oxidation. Xylenol orange assay, *Journal Agricultural and Food Chemistry*, 2007; 55(595):6-9.
- Delong JM, Prange RK, Hodges DM, Forney CF, Bishop MC, and Quilliam M, Using a Modified Ferrous Oxidation-Xylenol Orange (FOX) Assay for Detection of Lipid Hydroperoxides in Plant Tissue, *Journal of Agricultural and Food Chemistry*, 2002 ; 50(2):248-254.
- Oyedapo OO and Famurewa AJ, Antiprotease and membrane stabilizing activities of extracts of *Fagara Zanthoxyloides*, *Olax subscorpioides* and *Tetrapleura tetraptera*, *International journal of Pharmacognosy*, 1995 ; 33:65-69.
- Sakat SS, Juvekar AR, Gambhire MN, *In-vitro* antioxidant and anti-inflammatory activity of methanol extract of *Oxalis corniculata* Linn, *International Journal of Pharmacy and Pharmaceutical sciences*, 2010; 2(1):46-55.
- Irvine FR, *Woody Plants of Ghana*. Oxford University Press, London, UK.1961.
- Njayou FN, Moundipa PF, Tchana AN, Ngadjui BT, Tchouanguep FM, Inhibition of microsomal lipid peroxydation and oxideation by extracts from plants used in Bamum folk medicine (Cameroon) against hepatitis, *African journal of traditional, complementary and alternative medicines*, 2008; 278-289.
- Jegede IA, Ibrahim JA, Kunle OF, Phytochemical and pharmacognostic studies of the leaf and stem-bark of *Anthocleista vogelii* (Planch), *Journal of medicine plant research*, 2011; 5:6136-6139. <http://dx.doi.org/10.5897/jmpr10.404>.
- Leelaprakash G, Dass SM, *In-vitro* anti-inflammatory activity of methanol extract of *Enicostemma axillare*, *International Journal of Drug Delivery Research*, 2011; 3:189-196.
- Tabiri I, Henneh, Akrofi R, Ofori AE, Konja D, Owusu G, Abane B, Acquah-Mills J, Edzeameh JF and Tayman F, Stem Bark Extract of *Sterculia setigera* Delile Exhibits Anti-inflammatory Properties through Membrane Stabilization, Inhibition of Protein Denaturation and Prostaglandin E2 Activity, *International Journal of Pharmaceutical Research*, 2018 ; 22(5):1-11.
- Wisastra R, Dekker FJ, Inflammation, cancer and oxidative lipoxygenase activity are intimately linked, *Cancers (Basel)*, 2014; 6(1):500-521.
- Das SN, Chatterjee S. Long term toxicity study of ART-400, *Indian Journal of Indigenous Medicines*, 1995, 16(2):117-123.
- Mbiantcha M, Nguessom KO, Ateufack G, Oumar M, Kamanyi A, Analgesic properties and toxicological profile of aqueous extract of the stem bark of *Anthocleista vogelli* Planch, *International Journal of pharmaceutical, chemical and Biological Sciences*, 2013; 1:1-12.
- Baba H, Usifoh CO, Phytochemical investigation and anti-inflammatory property of ethanol-water extract of the roots of *Anthocleista djalonensis* A. Chev. (Gentianaceae). *African Journal of Biotechnology*, 2011; 10:6598-6600.
- Mulaudzi RB, Ndhlala AR, Kulkarni MG, Finnie JF, Staden Van J, Anti-inflammatory and mutagenic evaluation of medicinal plants used by Venda people against venereal and related diseases, *Journal of Ethnopharmacology*, 2013; 146(1):173-179.
- McGaw LJ, Elgorashi EE, Eloff JN, Cytotoxicity of African Medicinal Plants against Normal Animal and Human Cells, *Toxicological Survey of African Medicinal Plants*. Elsevier, London, 2014.
- Arya D and Patni V, Comparative analysis of *in vitro* anti-inflammatory and anti-arthritis activity in methanolic extract of *Pluchea lanceolata* Oliver & Hiern. *In vivo* and *in vitro*, *International Journal of Biological and Pharmaceutical Research*, 2013, 4(9):676-680.