


Available online on 15.05.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

Estimation of Metformin Hydrochloride in Bulk and Formulation by UV - Spectroscopic Area Under Curve Method

Kakade Vrushali B*, Lobhe G.A., Gandhat Snehah G., Dhabale S. M., Patel Salim G.

Vishal Institute of Pharmaceutical Education and Research, Ale, Tal- Junnar, Dist- Pune, (412411) Maharashtra, India

ABSTRACT

The current work is carried out for estimation of Metformin Hydrochloride by using area under curve (AUC) method using UV-visible spectrophotometer. For this purpose the wavelength range 221-241 nm was selected. Distilled water was used as a solvent throughout the work. Linearity was observed in concentration range 5-25 μ g/ml ($R^2 = 0.994$) for the method. The present method was found to be simple and linear which can be used for routine quality control analysis for spectrophotometric estimation of Metformin hydrochloride in bulk.

Keywords: Metformin Hydrochloride, Area under curve, Antidiabetic, λ_{max} .

Article Info: Received 24 March 2019; Review Completed 30 April 2019; Accepted 02 May 2019; Available online 15 May 2019



Cite this article as:

Kakade VB, Lobhe GA, Gandhat SG, Dhabale SM, Patel SG, Estimation of Metformin Hydrochloride in Bulk and Formulation by UV - Spectroscopic Area Under Curve Method, Journal of Drug Delivery and Therapeutics. 2019; 9(3):163-167
<http://dx.doi.org/10.22270/jddtv9i3.2632>

*Address for Correspondence:

Ms. Kakade Vrushali B., Department of M.Q.A., Vishal Institute of Pharmaceutical Education and Research, Ale, Tal- Junnar, Dist- Pune (412411) Maharashtra, India.

INTRODUCTION

Metformin hydrochloride is a hypoglycemic agent used in the treatment of non-insulin dependent diabetes mellitus. It acts as an antidiabetic agent by decreasing the uptake of glucose which enhance the glycemic control by increasing insulin sensitivity. Limitation associated with it is that, it does not show any response to dietary modification. It is basically a synthetic entity and chemically it is N,N-dimethylimidodicarbonimidic diamide hydrochloride (1,1-dimethyl biguanide hydrochloride). Generally used in combination with sulfonylureas, thiazolidinedione or other hypoglycemic agents or it can be used alone.¹⁻³




Figure 1: Chemical structure of Metformin HCl

MATERIAL AND METHOD

Chemicals:

Metformin hydrochloride was a gift sample from Flamingo Pharmaceutical, Taloja, Navi Mumbai, India. Metformin Hydrochloride Sustained Release Tablet 500 mg (Exermet 500) Cipla. All chemicals and reagents were of analytical (AR) grade.

Instrumentation:

A Shimadzu (Kyoto, Japan) UV-1800 double beam UV- Visible spectrophotometer attached with computer operated software UV probe 2.33 with spectral width of 2 nm, wavelength accuracy.

METHOD:

Area Under Curve Method:

The Area method is applicable where there is no sharp peak or when broad spectra are obtained. It involves the calculation of integrated value of absorbance with respect to the wavelength between two selected wavelengths λ_1 and λ_2 . Area calculation processing item calculates the area bound by the curve and the horizontal axis is selected by entering the wavelength range over which area has to be calculated. This wavelength range is selected on the basis of repeated observation so as to get the linearity between area curve and concentration. The above mentioned spectrums were

calculated to AUC. Thus, the calibration curve can be constructed by plotting concentration versus AUC.⁴⁻⁸

Experimental Work:

1. To check the solubility of Metformin Hydrochloride:

10 mg of Metformin was weighed and solubility of this sample was checked in distilled water, Methanol, Ethanol. In this Metformin HCL freely soluble in distilled water, slightly soluble in alcohol.⁴

2. To Identify the λ max of Metformin:

Accurately weigh 100 mg of pure drug and dissolve it in a small portion of distilled water and makeup the final volume up to 100 ml using distilled water to give a standard stock solution of 1000 μ g/ml. From above solution withdraw 10ml solution and transfer it into a 100ml volumetric flask and makeup final volume with distilled water to prepare 100 μ g/ml solution. Suitable dilutions made with distilled water to get working standard solutions of 5, 10, 15, 20, 25 μ g/ml.

3. Sample Preparation of Metformin:

Accurately weigh twenty tablets of metformin crush the tablets and measure the weight of powder equivalent to 100mg of drug. Then transfer it into a 100ml volumetric flask dissolve it into distilled water properly and make up the final volume with same distilled water to prepare 1000 μ g/ml solution. Then it filtered through whatman filter paper, pippetout 10ml filtrate transfer it into 100ml volumetric flask and makeup the volume up to 100ml with distilled water to prepare 100 μ g/ml solution. Suitable dilutions made with distilled water to get working standards of 5, 10, 15, 20, 25 μ g/ml.

Analytical Method Validation:⁵

1. Linearity:

Standard stock solution of Metformin, relevant amount of solution were pipette out into 25 ml volumetric flask and dilution were made with distilled water to be working standards of concentrations 5, 10, 15, 20, 25 μ g/ml. The difference in AUC of Metformin was measured area between 221.0-240.80 nm (Figure 6). The calibration curve of drug was plotted. The concentration range over which the drugs followed linearity was chosen as an analytical concentration range i.e 5-25 μ g/ml for Metformin.

2. Precision:

The precision of analytical procedure express closeness of agreement (degree of scattering) between a series of measurement obtained from multiple sampling of the same homogeneous sample under prescribed conditions. It may be considered at three levels: repeatability, intermediate precision and reproducibility. It is expressed as standard deviation or coefficient of variation.

3. Sensitivity:

The sensitivity of the method was determined in terms of limit of detection (LOD) and limit of quantitation (LOQ). The LOD and LOQ were calculated by using formula,

$$\text{LOD} = 3.3 \times \text{SD}/\text{slope}$$

$$\text{LOQ} = 10 \times \text{SD}/\text{slope}$$

Where, SD is the standard deviation of regression line.

RESULT AND DISCUSSION

The proposed UV- Spectroscopic, Area Under Curve method for determination of Metformin showed maximum absorbance at 233.00 nm (λ max). The result of UV-Spectroscopic analysis has been showed in Table.1 indicates that the representative calibration curve of the Metformin were plotted at 233.00 nm. A linear relationship was obtained for metformin in concentration range of 5-25 μ g/ml. Linear regression of absorbance on concentration gave the equation for Metformin,

$$\text{For Bulk, } y = 1.729x - 1.521$$

$$\text{Correlation coefficient (R}^2 = 0.992)$$

$$\text{For Formulation, } y = 1.356x - 1.024$$

$$\text{Correlation Coefficient (R}^2 = 0.994)$$

The area under curve of Bulk shown in Figure 5, and area under curve of formulation in figure 6.

Validation parameters results shown in Table.2 & 3. The overlay spectrum peak report shown in figure 7 and 8.

Table 1: Regression Analysis Data for Metformin by AUC Method

Parameter	AUC
Wavelength Range	221.0-240.80 nm
Concentration Range	5-25 μ g/ml
Slope (m)	1.3561
Intercept (c)	1.0245
Correlation Coefficient (R ²)	0.9941

Table 2: Summary of Data of Validation parameter

Sr. No.	Parameter	AUC Method
2	Linearity	5-25 μ g/ml
3	Regression Equation	$y = 1.356x - 1.024$
6	Correlation Co-efficient	$R^2 = 0.994$
7	LOD (μ g/ml)	0.0070
8	LOQ (μ g/ml)	0.0213
9	Precision	
9.1	Interday Precision	0.337% RSD
9.2	Intraday Precision	0.671% RSD

Table 3: Result of Intraday and Interday Precision

Parameter	\pm S.D	RSD	% RSD
Interday	0.002898	0.00167	0.337
Intraday	0.00145	0.00334	0.671


Figure 2: λ max of Metformin

Table 4: Calibration Curve of Pure Metformin (Bulk)

Concentration	Absorbance
5 ppm	0.325
10 ppm	0.720
15 ppm	1.073
20 ppm	1.469
25 ppm	1.782




Figure 3: Calibration Curve of Metformin (Pure Drug)

Table 5: Calibration Curve of Metformin (formulation)

Concentration	Absorbance
5 ppm	0.385
10 ppm	0.669
15 ppm	0.968
20 ppm	1.262
25 ppm	1.549




Figure 5: Calibration Curve of Metformin by AUC Method (Pure Drug)




Figure 4: Calibration curve of Metformin (Formulation)





Figure 4: Calibration Curve of Metformin by AUC Method (Formulation)

Spectrum Peak Area Report


15-04-2019 12:01:52

Data Set: File_190415_120047 - RawData


**Figure 5: Area between 223.40-240.00 nm selected for metformin (15 ppm) Bulk****Spectrum Peak Area Report**

15-04-2019 13:10:38

Data Set: File_190415_130904 - RawData

**Figure 6: Area between 221.0-240.80 nm selected for Metformin (15 μg/ml) Formulation****Overlay Spectrum Graph Report**

15-04-201 12:36:49

**Figure 7: Overlay Spectrum of Metformin (Bulk)**

Overlay Spectrum Graph Report

15-04-201 13:50:41




Figure 8: Overlay Spectrum of Metformin (Formulation)

CONCLUSION

The simple and economic UV spectrophotometric AUC methods have been developed and validated for the determination of Metformin Hydrochloride. The results of validation parameters show that the UV spectrophotometric method was found to be accurate, precise and sensitive. Because of cost- effective and minimal maintenance, the present UV-Spectrophotometric method can be preferred at small scale industries and successfully applied.

ACKNOWLEDGEMENT

The author are thankful acknowledge to Dr. Lobe .G. Head of Chemistry Department and Dr. Dhobale S.M., Head of Pharmaceutics Department Vishal Institute of Pharmaceutical Education and Research, Ale, Pune. For constant motivation and encouragement and also providing Metformin Hydrochloride drug as a gift sample from Flamingo Pharmaceuticals, Taloja, Navi Mumbai, India, for research work. We would like to thank our principle Dr. Jadhav S.L for providing us suitable environment for this work.

REFERENCES

- 1) Analytical Profiles of Drug Substances and Excipients Alison E. Bretnall, Graham S. Clarke, in Analytical Profiles of Drug Substances and Excipients, 1998
- 2) <https://en.m.wikipedia.org>
- 3) <https://pubchem.ncbi.nlm.nih.gov>
- 4) Indian Pharmacopoeia, Government of India, Ministry of Health and family welfare, Published by Indian Pharmacopoeial Commission, Ghaziabad, Volume-I, 161, Volume- II, 1657-1658.
- 5) ICH Harmonized Tripartite Guidelines: Validation Of Analytical method validation of analytical procedures: text and methodology Q2 (R1) . https://www.ich.org/fileadmin/public/website/ICH_products/Guidelines/QualityQ2-R1/Step4/Q2-R1_Guideline.pdf.
- 6) Beckett A.H, Stenlake J.B , Practical Pharmaceutical Chemistry, 4th edition, CBS publisher and Distributors, New Delhi, 275-295,2002.
- 7) Chistian G D, Analytical chemistry, six edition , John Wiley and sons, PA, 1-2,604,2003
- 8) Blessy M. A review, Development and stability indicating studies of drug, Journal of Pharmaceutical Analysis, 2013.
- 9) Saravanan V, Revathi R, Meera N, Method Development and Validation for the Simultaneous Estimation of Lycopene and Ubidecarenone By RP-HPLC in Combined Pharmaceutical Dosage Form. Journal of Drug Delivery and Therapeutics, 2016; 6(5):46-51.
- 10) Pachpute K, Jadhav S.L, Estimation of Disulfiram in Bulk and Formulation by UV- Spectrophotometric Area Under Curve method, Inventi Rpid:Pharma Analyis & Quality Assurance 2019; 1(2):1-6.
- 11) Chordiya BD, Patel SG, Pallavi PL, Area under curve UV-Specrophotometric Determination of Chlorcyclizine Hydrochloride in Bulk, Am.J. Pharm Tech Res.2019; 9(1):308-314.
- 12) Loni AB, Ghante MR, Sawant SD, Simultaneous UV-Spectroscopic Method for Estimation of Sitagliptin Phosphate and Metformin Hydrochloride in Bulk and Tablet Dosage , Scholar Research Library ,Der Pharm Chemica, 2012,4(3):854-859.
- 13) Skoog D.A, West, D.M. Holler F.J and Crouch S.R, In: Fundamentals of Analytical Chemistry, 5th edition, Thomas Books/cole, 971-995.