




Available online on 15.09.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

Anti-Inflammatory, Analgesic and Antipyretic Activity of *Catunaregam spinosa* (Thumb.) Tirveng Extracts

Hansa Saini ^{1*}, Jayesh Dwivedi ², Himanshu Paliwal ³, Udichi Kataria ⁴, Priyanka Chauhan², Rahul Garg ⁵

¹ Department of Pharmacy, Bhagwant University, Ajmer, Rajasthan (India)

² Pacific College of Pharmacy, Pacific University, Udaipur, Rajasthan (India)

³ Sunrise College of Pharmacy, Udaipur, Rajasthan (India)

⁴ Geetanjali Institute of Pharmacy, Udaipur, Rajasthan (India)

⁵ Venkateshwar Institute of Pharmacy, Udaipur, Rajasthan (India)

ABSTRACT

Catunaregam spinosa leaves have been ethnopharmacologically accounted for acquiring various pharmacological properties. The present study was undertaken to evaluate anti-inflammatory, analgesic and antipyretic potential of leaves of *C. spinosa*. The ethanolic extract was selected for this purpose based on phytochemical screening. Inflammation was inhibited at the dose of 200 mg/kg with percent inhibition of inflammation 32.06, 37.28 and 43.16 %, respectively, at 1, 3 and 5 h, while in egg albumin model % inhibition was found to be 47.81%. There was no significant analgesic activity seen in acetic acid induced writhing response method while significant effects were observed in the doses of 25 and 100 mg/kg on hot plate test. No antipyretic activity was shown by ethanolic extracts(25, 100 and 200 mg/kg) against Brewer's yeast induced pyrexia in rats.

Keywords: *Catunaregam spinosa*, Anti-inflammatory activity, Phytochemical screening, Ethanolic extract.

Article Info: Received 10 June 2019; Review Completed 18 Aug 2019; Accepted 29 Aug 2019; Available online 15 Sep 2019



Cite this article as:

Saini H, Dwivedi J, Paliwal H, Kataria U, Chauhan P, Garg R, Anti-Inflammatory, Analgesic and Antipyretic Activity of *Catunaregam spinosa* (Thumb.) Tirveng Extracts, Journal of Drug Delivery and Therapeutics. 2019; 9(5):89-94
<http://dx.doi.org/10.22270/jddt.v9i5.3363>

*Address for Correspondence:

Ms. Hansa Saini (M. Pharm.), Sikar Road, Ajmer-305004, Rajasthan (INDIA)

INTRODUCTION

Ever since ancient times, in search of rescue for their disease, the people looked for drugs in nature. According to the World Health Organisation (WHO), more than 80% of the World's population relies on traditional herbal medicine for their primary health care needs. In last few years interest of researches in medicinal plants has increased significantly. Identification and validation of plant derived substances for the management of several diseases is the goal of current researchers. It is estimated that about 25% of the western medicines are directly or indirectly derived from plant sources¹.

Catunaregam spinosa (Thumb.) Tirveng is also known as a *Randia dumetorum* belongs to family Rubiaceae. It occurs in almost throughout India up to 4,000 ft altitude^{2,3}. *C. spinosa* is known to show alexiteric and antipyretic activity with also significant in skin diseases, inflammations, ulcers, wounds and tumours like diseases⁴. It contains secondary metabolites like triterpenoidal saponins, tannins, essential

oil, resin and veleric acid. A number of Phytoconstituents are reported like D-mannitol, 1-keto-3 α hydroxy Oleanane, saponins, scopoletin, iridiodal and glycosides, such as garenoside, randidoside and geniposide. Also, steroids and fatty acid such as caprylic acid, capric acid, tauric acid, myristic acid has been reported in literature^{5,6}. The main aim of the current study is to explore presence of phytochemical and secondary metabolites profile analysis of *C. spinosa* leaves and investigate the anti-inflammatory, analgesic and antipyretic activity of the plant.

METHODOLOGY

Collection of plant material and authentication

The leaves of the *C. spinosa* were collected and dried. The dried material was powdered in the coarse powder by mechanical grinder. The resulting coarse powder was used for the studies. The leaves of the plant were authenticated by the Dr. Dilip Gena, Botanist, Pteridophyte biology lab Department of Botany, Government College, Ajmer.

Extraction procedure for collected plant material

The extract of leaves of *Catunaregam spinosa* was obtained different types of solvents like, Methanol, Ethanol, Acetone, Petroleum ether, Benzene and Distilled water. 10 gm of powdered leaves of *C. spinosa* were subjected Soxhlet extraction and maceration method using different solvents⁷. The macerated extract was kept on the shaker for 24 hours, later it was centrifuged and the supernatant was taken for the excess solvent evaporation in laboratory conditions. After evaporation of excess solvent the crude extract was stored in refrigerator till further analysis.

Phytochemical analysis of *Catunaregam spinosa*

Phytochemical tests for the identification of amino acids, carbohydrates, saponins, tannins, phytosterols, alkaloids, proteins, glycosides, flavonoids and phenolic compounds were carried out for all the extracts. The present investigation was planned with an objective to establish Pharmacognostic standards and to evaluate preliminary phytochemical data that can facilitate the authentication and the isolation of the desired constituent from the correct extract⁸.

Characterization of extract using Fourier Transform Infrared Spectrophotometer

FTIR is the most powerful and applicable tool for identification of chemical bonds (functional groups), and their types present in sample. The wavelength which absorbed light in different variation is characteristic and predictable information for the chemical bond which can be seen in the annotated spectrum of FTIR. By interpreting the infrared absorption spectrum, the chemical bonds present in molecule can be determined. Dried powder of plant sample for different solvents extracts were used in FTIR analysis with 100 mg KBr pellet as encapsulate in sample discs. The powdered plant part sample was loaded in FTIR spectroscope (Shimadzu, IR Affinity 1, Japan), with the Scanning range from 400 to 4000 cm⁻¹ with a 4 cm⁻¹ resolution^{9,10}.

Anti-inflammatory activity

Carrageenan induced rat paw edema

The anti-inflammatory activity of ethanolic extract of *C. spinosa* was determined by carrageenan induced inflammation test in hind paw of rats. Rats were fasted for 24 h before the commencement of experiment. Edema was induced by injecting 0.1 ml of 1% (w/v) carrageenan in saline. Rats were divided into different treatment groups (n = 5) viz., group I (vehicle, 10 mL/kg), group II (indomethacin, 10 mg/kg), group III (Ethanolic extract, 100 mg/kg), group IV (Ethanolic extract, 200 mg/kg). The extracts and indomethacin (IM) were suspended in 1% CMC and administered orally by intragastric tube 1 h before the carrageenan injection. After the administration of the phlogistic agent, paw volume (in mm) of individual rat was measured at 1, 3 and 5 h¹¹.

Egg albumin induced rat paw edema

This method has been applied in the screening of antiinflammatory activity of African spices and herbs by some researchers. Egg albumin is reported to induce inflammation in the hind paw of rats and therefore, this model was employed to evaluate the anti-inflammatory activity of extracts. Acute inflammation was induced by

injecting 0.1 ml/kg of fresh egg albumin into the plantar region of the hind paw of rats. Separate sets of rats (n = 5) were employed for control, standard and treatment groups as described in carrageenan induced inflammation experiment. The change in paw volume (mm) was measured up to 120 min, at 20 min intervals after egg albumin injection^{12,13,14}.

Analgesic activity

Analgesic activity was tested in mice (King Instt. Strain) weighing between 20-25 g with six number of animals in each group using Acetic acid induced writhing response and Hot plate response methods.

Acetic acid induced writhing response

Different groups of six mice each received orally normal distilled water (2 ml/kg) (i.e. control), Ethanolic extract of *C. spinosa* (25, 100 and 500 mg/kg) and Analgin (500 mg/kg) (i.e. standard). Thirty minutes later, 0.7% acetic acid (10 ml/kg) solution was injected intraperitoneally to all the animals in different groups. The stretching episodes of individual mice were counted for 30 min post-administration of acetic acid. Analgin in a dose of 500 mg/kg orally served as standard. The formula used for computing percent inhibition was: average writhes in the control group minus writhes in the drug group divided by writhes in the control group times 100. Compounds with less than 70% inhibition are considered to have minimal antinociceptive activity¹⁵.

$$\% \text{ Inhibition} = [(W_c - W_t) \times 100] / W_c$$

Where, W_c = average number of writhes in control c group; W_t = average number of writhes in test t group

Hot-plate response in mice

Mice were retained on a hot plate having a stable temperature of 55 ± 1°C. The time taken for either paw licking or jumping was recorded. Each mouse was individually placed on the hot plate in order to find the animal's reaction to electrical heat-induced pain (licking of the forepaws and eventually jumping). The latency until mice showed first signs of discomfort (hind paw lifting, hind paw licking, or jumping) was recorded, before (baseline), and response was determined at 30, 60, 90, 120, 150 and 180 min after the administration of *C. spinosa* (25, 100, and 500 mg/kg), and analgin (500 mg/kg)¹⁶.

Anti-pyretic activity

The rats were made pyretic by a sub-cutaneous injection of 20% suspension of dried Brewer's yeast in normal saline in a dose of 1 ml/100 g. The rectal temperature was recorded by introducing a clinical thermometer about 2 cm deep into the rectum for the duration of one min before and 4 h after yeast injection. Animals showing pyrexia were selected and the drugs were administered. The rectal temperature was recorded at hourly intervals for 5 h. Animals that indicated a rise in body temperature of at least 0.6 °C were considered eligible for carrying out further experiments.

RESULTS AND DISCUSSION

Phytochemical analysis of *Catunaregam spinosa*

The results of phytochemical analysis of leaves of *C. spinosa* are tabulated below in Table.1. The ethanolic extract was selected for further investigations based on the results obtained phytochemical analysis.

Table.1: Phytochemical analysis of different extracts of *Catunaregum spinosa* leaves

S. No.	Name of the constituent	(P)	(B)	(C)	(A)	(E)	(M)	(Aq)
1	Alkaloids	—	—	—	—	—	—	—
2	Glycosides	—	—	—	+	+	+	—
3	Carbohydrates	—	—	—	+	+	+	+
4	Phytosterols/ Triterpenoids	+	+	+	+	+	+	—
5	Proteins & Amino acids	—	—	—	—	—	—	+
6	Saponins	—	—	—	—	+	+	+
7	Flavonoids	—	—	—	+	+	+	—
8	Fixed oils & Fats	+	—	—	—	—	—	—
9	Gums/Mucilage	—	—	—	—	—	—	—
10	Volatile oil	+	—	—	—	—	—	—
11	Phenolics/Tannins	—	—	—	+	+	+	+



(P):Petroleum ether (60-80°C); (B):Benzene; (C): Chloroform; (A): Acetone; (M): Methanol; (E): Ethanol; (Aq): Aqueous; +: Positive; -: Negative

FTIR Spectral Analysis

The methanolic extract of *C. spinosa* leaves showed typical absorption bands at 3295 cm⁻¹ for a hydroxyl (-OH) group 2829 cm⁻¹, 2413 cm⁻¹ (for C-H stretching), 1386 cm⁻¹ (for C-H bending), and at 1620 cm⁻¹ for C=C group. The ethanolic extract illustrated the observation of absorption bands at 2919 cm⁻¹ (for C-H stretching), 1433 cm⁻¹ (for C-H bending) for C-H group and at 1709 cm⁻¹, for a carbonyl group (C=O).

Table.2: FTIR spectral peak values and functional groups obtained for the leaf extract

Extracts	Peak values	Functional groups
Methanol	1386	C-H bending
	1620	C=C group
	2413	C-H stretching
	2829	C-H stretching
	3295	-OH group
Ethanol	1433	C-H bending
	1614	C=C group
	1709	C=O carbonyl group
	2919	C-H stretching
	3301	-OH group

**Figure.1: FTIR of *C. spinosa* Leaves in Methanol****Figure.2: FTIR of *C. spinosa* Leaves in Ethanol**

Anti-inflammatory activity**Anti-inflammatory activity against carrageenan- induced inflammation**

Ethanol extract of *C. spinosa* at the dose of 100 mg/kg inhibited only the initial phase of inflammation, while initial as well as lateral phases of inflammation were inhibited at the dose of 200 mg/kg with percent inhibition of inflammation 31.69, 35.11 and 41.62 %, respectively, at 1, 3 and 5 h. Standard drug, indomethacin attenuated the inflammation at 10 mg/kg with 18.97, 57.72 and 43.53% inhibition, respectively, at 1, 3 and 5 h (Table.3). These results were found to be dose dependent.

Anti-inflammatory activity against egg albumin- induced inflammation

To substantiate the anti-inflammatory activity of the extract, the efficacy was also studied against egg albumin-induced inflammation (Table.4). The ethanol extract (100 mg/kg, p.o.) failed to exhibit anti-inflammatory effect ($p > 0.05$) at 20 and 40 min time interval (initial phase) but showed significant inhibition of inflammation at 60, 80, 100 and 120 min (lateral phase) after egg albumin administration as compared to control. On the other hand, the extract at higher dose (200 mg/kg, p.o.) significantly inhibited ($p < 0.05$) egg albumin-induced inflammation at all time interval (initial as well as lateral phase).

Table.3: Anti-inflammatory activity of Ethanol extract of *C. spinosa* in carrageenan-induced inflammation model

Treatment	Dose (mg/kg)	Edema inhibition against carrageenan-induced inflammation (h)		
Measurement time interval →		1	3	5
Control		6.23 ± 0.13	7.45 ± 0.14	6.50 ± 0.19
Ethanol extract	100	5.10 ± 0.21 [17.30] #	7.11 ± 0.15 [4.45] ns	6.33 ± 0.10 [3.35] ns
	200	4.26 ± 0.25 [32.06] #	4.83 ± 0.18 [37.28] #	1.79 ± 0.11 [43.16] #
Indomethacin	10	5.05 ± 0.17 [18.97] #	3.15 ± 0.15 [57.72]	3.67 ± 1.02 [43.53] #

Table.4: Anti-inflammatory activity of Ethanol extract in egg- albumin induced inflammation model

Treatment	Dose (mg/kg)	Edema inhibition against egg albumin-induced inflammation (min)					
Measurement time interval →		20	40	60	80	100	120
Control		5.45 ± 0.12	5.96 ± 0.23	6.62 ± 0.17	7.67 ± 0.21	7.30 ± 0.33	7.14 ± 0.19
Ethanol extract	100	5.28 ± 0.23 [3.11] ns	5.76 ± 0.26 [3.45] ns	6.31 ± 0.32 [4.60] *	5.27 ± 0.34 [31.30] #	4.86 ± 0.23 [33.29] #	5.32 ± 0.22 [25.59] #
	200	4.16 ± 0.15 [21.44] #	4.69 ± 0.16 [20.94] #	5.41 ± 0.16 [18.12] #	5.43 ± 0.27 [30.02] #	4.80 ± 0.21 [33.94] #	3.69 ± 0.11 [47.81] #
IM	100	4.99 ± 0.13 [8.41] #	5.27 ± 0.12 [11.58] #	5.53 ± 0.23 [16.43] #	5.76 ± 0.16 [24.96] #	3.93 ± 0.18 [46.15] #	3.25 ± 0.14 [54.48] #

Values represent the mean±SEM. of five animals for each group; Values in square brackets indicate the percentage inhibition rate of inflammation. * $p < 0.01$ and # $p < 0.001$ indicate different levels of statistically significant values against control and ns stands for non significant. % Inhibition calculated using formula: paw edema volume control-sample/control.

Analgesic activity

The compound did not show significant analgesia on stretching episodes induced by acetic acid in any dose (Table.5), whereas, it showed significant effects in the doses of 25 and 100 mg/kg on hot plate test. The analgesia

produced by the former doses was for a brief period (between 60 to 90 min) whereas, the effect produced by 100 mg/ kg was more prolonged and with an earlier onset comparable to the effect produced by Analgin in a dose of 500 mg/kg (Table.6).

Table.5: Effect of extract of *C. spinosa* on acetic acid induced stretching episodes in mice

(Values are Mean ± S.E., Number of observations in parenthesis)

Groups	Dose in mg/kg oral	Stretching episodes for 30 min	% Inhibitory activity	P-value
Control (6)	Distilled water	36.33 ± 5.77	---	---
Analgin (6)	500	3.16 ± 1.19	91.31	<0.001
Extract (6)	25	40.16 ± 8.19	---	ns
Extract (6)	100	46.16 ± 9.33	---	ns
Extract (6)	200	26.00 ± 4.46	28.44	ns

Table.6: Analgesic Activity of extract of *C. spinosa* on Hot plate in Mice(Figures are Mean \pm S. E., Number of observations in parenthesis)

Groups	Dose in mg/kg oral	Initial response time (sec)	Response time in seconds after					
			30 min	60 min	90 min	120 min	150 min	180 min
Analgin (6)	500	5.05 \pm 0.48	7.41 \pm 1.21 ^{ns} p<0.001	14.46 \pm 1.83 1.24 p<0.001	18.55 \pm 1.09 0.96 p<0.001	19.14 \pm 1.09 0.96 p<0.001	14.63 \pm 1.87 0.96 p<0.001	7.55 \pm 0.88 0.94 p<0.05
Extract (6)	25	5.01 \pm 0.68	6.51 \pm 0.45 ^{ns} p<0.02	8.80 \pm 1.20 0.96 p<0.02	7.18 \pm 0.29 0.96 p<0.01	6.86 \pm 1.07 ^{ns} 0.96 p<0.01	5.96 \pm 0.74 ^{ns} 0.91 ^{ns} p<0.01	6.51 \pm 0.64 ^{ns} 0.94 p<0.05
Extract (6)	100	5.11 \pm 0.32	7.20 \pm 0.53 0.53 p<0.001	7.38 \pm 0.96 0.96 p<0.01	7.10 \pm 1.06 0.96 p<0.01	6.75 \pm 0.86 0.86 p<0.01	5.41 \pm 0.91 ^{ns} 0.91 ^{ns} p<0.01	6.20 \pm 0.94 0.94 p<0.05
Extract (6)	200	5.21 \pm 0.65	7.63 \pm 0.98 ^{ns} 0.98 ^{ns}	6.11 \pm 1.11 ^{ns} 1.11 ^{ns}	7.75 \pm 1.06 ^{ns} 1.06 ^{ns}	6.48 \pm 0.73 ^{ns} 0.73 ^{ns}	6.35 \pm 0.41 ^{ns} 0.41 ^{ns}	5.43 \pm 0.55 ^{ns} 0.55 ^{ns}

ns stands for non significant.

Antipyretic activity

There was no 'antipyretic effect observed in this model of experiment in any of the doses employed (Table.7).

Table.7: Effect of extract of *C. spinosa* On Brewer's Yeast Induced Pyrexia in Rats(Values are Mean \pm S. E., Number of animals in parenthesis)

Groups	Body Weight (in g)	Fasting rectal temp. (°F)	Initial Pyrexia	Mean rectal temperature in °F at				
				1 st h	2 nd h	3 rd h	4 th h	5 th h
Control(6)	131 \pm 6.57	99.45 \pm 0.05	102.75 \pm 0.29	102.95 \pm 0.13	103.10 \pm 0.17	103.70 \pm 0.33	103.85 \pm 0.20	103.26 \pm 0.27
Paracetamol 500 mg/kg(6)	163.75 \pm 8.50	99.30 \pm 0.15	102.35 \pm 0.24	100.85* \pm 0.11	99.70* \pm 0.21	100.20# \pm 0.63	101.25# \pm 0.25	101.80 \pm 0.21 ^{ns}
Extract 25 mg/kg (6)	148.75 \pm 13.90	99.30 \pm 0.23	102.80 \pm 0.25	102.50 \pm 3.28 ^{ns}	102.00 \pm 3.28 ^{ns}	102.85 \pm 0.26 ^{ns}	103.85 \pm 0.65 ^{ns}	102.25 \pm 0.41 ^{ns}
Extract 100 mg/kg (6)	111.25 \pm 9.43	99.05 \pm 0.22	102.60 \pm 0.29	103.00 \pm 0.25 ^{ns}	103.40 \pm 0.14 ^{ns}	102.00 \pm 0.80 ^{ns}	103.05 \pm 0.17 ^{ns}	103.60 \pm 0.42 ^{ns}
Extract 200 mg/kg(6)	125.00 \pm 8.89	98.90 \pm 0.17	101.85 \pm 0.62	102.50 \pm 0.97 ^{ns}	102.20 \pm 0.39 ^{ns}	101.95 \pm 1.01 ^{ns}	102.50 \pm 0.4 ^{ns}	101.10 \pm 2.03 ^{ns}

CONCLUSION

The multifunctional properties of *C. spinosa* exposed in previous studies about its antioxidant and anticancer activities might be useful in explaining ethnopharmacological claim of its therapeutic potential in inflammation, pain and fever, showing no signs of oral toxicity.

The extract of *C. spinosa* showed significant anti-inflammatory effect both in exudative and proliferative phases of inflammation. In carrageenin induced edema, the effect produced by the dose of 100 mg/kg was little lesser than Indomethacin in the dose of 10 mg/kg with further higher increase in anti-inflammatory activity with higher dose of 200 mg/kg. In case of egg-albumin induced inflammation, dose of 200 mg/kg showed anti-inflammatory activity which was comparable to Indomethacin (10mg/kg).

The compound did not show significant analgesia in chemical writhing test whereas, on hot-plate, significant effect was observed in the doses of 25 and 100 mg/kg. The analgesia produced by the former doses was for a brief period (between 60 to 90 min) whereas, the effect produced by 100 mg/ kg was more prolonged and with an earlier onset comparable to the effect produced by Analgin in a dose of 500 mg/kg.

The compound is devoid of antipyretic effect in this model of experiment. The maximum anti-inflammatory and analgesic

activities are almost around the dose of 200 mg/ kg which certainly may make it interesting to study further in detail therefore it could be clinically tried.

REFERENCES

1. Pallab naity,Dipak Mishra et al, Biological activities of crude extracts and chemical constituents of Bael, *Aegle marmelos* (L.), Indian Journal of experimental biology : November 2009, 47, pp. 849-861.
2. Vaidya. V. M. Gogte, Ayurvedic Pharmacology and Therapeutic Uses of medicinal Plants(Dravyaguna vignyan) Part II, medicinal plants, Plant no.(87), First English Edition, Bhartiya Vidya Bhavan, Oct 2000, Pg 454.
3. Kirtikar K.R., Basu B.D. Indian Medicinal Plants. Panni office,Bhuwaneswari Ashrama, Bahadurganj, Allahabad. (1991) 648-652.
4. Agrawal SS, Singh VK, Immuno modulators-A review of studies on Indian medicinal plants and synthetic peptides, Part- 1, Medicinal plants, Proc Indian Natl Sci Acad, 62, 1999, 179-204.
5. Sharma PC, Yelne MB, Denis JJ, Data base on Medicinal plants used in Ayurveda, Central Council for Research in Ayurveda & Siddha, 2000, 380-383.
6. Iyengar MA, Bibliography of Investigated Indian Medicinal Plants, (1950-1975), Manipal, Manipal Power Press, 1976, 1-144.
7. World Health Organisation. 1998; Quality Control Methods for Medicinal Plant Materials, WHO, Geneva.

8. Harborne JB. Phytochemical methods, *Chapman and Hall*, London (1998), Edn 3, 90 - 203.
9. Yang J, Yen HCE, Early Salt Stress Effects on the Changes in Chemical Composition in Leaves of Ice Plant and Arabidopsis, A Fourier Transform Infrared Spectroscopy Study, *Plant Physiology*, 130, 2002, 1032-1042.
10. Martin JA, Solla A, Woodward S, Gil L, Fourier transform-infrared spectroscopy as a new method for evaluating host resistance in the Dutch elm disease complex, *Tree Physiology*, 25, 2005, 1331-1338.
11. Winter C.A., Risley E.A., Nuss G.W., *Proc. Soc. Exp. Biol. Med.* 111, 544 (1962).
12. Otimenyin, S.O., Uguru, M.O., Auta, A., 2008. Anti-inflammatory and analgesic activities of Cassia goratensis and *Sacracephalus esculentus* extracts. *J. Herbs Spices Med. Plants* 13, 59-67.
13. Chinasa, E.C., Ifeoma, I.A., Obodoike, E.C., Chhukwuemeka, E.S., 2011. Evaluation of anti-inflammatory property of the leaves of *Sansevieria liberica* ger. and labr. (fam: Dracaenaceae). *Asian Pac J Trop Med* 4, 791-795.
14. Winter C.A., Risley E.A., Nuss G.V., *J. Pharmacol. Exp. Ther.* 141, 369 (1963).
15. Turner R. A. (1965a). Non-narcotic analgesics. In Screening Methods in *Pharmacology*. Vol. 1, pp. 114-115, New York : Academic Press.
16. Eddy, N.B. and Leimbach, D.J., (1953) Synthetic analgesics. II. Dithie-nylbutenyl- and dithienylbutylamines. *Journal of Pharmacology and Experimental Therapeutics*, 107, 385-393.

