

Available online on 30.08.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited

Open  Access

Review Article

Biological Activities and Phytochemical Constituents of Trailing Daisy Trilobata: A Review

Sai Prasanna K^{*1}, Jyothi Reddy G¹, Kiran M¹, Thyaga Raju K²

1. Department of Pharmacology, SVU College of Pharmaceutical Science, SV University, Tirupati - 517501, Andhra Pradesh, India.

2. Department of Biochemistry, SVU College of Pharmaceutical Science, SV University, Tirupati - 517501, Andhra Pradesh, India.

ABSTRACT

The traditional system of medicinal plants have been found to possess significant anti-inflammatory, antibacterial, anti-fungal, anti-diabetic, analgesic properties etc. Plant-derived drugs are used to cure mental illness, skin diseases, tuberculosis, diabetes, jaundice, hypertension, and cancer. *Wedelia Trilobata* belongs to family Asteraceae. Leaf, stem, and flower of *Wedelia trilobata* show anti-microbial, anti-oxidant and anti-inflammatory activity, analgesic activity. Phytochemical screening of the extract has been reported to show the presence of tannins, cardiac glycosides, flavonoids, terpenoids, phenols, saponins, and coumarins. *Wedelia Trilobata* is also used in reproductive problems, amenorrhea, chest cold, dry cough, and fever. The present review aims to the study was phytoconstituents, biological and pharmacological activities of *Wedelia trilobata*. This study suggested a possible use of *Wedelia trilobata* as a source of natural medicines as an anti-inflammatory, anti-oxidant, anti-microbial, hepatoprotective, antidiabetic agents.

Keywords: *Wedelia Trilobata*, Trailing daisy trilobata, Complaya trilobata(L), Sphagneticola Trilobata, pharmacological review.

Article Info: Received 16 June 2019; Review Completed 13 August 2019; Accepted 17 August 2019; Available online 30 August 2019



Cite this article as:

Sai Prasanna K, Jyothi Reddy G, Kiran M, Thyaga Raju K, Biological Activities and Phytochemical Constituents of Trailing Daisy Trilobata: A Review, Journal of Drug Delivery and Therapeutics. 2019; 9(4-s):888-892
<http://dx.doi.org/10.22270/jddt.v9i4-s.3578>

*Address for Correspondence:

Sai Prasanna. K, Department of Pharmacology, SVU College of Pharmaceutical Science, SV University, Tirupati - 517501, Andhra Pradesh, India.

Introduction:

The introduction of plant-derived drugs in modern medicine has been linked to the uses of plant-derived materials as an indigenous cure in the traditional system of medicine. Some of the plants have been used for the treatment of antibacterial, antifungal, anticancer, anti-diuretic, anti-inflammatory, anti-diabetic medicines at a lower cost and lesser side effects than marketed drugs. Various chemical constituents have been isolated from medicinal plants. Some plant-derived drugs have been used to various diseases like mental illness, skin diseases, tuberculosis, diabetes, jaundice, hypertension, cancer etc¹.

Wedelia trilobata belongs to family Asteraceae is a highly resistant, creeping, perennial plant, with succulent stems. *Wedelia trilobata* is also known as Trailing daiay trilobata². It has a very old tradition of wide medicinal use it a high reputation in Ayurveda, Unani, Siddha, and Traditional Chinese medicine and also in traditional systems of healing in the Carribean, Central and South America. The extracted essential oil of *Wedelia trilobata* has been found to act itself as an antioxidant, antibacterial, antifungal, anti-inflammatory, cough relieving agent, hepatoprotective,

febrifuge, immuno-stimulatory and analgesic agent. The present review aims to address the ethnopharmacological uses, phytochemical constituents and reported pharmacological activity of Trailing daisy trilobata³.

Description of *Wedelia trilobata*:

Habitat: A weed of urban bushland, closed forest margins, open woodland, waterways, lake margins, wetlands, roadsides, disturbed sites, waste areas, vacant lots and coastal sand dunes in tropical and sub-tropical regions. It may also encroach into lawns, footpaths, and parks from nearby gardens.

Geographical Distribution: Native to India, A.P, Chittor district, Mexico, Central America (i.e. Belize, Costa Rica, Guatemala, Honduras, Nicaragua and Panama) the and throughout the Caribbean, where it is noted as a weed in Trinidad, Puerto Rico, the Dominican Republic, Jamaica, Panama and tropical South America(i.e. French Guiana, Guyana, Surinam, Venezuela, Brazil, Bolivia, Colombia, Ecuador, and Peru). Naturalized in South Africa, Florida, Louisiana, Hawaii, Puerto Rico, and the Virgin Island. Escaped in many tropical regions of the world including

Australia (south-eastern Queensland and north-eastern New South Wales). The Pacific Island (i.e. American Samoa, the Cook Island, Fiji, French Polynesia, Guam, Kiribati, the Marshall Islands, Nauru, Niue, New Caledonia, Palau, Western Samoa, Tonga, and Hawaii), Malaysia, Indonesia, Thailand, India, Papua New Guinea.

Taxonomical Classification:

Domain: Eukaryota
 Kingdom: Plantae
 Phylum: Spermatophyta
 Subphylum: Angiospermae
 Class: Dicotyledonae
 Order: Asterales
 Family: Asteraceae
 Genus: *Sphagneticola*
 Species: *Sphagneticola trilobata*

Vernacular Names:

Telugu: Guntagalagara.
 English: Bay Biscayne creeping-oxeye.
 Hindi: Pilabhangara, Bhangra.
 Tamil: Manjalkarilamkanni, patalai kavyantakara.
 Malayalam: Mannakkannunni.
 Kannada: Gargari, kalsarji.

Synonyms:

Wedelia trilobata (Rich.) Bello
Complaya trilobata (L.)
Strother, *Silphium trilobatum* L.,
Thelechitonia trilobata (L.) H. Rob. & Cuatrec.,
Wedelia carnosa Rich.,



Figure: 01 *Sphagneticola trilobata*

Pharmacological Activities

1. Anti-inflammatory Activity:

Meena *et al.* (2011) investigated four herbal drugs used in traditional Medicine including *Wedelia trilobata* on the anti-inflammatory activity. They reported that all the extracts reduced croton oil-induced ear dermatitis. Results suggested that lipophilic extracts are potential source was found to possess have significant anti-inflammatory activity⁴.

Maldini *et al.* (2009) studied anti-inflammatory activity using in-vitro models such as albumin denaturation and heat-induced hemolysis. An ethanol extract of *Wedelia trilobata* demonstrated anti-inflammatory activity, vice suppression of lipopolysaccharide-induced cytokine and nitric oxide production in peripheral blood mononuclear cells⁵.

2. Anti-microbial:

Maori *et al.* (2009) conducted a study on Anti-microbial activity of *Wedelia trilobata*, by using Disc- diffusion method. The activity of curd extract from *Wedelia trilobata* against gram-positive, gram-negative bacteria, yeast, and fungi was evaluated. The n-hexane extract showed antibacterial activity against *Bacillus subtilis*, *Mycobacterium smegmatis*, and *Pseudomonas aeruginosa*, *Salmonella* group C, *S Paratyphi* and *Shigella sonnei*. The aqueous extract was active only against salmonella group C, inactive against other tested bacteria. None of the tested extracts showed biological activity against yeast and fungi⁶.

Mandal *et al.* (2005) conducted a study on the antimicrobial activity of *Wedelia trilobata*. Aqueous extract of leaf, stem, flower inhibited the growth of all the bacterial isolates, but the extracts did not show any similar significant effect on fungi isolates. The leaf extract showed more potent activity against *Pseudomonas aeruginosa*, *E.coli*, *Pseudomonas fluorescens*, *Xanthomonas oryzae* *Proteus Vulgaris.oryzae*, *Xanthomonas axanopodis* *Proteus Vulgaris.malvacearum* moderately inhibited the *Clavibacter minchiganensis* sub sp. *Minchiganensis* but less activity was observed on *Staphylococcus aureus*. All the extracts exhibited less activity on all species of *Fusarium* and *Aspergillus*⁷.

3. Analgesic:

Suresh Kumar *et al.* (2007) conducted a study on *Wedelia trilobata* by acetic acid-induced writhing method and hot plate assay to assess analgesic activity in mice. It was found that the extract caused an inhibition of the writhing response induced by acetic acid a dose depended manner (500mg/kg). The results reflect analgesic effects and the therapeutic efficacy of the extract in animal models was comparable with that of standard drugs such as aspirin and morphine⁸.

Teerapol srichana *et al.* (2014) conducted a study in mice on the analgesic activity of ethanol extract of *Wedelia trilobata* by the acetic acid-induced writhing method. The extract showed dose depended (500mg/kg) blocking of writhing response. kaurenoid acid(10mg/kg) obtained from *Wedelia trilobata* inhibited over nociception, acute carrageenin, & PGE2 induced complete Freund's adjuvant (CFA) and chronic induced mechanical hyperalgesia. The results reflect analgesic effects and therapeutic efficacy of the extract on animal models which are comparable with that of standard drug aspirin⁹.

4. Hepatoprotective activity:

Karamagam *et al.* (2008) conducted a study on ethanolic leaf extract of *Wedelia trilobata* against carbon tetrachloride CCl₄ induced acute hepatotoxicity in rats. There was shown a significant reduction (P<0.05) in the levels of protein, bilirubin and organ weight including liver, heart, left lung, spleen, and kidney without any significant changes in body weight. The treatment of ethanol leaf extract showed a dose-dependent reduction of CCl₄ induced elevated serum levels of enzyme activities with a parallel increase in total protein & bilirubin¹⁰.

Lin *et al.* (1994) investigated the hepatoprotective effect of *Wedelia trilobata*. Acute hepatitis was induced by three hepatotoxins, such as carbon tetrachloride, acetaminophen

in mice and galactosamine in rats after treatment with *W. trilobata* (300 mg/kg) a reduction in the elevated serum glutamate oxaloacetate transaminase and glutamate pyruvic transaminase levels was observed at 24 hours after hepatotoxins were administration. It was concluded that *Wedelia trilobata* has a definite hepatoprotective effect against liver injuries¹¹.

5. Anti-diabetic:

Sunita Kanikaram *et.al.* (2018) determined the anti-diabetic activity of crude extracts of *Wedelia trilobata* along with *Brassica oleraceae* by applying in vitro α -amylase inhibition method. The methanol extract of *Wedelia trilobata* significantly showed DPPH inhibition with IC₅₀ 20 μ g/mL and

in vitro α -amylase inhibition IC₅₀ 50 μ g/mL. It was concluded that methanol extract of *Wedelia trilobata* can be a potential source of anti-oxidant and as good anti-diabetic¹².

Pradeep *et al.*(2014) conducted a study on methanolic extract of *Wedelia trilobata* using in- vitro glucose diffusion, α -amylase, α -glycosidase & angiotensin I converting enzyme inhibition method. They showed an inhibitory effect on the α -glucosidase enzyme. Glucose movement from sealed dialysis tube to an external solution was inhibited by *Wedelia trilobata* extract. The methanol extract of *Wedelia trilobata* inhibited the rabbit lung angiotensin I converting ¹³ enzyme with IC₅₀ of 30 μ g/mL. Biological and pharmacological activities are explained in Table.2.

Table 2: Biological and pharmacological activities reported

Activity	Method	Parameters	Reference
1. Anti inflammatory activity	a. In vitro Heat-induced albumin denaturation.	Albumin denaturation	GovindappaM.Naga (<i>et.al</i> 2011) ¹⁴ .
	b. In vivo Croton oil-induced ear oedema in rats.	Inhibition of plasma extravasations and neutrophil migration	Giovanafucina(<i>et.al</i> 2016) ¹⁵ .
	c. In vitro Heat-induced albumin denaturation and red blood cell membrane denaturation.	protein denaturation	Naga sravya s (<i>et.al</i> 2011) ¹⁶ .
	d. In vitro Lipo poly saccharide: (LPS) induced nitric oxide	A cell as well as cytotoxic activity	Nguyen Phuong thao (<i>et.al</i> 2019). ¹⁷
	e. In vitro nuclear staining and DNA fragmentation assays.	Induces apoptosis in MEG-01 cells	Satish Kumar Murari (<i>et.al</i> 2016). ¹⁸
2. Antimicrobial	a. In vitro disc diffusion method.	It shows a zone of inhibition on <i>Staphylococcus aureus</i> , <i>X. oryzaepv. Oryzae</i>	Taddei A. (<i>et.al</i> 1999). ¹⁹
	b. In vitro Agar well as disk diffusion method and minimum inhibitory concentration method	A moderate inhibitory activity against all bacterial species with zones of inhibition	Diptanu biswas (<i>et.al</i> 2013). ²⁰
	c. In vitro antimicrobial activity of n-hexane inhibited by bacillus subtilis	Inhibited the growth of bacillus subtilis shows yeast and fungi	Antonietataddei (<i>et.al</i> 2006). ²¹
	d. In vitro agar disc diffusion method	Human pathogenic bacteria was tested	J. Chethan (<i>et.al</i> 2012). ²²
	f. In vivo broth microdilution method in dogs.	Activity against bacteria Escherichia coli in human and dog skin	Ana greiceborbaleite (<i>et.al</i> 2019). ²³
4. Analgesic activity	a. In vivo Kaurenoic acid-induced, PGE2 method in mice.	Blocked the writhing response	Suresh kumar (<i>et.al</i> 2007). ²⁴
	b. In vivo isoprenaline-induced myocardial infarction in rats	Blocking the mitochondrial lipid peroxidation (LPO).	Rangasamy Rajesh (<i>et.al</i> 2007). ²⁵
	c. In vivo acetic acid-induced writhing method and hot plate in mice.	Blocking writhing response	S.Bhama (<i>et.al</i> 2007). ²⁶
5. Hepato protective activity	a. In vivo Croton oil-induced ear dermatitis in rats	Inhibition of writhing response	Meena (<i>et al</i> 2011). ²⁷
	b. In vivo CCl ₄ induced acute hepatotoxicity in rats	Serum aspartate transaminase (AST) and alanine transaminase (ALT)	Karmegam (<i>et al</i> 2008). ²⁸
6. Anti diabetic	a. In vitro α -amylase inhibition method.	IC ₅₀ of α -amylase shows Insulin dysfunction.	Sunita Kanikaram (<i>et al</i> 2018). ²⁹
	b. In vitro Heat-induced albumin denaturation	Protein denaturation.	Pradeep (<i>et.al.</i> 2014). ³⁰

Phytochemical Constituents Reported in *Wedelia trilobata*:

The secondary metabolites from these plants mainly include terpenoids, flavonoids, and polyacetylenes as well as steroids. The leaves and stem contain eudesmanolide lactones, Luteolin, and kaurenoic acid. Different classes of phytoconstituents such as sesquiterpenoids, triterpenoids, and diterpenoids have been reported from the aerial parts,

and the flower of *Trailing daisy trilobata* shown the presence of sterols, flavonoids, Benzene derivatives. Most constituents of oils belong to the large group of terpenes. The essential oil obtained from the leaves of *Wedelia trilobata* was analyzed by GC/MS. Which shows α -pinene (above 30%), α -phellandrene (17.4%) and limonene (16.3%) as major components. Phytochemical constituents are explained in Table.3

Table 3: Phytochemical Constituents reported in *Wedelia trilobata*

Plant part	Phytoconstituents	Reference
Aqueous extract of aerial parts	Terpenoids, flavonoids, polyacetylenes and Steroids.	Sadananda (et.al 2011). ³¹
Aqueous extract of Leaves and stem	Eudesmanolide lactone, luteolin and Arachidonic acid.	Sharanappa (et.al 2011). ³²
Ethanol extract of aerial parts	Sterols, flavonoids, Benzene derivatives.	Guaratini (et.al 2016). ³³
A methanol extract of Leaves	Terpenes	Hoepers (et. al. 2016). ³⁴
Essential oils obtained from leaves analyzed by GC/MS	α -pinene. α -phellandrene. Limonene	Govindappa et.al. (2011). ³⁵

Conclusion:

Presently there is an increase in the significance of herbal medicines with amplified laboratory investigations of biological and pharmacological properties. The plant *Wedelia trilobata* emerged as a good source of medicine with anti-inflammatory anti-microbial, analgesic, anti-oxidant, hepatoprotective and anti-diabetic properties. A large number of phytoconstituents have been isolated and identified from different parts of *Wedelia trilobata* which include flavanoids, triterpenoids luteolin, arachidonic acid, sterols and other constituents of the essential oil reported are α -pinene, α -phellandrene and limonene. The current review on *Wedelia trilobata* can be a good source of information for further research to explore their full therapeutic activity.

Conflicts of Interest: None

References:

1. Oladejo. The characteristics and roles of medicinal plant some important medicinal plants in Nigeria. *Nat Prod Ind J*, 2016; 12(3):102.
2. Adam arsenic, Herbs from distant lands, herbsfromdistantlands.blogspot.com, 2018; 5(58):5-6.
3. P Nagaveni, K Saravana Kumar, Y Ramesh, Pharmacognostic properties and analgesic activity studies of *Abrus precatorius* leaves, *JITPS*, 2012, 3(1): 18-23.
4. Neelam balekar, titpawan nakpheng, and terapal srichana, *Wedelia trilobata* a phytochemical and pharmacological, *Chiang Mai J. Sci*, 2014; 41(3):590-605.
5. Meena AK., mm Rao, RP Meena, p. Panda and renu, Pharmacological and phytochemical evidence for the plants of the *wedelia genus*, *J. Pharm. Reviews*, 2011; 1(1):1-6
6. Maldini M, S. Sosa, P. Montoro, A. giangaspero, M.J. balick, C.pizza, R. Della loggia. Screening of the topical anti-inflammatory activity of the leaves, *sphagneticola trilobata* Hitchcock., *Journal of ethnopharmacology*, 2009; 122:430-433.
7. Taddei A, A.J. Rosas-Romero, Anti-microbial activity of *Wedelia trilobata* crude extracts, *Phytomedicine*, 1999; 6(2):133-4.
8. Mandal p, babu's sp, Mandal NC, Anti-microbial, anti-oxidant and anti-inflammatory activity and phytochemical screening of water extract of *Wedelia trilobata* (L) hitch.journal of medicinal plants, 2011; 5(24): 462-465.
9. Suresh Kumar S, S. bhama, T.S. Kumar, M.J.N Chandrasekar, Analgesic activities of the medicinal plant of *Wedelia trilobata*, w.biflora and eclipta Alba in standard experimental animal models, *Worldwide journal of multidisciplinary research and development*, 2007; 19(6):4674-4678.
10. Teerapol srichana, titpawan nakpheng, Neelam balekar. *Wedelia trilobata* (L), A phytochemical and pharmacological, *Chiang Mai J. Sci*, 2014; 41(3):590-605.
11. Murugaian P, V. Ramamurthy, and N. Karmegam, Hepatoprotective activity of *wedelia calendoulacea* against acute hepatotoxicity in rats, *Research Journal of Agriculture and Biological Sciences*, 2008; 4(6):685-687.
12. Lin sc, Lin cc, gin yh, shyyu SJ, Hepatoprotective effects of Taiwan folk medicine: *wedelia Chinensis* on three hepatotoxins induced hepatotoxicity, *Is J Chin Med*, 1994; 22(2):155-68.
13. Yerikala Ramesh, Koorapati Balasaradhi, Kaki Rohan Abhilash, 2019, Formulation and evaluation of Albendazole Nanoparticle, *Journal of Drug Delivery & Therapeutics*, 9(1):16-22.
14. Govindappa M.Naga, Lan C, *wedelia trilobata*, A phytochemical and pharmacological review, *J. Pharmacog. Phytother*, 2014; 41(3):590-605.
15. Giovana fucina, Lilian wunseh Rocha, gislaine francigli da silva, Topical anti-inflammatory phytomedicine based on *sphagneticola trilobata* dried extract, *pharmaceutical biology*, 2016; 54(11):2465-2474.
16. Nagasravya S, Poojasri M.N, Sadanand T.S, chandrappa C.P, Anti-microbial, anti-oxidant and in-vitro anti-inflammatory activity and phytochemical screening of water extract of *Wedelia trilobata* (L) Hichc, *journal of medicinal plant research*, 2011; 5(24):5718-5729.
17. Nguyen Phuong Thao, Pham Thanh Binh, Nguyen thi luyen, Nguyen Duy Cong, Anti-inflammatory and cytotoxic activities of constituents from *wedelia trilobata* (L). Hitchc, *Vietnam journal of chemistry*, 2019; 57(1):121-127.
18. Satish Kumar Murari, Shankar Jayaram, Chetan javara Shetty, shiva Prasad Koller: Methanolic extract of *Wedelia trilobata* on antiproliferation and apoptotic activity, *Natural products chemistry and research*, 2016; 4(2):2329-6836.
19. Taddei A, A.J. Rosas-Romero: Anti-microbial activity of *Wedelia trilobata* crude extracts, *Phytomedicine*, 1999; 6(2):133-4.
20. Diptanu Biswas, Prakash Yogananda, avnitabha dey, Lokesh deb: Evaluation of anti-microbial and wound healing potential of ethanol extract of *wedelia bilfora* Linn D.C.leaves, 2013; 75(2):156-61.
21. Antonieta taddei, Maria. Jose Rodriguez, Ernesto marquez-vilchez, Cristina castelli, Isolation and identification of streptomycin lep spp: from Venezuelan soils, Morphological and biochemical studies *sphagneticola trilobata*, 2006; 161(3):222-31.
22. Chethan J, K.K. Sampath Kumara, Shailasree Sekhar, HS. Prakash, Anti-oxidant, anti-bacterial and DNA protecting

- activity of selected medicinally important Asteraceae plants, 2012; 4(2):257-261.
23. Ana greice Borba Leite, eulina Tereza nery Farias, ana Paula de Oliveira, and ruan Emmanuel franco de Abreu, Mateus matiuzzi da costa, Phytochemical screening and antimicrobial activity of testing crude hydroalcoholic extract from leaves of sphagneticola trilobata. (Asteraceae), 2019; 49(4):1678-4596.
24. Sureshkumar S, bhama S, Kumar T.S, M.J.N.chandrasekar: Analgesic activities of the medicinal plants of *wedelia trilobata*, *wedelia biflora*, and *eclipta Alba* in standard experimental animal models: Worldwide journal of multidisciplinary research and development, 2007; 19(6):4674-4678.
25. Rangasamy Rajesh, Balarama Ganesan, Rangasamy anandan, Nanjappan dhandapan, Biochemical studies on the protective effect of betaine on mitochondrial function in experimentally induced myocardial infarction in rats (*wedelia trilobata*), 2007; 53(6):671-681
26. Bhama S, M.J.N.chandarasekhar, and Kumar T.S, Suresh Kumar S, Analgesic activities of the medicinal plants of *Wedelia trilobata*, *wedelia biflora*, and *eclipta Alba* in standard experimental animal models, worldwide journal of multidisciplinary research and development, 2007; 19(6):4674-4678.
27. Meena RP, Rao MM, renu, panda P, Pharmacological and phytochemical evidence for the plants of *wedelia* genus, J. Pharm. Reviews, 2011; 1(1):2231-5683.
28. Karmegan N, Ramamurthy V, Murugaian P, Hepatoprotective activity of *wedelia calendulacea* L. Against acute hepatotoxicity in rats, Research Journal of Agriculture and Biological Sciences, 2008; 4(6):685-687.
29. Sunita kanikaram, Hari babu Bollikolla, Swarna Latha kurapati, Ramya Krishna Pallapatti, A Quantitative estimation of phytochemicals, anti-diabetic and antioxidant activities of crude extracts of *sphagneticola trilobata* (L) and *adathoda vasica* Linn, J. Nat. Prod. Resour, 2018; 4(1):155-159.
30. Pradeep Kumar P.M, Prakash H.S, Chetan J, Anti-diabetic and antihypertensive potential of selected Asteraceae plant species, American journal of advanced drug delivery, 2014; 2(3):355-363.
31. Sadananda T.S, chandrappa C.P, Gustavo, Santoyo, Anil Kumar N.V. Anti-inflammatory activity and phytochemical screening of water extract of *Wedelia trilobata* (L) hitch, journal of medicinal plants research, 2011; 5(24):5718-5729.
32. Sharanappa P, Anil Kumar N.V, Naga sravya S, Poojashri M.N. In vitro anti-inflammatory activity and phytochemical screening of water extract of *Wedelia trilobata* (L). Hitch, journal of medicinal plant research, 2011; 5(24):5718-5729.
33. Guaratini Thais, Marcelo P. Barros, Vanes r. Falcao, Karina H.M. Cardozo. Metabolites from algae with the economical impact of *wedelia trilobata*, Comparative biochemistry and physiology, science direct.com, 2007; 60-78:146.
34. Hoeper's S.M, Ferreira FP, Guaratin T, Fucina G, and Rochon LW, Topical anti-inflammatory phytomedicine based on *sphagneticola trilobata* dried extract, pharmaceutical biology, 2016; 54(11):2465-2474.
35. Govindappa M, Naga S.S, Poojasri M.N, Sadanand T.S, and Chandrappa C.P, Anti-microbial, the anti-oxidant and in-vivo anti-inflammatory activity of ethanol extract and active phytochemical screening of *Wedelia trilobata* (L), Journal of medicinal plant research, 2011; 3:43-51.

