

Available online on 30.07.2019 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

© 2011-18, publisher and licensee JDDT, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

QSAR Studies of Flavonoids Derivatives for Antioxidant and Antimicrobial Activity

Sharma Mukesh^{1*}, Ahuja Dharmendra²¹ Research Scholar, Faculty of Pharmaceutical Sciences Jayoti Vidyapeeth Women's University, Jaipur, Rajasthan, India² Professor, Faculty of Pharmaceutical Sciences Jayoti Vidyapeeth Women's University, Jaipur, Rajasthan, India

ABSTRACT

The biological activity and the molecular structures are the two main aspects for the QSAR study of a specific compound, which leads to generate a new chemical moiety. The Quantitative Structure Activity Relationship (QSAR) paradigm is based on the assumption that there is an underlying relationship between the molecular structure and biological activity. Physico-chemical properties were calculated employing the modeling software Win CAChe 6.1. The calculated descriptors were conformational minimum energies (CME), HOMO energy, LUMO energy, heat of formation (HF), ionization potential (IP), molar refractivity (MR), Shape Index (basic kappa, order 1) (SI1), log P, electron affinity (EA), solvent accessible surface area (SAS). In random selection, 18 compounds were in training set and 10 in test set. Subsequently with the stepwise multiple linear regression analysis was carried out to achieve the best models. The equation generated was validated. The selected QSAR model showed correlation coefficient R^2 0.7609, and cross-validated squared correlation coefficient Q^2 of 0.5041 for antioxidant activity ANOVA of predicted value for Variance were found as 0.063638, 201.73 and 0.112396 for group -1.51851, 86.27 and -8.151 respectively. Source of Variation for between and within group i.e. SS value 40522.91 df value 2, MS value 20261.45 F value 301.0527 and p-value 1.01E-17. The HOMO-LUMO range were 73.149 to 84.775, Molar refractivity range was observed -7.409 to -8.84, Predicted value range was -2.52559 to -2.98191 for compound V1 to V5 and R1, R2. The result of study indicated that C5 [1-(2-hydroxyphenyl)-5-phenylpenta-2,4-dien-1-one]; is only inactive against *Streptococcus mutans*. All 3-hydroxyflavone derivatives exhibited their MIC to be in range of 250-125 μ g/ml., 2,3-dihydroflavan-3-ol derivatives exhibited their MIC to be in range of 1000- 500 μ g/ml. The chalcone derivatives exhibited their MIC to be at 250 μ g/ml.

Keywords: QSAR, *Streptococcus mutans*, Win CAChe 6.1 and antioxidant activity

Article Info: Received 25 May 2019; Review Completed 13 July 2019; Accepted 22 July 2019; Available online 30 July 2019



Cite this article as:

Sharma M, Ahuja D, QSAR Studies of Flavonoids Derivatives for Antioxidant and Antimicrobial Activity, Journal of Drug Delivery and Therapeutics. 2019; 9(4):765-773

DOI: <http://dx.doi.org/10.22270/jddt.v9i4.3967>

*Address for Correspondence:

Sharma Mukesh, Research Scholar, Faculty of Pharmaceutical Sciences Jayoti Vidyapeeth Women's University, Jaipur, Rajasthan, India

INTRODUCTION:

Drug discovery is the extremely time consuming and expensive processes. With rapidly increasing costs, diminishing resources, more intense public scrutiny, greater government regulations, and higher expectations, this hit and- miss approach is not effective on economical and efficiency basis. The combined application of molecular graphics, computational pharmaceutical chemistry and biological data studies are comes in the field of computer added drug design (CADD). These advance computer based techniques are helpful for the achieving to long-term goal for a medical analyst: the prediction of pharmacological activity prior to extensive chemical synthesis and biological data testing.

The biological activity and the molecular structures are the two main aspects for the QSAR study of a specific compound, which leads to generate a new chemical moiety. The Quantitative Structure Activity Relationship (QSAR) paradigm is based on the assumption that there is an underlying relationship between the molecular structure and biological activity. On the basis of this assumption QSAR study establish a correlation between various molecular properties of a series of molecular compound with their well known biological activity.

The antioxidant effects of flavonoids lead to many effective and observable pharmacological responses as cardiovascular, antithrombogenic, antitumor, anti-inflammatory, antidiarrheal, antiviral, antiplatelet etc. Flavonoids are the well known biological response. They have the ability of

CODEN (USA): JDDTAO

modification to the metabolic reaction and activity of body to allergic molecules, viruses and carcinogens. Thus, goal of this work is to syntheses flavonoids starting from commonly available raw materials and its evaluation for antioxidant and antibacterial activity.

EXPERIMENTAL WORK:

Selection of training and test set:

QSAR models are used largely to screen chemical databases and/or virtual chemical libraries for potentially biological active molecules. These developments highlight the importance of rigorous model validation to ensure that the models have both the ability to explain the difference in the biological activity (internal validation) and the acceptable anticipate power e.g. external validation. For model validation the dataset is required to be divided into test set for making the QSAR model and test set for screening its predictive power. For any QSAR model, it is of very importance that the training set selected to calibrate the model exhibits a well-balanced distribution and contains representative molecules.

Evaluation of the Model:


The numbers of statistical methods are available for evaluation of the significance of the model, these methods are;

1. **n** number of molecules (> 20 molecules)
2. **k** number of descriptors in a model (statistically $n/5$ descriptors in a model)

3. **df** degree of freedom ($n-k-1$) (higher is better)
4. **r₂** coefficient of determination (> 0.7)
5. **q₂** cross-validated r^2 (>0.5)
6. **pred_r₂** r^2 for external test set (>0.5)
7. **SEE** standard error of estimate (smaller is better)
8. **F-test** F-test for statistical significance of the
9. **F_prob.** Alpha error probability (smaller is better)
10. **Zscore** Z score calculated by the randomization test (higher is better)
11. **best_ran_q₂** highest q^2 value in the randomization test (as low as compared to q^2)
12. **best_ran_r₂** highest r^2 value in the randomization test (as low as compared to r^2) alpha test (<0.01) statistical significance parameter by randomization

Interpretation of Model:


Multiple regressions are largely used method for construction of QSAR model. It is very easy to interpret a regression procedure modal, which contribution of each descriptor could be seen by the magnitude and sign of its regression coefficient. A descriptor coefficient magnitude shows its relative contribution w.r.t other descriptors and sign indicates whether it is directly (positive) or inversely (negative) proportional to the activity.



Procedure:


2.5 mmol of the benzaldehyde derivative and 2.5 mmol of the acetophenone derivative were measured accurately and placed in a mortar. To this added 3 pellets of NaOH (7.5 mmole). Mixture was ground for about 5-10 min. The mixture was turned yellow and pasty after a few times of

grinding. Grinding was continued until the mixture turn into a solid mass and the solid became powdery. 10 mL of water added to the mortar at the last minute of the grinding period. Product was mixed properly with the water, using the spatula solid product was dislodged from the mortar's wall and filtered.



Procedure:

To a mixture of chalcone [0.01mole] in ethyl alcohol [85ml] was added 20 present aqueous sodium hydroxide [10ml] with continues mixing, thereafter by careful addition of hydrogen peroxide [18ml] 20% solution over a period of 0.5hr. The reaction mixture was mixing by continues stir for 3 hrs. at 300c temperature and then put it onto crushed ice containing 5M Hydrochloric acid. The precipitate then filtered. It washed with water and then dried in hot oven. The dried powder was crystallized from chloroform: methyl alcohol [9:1].



Procedure:

A mixture of powdered 2-hydroxychalcone (0.45 mmol) and aq. Sodium hydroxide solution (8 N, 1.0 ml) and a 30% H₂O₂ solution (0.25 g) was mixed with continuous stirring at room temperature for 2 hours. The raw product was filtered off. It was washed with water. The product was dried in a desiccator to give flavanol and recrystallization of the crude product from methyl alcohol.


Characterization of Final Compounds:

The identification & characterization of the prepared compounds were performed on the basis of their physicochemical and spectral data. Such as thin layer chromatography (TLC), melting point (MP), UV-visible analysis(UV), infrared spectroscopy (IR) and Nuclear Magnetic resonance method (¹H NMR).

RESULT AND DISCUSSION:

Physico-chemical properties were calculated employing the modeling software Win CAChe 6.1. The calculated

descriptors were conformational minimum energies(CME) , HOMO energy, LUMO energy, heat of formation (HF), ionization potential (IP), molar refractivity(MR), Shape Index (basic kappa, order 1) (SI1), log P, electron affinity (EA), solvent accessible surface area (SAS). The above properties served as source data for the statistical software and a correlation matrix was prepared to select the parameters with a very low inter-correlation and maximum correlation with the studied compounds anti- oxidant activity. In random selection, 18 compounds were in training set and 10 in test set. Subsequently with the stepwise multiple linear regression analysis was carried out to achieve the best models.


**Table 1: Substitution pattern of the series of flavonoids examined for their antiradical activity**

Compound	RSAexp. /%	R3	R5	R7	R8	R2'	R3'	R4'	R'5	C2=C3
Morin	96.5	OH	OH	OH	H	OH	H	OH	H	+
Taxifolin	94.8	OH	OH	OH	H	H	OH	OH	H	-
Kaempferol	93.5	OH	OH	OH	H	H	H	OH	H	+
Fustin	91.9	OH	H	OH	H	H	OH	OH	H	-
Galangin	91.8	OH	OH	OH	H	H	H	H	H	+
Rutin	90.9	Ogl	OH	OH	H	H	OH	OH	H	+
Quercetin	89.8	OH	OH	OH	H	H	OH	OH	H	+
luteolin 7-gl	87.6	H	OH	Ogl	H	H	OH	OH	H	+
quercetin 3,7,-digl	86.8	Ogl	OH	Ogl	H	H	OH	OH	H	+
laricytrin	84.6	OH	OH	OH	H	H	OH	OH	OMe	+
laricetrin-3'-gl	83.8	OH	OH	OH	H	H	Ogl	OH	OMe	+
robinetin	82.3	OH	H	OH	H	H	OH	OH	OH	+
fisetin	79	OH	H	OH	H	H	OH	OH	H	+
myricetin	72.8	OH	OH	OH	H	H	OH	OH	OH	+
kaempferol	70.6	Ogl	OH	Ogl	H	H	H	OH	H	+
Apigenin-7-gl	34.8	H	OH	Ogl	H	H	H	OH	H	+
hesperetin	30	H	OH	OH	H	H	OH	OMe	H	-
vitexin	21	H	OH	OH	Ogl	H	H	OH	H	+
3,5,7,3',4',5'-hexamethoxyflavone	12.6	OMe	OMe	OMe	H	H	OMe	OMe	OMe	+
naringenin	6.3	H	OH	OH	H	H	H	OH	H	-
Naringin	4.7	H	OH	Ogl	H	H	H	OH	H	-
7-hydroxyflavone	2.8	H	H	OH	H	H	H	H	H	+
Flavanone	2.6	H	H	H	H	H	H	H	H	-
Flavones	1.5	H	H	H	H	H	H	H	H	+
Chrysin	1.1	H	OH	OH	H	H	H	H	H	+
Apigenin	0.7	H	OH	OH	H	H	H	OH	H	+
8-methoxyflavone	0.7	H	H	H	OMe	H	H	H	H	+
5-hydroxyflavone	0.6	H	OH	H	H	H	H	H	H	+

Experimental values as RSA, radical scavenging activity (percents) were taken from ref. 26, gl: glycosyl

Table 2: physic-chemical parameters of flavonoids

Flavonoids	Log (1/Activity)	eA	HF	HOMO	IP	Log P	LUMO	MR	SI1	CME	SAS
3,5,7,3',4',5'-Hexamethoxyflavone	-1.1	0.76	206.16	-8.82	8.82	0.19	-0.76	105.74	23.66	-206.16	398.7
5-hydroxyflavone	0.22	0.75	-33.54	-9.13	9.13	2.03	-0.75	68.67	13.01	-33.54	252.7
7-hydroxyflavone	-0.44	0.87	-37.80	-9.34	9.34	2.03	-0.87	68.67	13.01	-37.80	255.0
8-methoxyflavone	0.15	0.78	-27.74	-8.98	8.98	2.07	-0.78	73.43	13.96	-27.74	272.5
Naringin	-0.67	0.29	-567.09	-9.12	9.12	0.09	-0.29	133.73	32.40	-567.09	520.9
Apigenin Glucoside	-1.54	0.89	-360.20	-9.08	9.08	0.00	-0.89	104.49	24.14	-360.20	406.7
Apigenin	0.15	0.78	-123.03	-9.05	9.05	1.47	-0.78	72.05	14.92	-123.03	273.1
Chrysin	-0.04	0.78	-78.45	-9.23	9.23	1.75	-0.78	70.36	13.96	-78.45	262.7
Fisetin	-1.89	0.98	-154.49	-8.74	8.74	0.56	-0.98	73.73	15.88	-154.49	280.7
Flavanone	-0.41	0.41	-16.59	-9.30	9.30	2.84	-0.41	65.35	12.06	-16.59	243.4
Flavones	-0.17	0.85	6.90	-9.27	9.27	2.32	-0.85	66.97	12.06	6.90	244.7
Fustin	-1.96	0.51	-184.42	-9.09	9.09	1.56	-0.51	71.77	15.88	-184.42	281.8
Galangin	-1.96	0.79	-111.31	-8.85	8.85	0.85	-0.79	72.04	14.92	-111.31	268.7
Hesperitin	-1.47	0.36	-181.60	-8.84	8.84	1.74	-0.36	76.89	16.84	-181.60	298.5
Kaempferol dirhamnose	-1.84	0.82	-532.89	-8.94	8.94	-2.96	-0.82	134.13	32.40	-532.89	517.8
kaempferol	-1.97	0.79	-155.60	-8.71	8.71	0.56	-0.79	73.73	15.88	-155.60	278.8
laricetin 3-O-glucoside	-1.92	1.01	-497.00	-8.75	8.75	-1.72	-1.01	114.68	28.02	-497.00	443.2
Laricytrin	-1.93	0.86	-225.57	-8.69	8.69	0.03	-0.86	81.89	18.78	-225.57	314.6
luteoline glucoside	-1.94	0.87	-395.90	-8.91	8.91	-0.28	-0.87	106.18	25.10	-395.90	413.1
Morin	-1.98	0.68	-195.33	-8.53	8.53	0.28	-0.68	75.43	16.84	-195.33	285.8
Myricetin	-1.86	0.95	-235.10	-8.79	8.79	-0.01	-0.95	77.12	17.81	-235.10	296.7
Naringenin	-0.80	0.35	-147.57	-9.24	9.24	1.99	-0.35	70.43	14.92	-147.57	270.8
Quercetin Diglucoside	-1.94	1.74	-559.06	-7.84	7.84	-4.41	-1.74	130.08	34.34	-559.06	517.5
Quercetin	-1.95	0.88	-195.45	-8.68	8.68	0.28	-0.88	75.43	16.84	-195.45	288.5
Robenitin	-1.92	1.04	-194.10	-8.85	8.85	0.28	-1.04	75.43	16.84	-194.10	289.1
Rutin	-1.96	0.48	-617.39	-9.00	9.00	-4.53	-0.48	137.27	34.34	-617.39	513.6
Taxifolin	-1.98	0.43	-225.85	-9.05	9.05	1.27	-0.43	73.46	16.84	-225.85	290.2
Vitaexine	-1.32	0.92	-399.67	-9.01	9.01	-0.28	-0.92	106.18	25.10	-399.67	398.6


**Figure 1: Experimental vs predicted****Table 3: ANOVA of predicted value**

Groups	Count	Sum	Average	Variance
-1.51851	9	-14.6228	-1.62475	0.063638
86.27	9	694.379	77.15322	201.73
-8.151	9	-72.489	-8.05433	0.112396


Source of Variation	SS	df	MS	F	p-value	F-crit
Between Groups	40522.91	2	20261.45	301.0527	1.01E-17	3.402826
Within Groups	1615.248	24	67.30202			
Total	42138.16	26				

Table 4: predicted activity of synthesized compounds

Comp. code	HOMO-LUMO	Molar refractivity	Prediction
V1	68.652	-7.927	-2.79822
V2	75.115	-7.769	-2.76108
V3	83.08	-7.43	-2.52559
V4	84.775	-7.409	-2.62972
V5	78.97	-7.595	-2.67344
R1	66.685	-8.841	-2.98191
R2	73.149	-8.778	-2.87362

**Figure 2: General structure of 3-hydroxy flavonoid derivatives****Table 5: Different substitution of synthesized compounds**

Compound code	COMPOUND NAME	SUBSTITUENTS				
		X	R 5	R 6	R 7	R 8
V 1	3-hydroxy-2-phenyl-4H-chromen-4-one	2 - Hydroxyphenyl	H	H	H	H
V 2	3-hydroxy-2-(4-methoxyphenyl)- 4H-chromen-4-one	4 - Isopropylphenyl	H	H	H	H
V 3	3-hydroxy-2-p-tolyl-4H-chromen-4-one	4 - Methylphenyl	H	H	H	H
V 4	3,7-dihydroxy-2-(2-hydroxyphenyl)- 4H-chromen-4-one	2 - Hydroxyphenyl	H	H	OH	H
V 5	3,7-dihydroxy-2-(4-isopropylphenyl)- 4H-chromen-4-one	4 - Isopropylphenyl	H	H	OH	H
V 6	3,7-dihydroxy-2-p-tolyl-4H-chromen-4-one	4 - Methylphenyl	H	H	OH	H

**Figure 3: General structure of 2, 3-dihydroxy flavan-3-ol derivatives****Table 5: Different substitution of 2, 3-dihydroxy flavan-3-ol derivatives**

Compound Code	Compound Name	Substituents				
		X	R 5	R 6	R 7	R 8
R1	3-hydroxy-2-phenyl-2,3-dihydro- 4H-chromen-4-one	2 - Hydroxyphenyl	H	H	H	H
R2	2,3 -dihydro-3-hydroxy-2-(4-isopropylphenyl)chromen-4-one	4 - Isopropylphenyl	H	H	H	H
R3	2,3-dihydro-3-hydroxy-2-p- toylchromen-4-one	4 - Methylphenyl	H	H	H	H
R4	2,3-dihydro-3,7-dihydroxy-2-(2-hydroxyphenyl)chromen-4-one	2 - Hydroxyphenyl	H	H	OH	H
R5	2,3-dihydro-3,7-dihydroxy-2-(4-isopropylphenyl)chromen-4-one	4 - Isopropylphenyl	H	H	OH	H
R6	3, 7-dihydroxy-p-tolylchromen-4- one	4 - Methylphenyl	H	H	OH	H

Minimum inhibitory concentration:

On observation of positive results of antimicrobial activity, the MICs of synthesized compounds were determined by tube dilution method. Studies indicated the requirement of inhibitory concentration in order "chalcone derivatives > 2,3-dihydroflavan-3-ol derivatives > 3-hydroxyflavone

derivatives. The result of study indicated that C5 [1-(2-hydroxyphenyl)-5-phenylpenta-2,4-dien-1-one]; is only inactive against *Streptococcus mutans*. All 3-hydroxyflavone derivatives exhibited their MIC to be in range of 250-125 µg/ml., 2,3-dihydroflavan-3-ol derivatives exhibited their MIC to be in range of 1000- 500 µg/ml. The chalcone derivatives exhibited their MIC to be at 250 µg/ml.

Table 6: MIC of flavonoid derivatives with *Streptococcus mutans*

Compound code	% Transmittance			
	1000 µg/ml	500 µg/ml	250 µg/ml	125 µg/ml
J1	37	34	30	17
J2	24	26	29	16
J3	18	27	17	19
J4	33	35	28	19
J5	30	31	29	26
J6	31	39	34	27
V1	30	37	41	25
V2	27	33	30	27
V3	32	34	36	12
V4	33	36	25	15
V5	33	35	28	19
V6	32	33	30	28
R1	30	39	35	27
R2	47	24	21	25
R3	34	15	18	15
R4	58	43	23	27
R5	44	48	32	30
R6	53	16	13	18




Figure 4: MIC of 2-hydroxy Chalcone derivatives




Figure 5: MIC of 3-hydroxy flavone derivatives




Figure 6: MIC of 2, 3-dihydroxy flavan-3-ol derivatives

CONCLUSION:

A QSAR study was performed on a series of flavonoids using WIN CAChE 6.1 and Vlife MDS 3.5 to explore the physico-chemical parameters responsible for their antioxidant activity. The equation generated was validated. The selected QSAR model showed correlation coefficient R^2 0.7609, and cross-validated squared correlation coefficient Q^2 of 0.5041 for antioxidant activity. The developed model indicated that positive dependence of molar refractivity & HOMO-LUMO gap on the antioxidant activity of flavonoids. The newly synthesized flavonoid derivatives were screened for antioxidant, antimicrobial activity. The results of pharmacological screenings are satisfactory. Most of the derivatives have shown comparable antioxidant activity in relation to standard Ascorbic acid. These derivatives were found to have inhibitory effect against *Streptococcus mutans*. The result of antimicrobial study against microorganism *Streptococcus mutans* shows flavonoid derivatives possess antibacterial activity.

REFERENCES:

1. Bilkis jahan Lumbiny1 , Zhang hui, Antiaging, Antioxidant Flavonoids; Synthesis, Antimicrobial Screening as well as 3d qsar COMFA Models for the Prediction of Biological Activity, *j. asiat. soc. bangladesh, sci.* 2013; 39(2):191-199.
2. Mostahar, S., P. Khatun and M. A. Islam. Synthesis of two vanillin ring containing flavones by different methods and studies of their antibacterial and antifungal activities. *Journal of Biological Science.* 2007; 7(3):514-519.
3. Patrick, G. L. 2001. An Introduction to Medicinal Chemistry. *Oxford University Press.* U. K. 450- 465.
4. Putambaker, D. S., R. Giridhar and M. 3D-QSAR CoMFA/CoMSIA studies on 5- aryl-2, 2-dialkyl-4-phenyl-3 (2H)-furanone derivatives as selective COX-2 Inhibitors. *Acta Pharm.* 2006; 56:157-174.
5. Samee, W., J. Ungwitayatorn and J. Pimthon. 3D-QSAR Studies on Phthalimide Derivatives as HIV-1Reverse Transcriptase Inhibitors. *Science Asia.* 2004; 30:81- 88.
6. Shirley, B.W. Flavonoid Biosynthesis. A colorful Model for Genetics, Biochemistry, Cell Biology, Biotechnology. *Plant Physiology.* 2001; 126:485-493.

7. Tapas, A. R., D. M. Sharker and R.B. Kakde. Flavonoids as Neutraceuticals: A Review. *Tropical Journal of Pharmaceutical Research*. 2008; 7(3):1089-1099.
8. Young, D. C. Computational Chemistry; A Practical Guide for Applying Techniques to Real World Problems; John Wiley and Sons. 2001; 243-249.
9. Chandra, P. Yadav, A. Patni, B.; Mishra, A.P.; Nautiyal, A.R. Antimicrobial resistance and the alternative resources with special emphasis on plant-based antimicrobials-A review. *Plants* 2017; 6:16.
10. Gupta, P.D.; Birdi, T.J. Development of botanicals to combat antibiotic resistance. *J. Ayur. Integr. Med.* 2017; 8:266-275.
11. Zheng, J.; Huang, C.; Yang, B.; Kallio, H. Regulation of phytochemicals in fruits and berries by environmental variation—Sugars and organic acids. *J. Food Biochem.* 2019, 43, e12642. *J. Clin. Med.* 2020, 9, 109:13 of 17
12. Özcelik, B. Kartal, M. Orhan, I. Cytotoxicity, antiviral and antimicrobial activities of alkaloids, flavonoids, and phenolic acids. *Pharm. Biol.* 2011; 49:396-402.
13. Barbieri, R. Coppo, E.; Marchese, A. Daglia, M. Sobarzo-Sánchez, Nabavi, S.M. Phytochemicals for human disease: An update on plant-derived compounds antibacterial activity. *Microbiol. Res.* 2017; 196:44-68.
14. Fialova, S. Rendekova, K. Mucaji, P. Slobodnikova, L. Plant natural agents: Polyphenols, alkaloids and essential oils as perspective solution of microbial resistance. *Curr. Org. Chem.* 2017; 21:1875-1884.
15. Khameneh, B. Iranshahy, M.; Soheili, B.S.F. Review on plant antimicrobials: A mechanistic viewpoint. *Antimicrob. Resist. Infect. Control* 2019; 8:118.
16. Gutiérrez-Grijalva, E.P. Picos-Salas, M.A. Leyva-López, N. Criollo-Mendoza. Flavonoids and phenolic acids from oregano: Occurrence, biological activity and health benefits. *Plants* 2018, 7, 2.
17. Jungbauer, A. Medjakovic, S. Anti-inflammatory properties of culinary herbs and spices that ameliorate the effects of metabolic syndrome. *Maturitas* 2012, 71, 227-239.
18. Goncalves, S. Moreira, E. Grosso, C. Andrade, Phenolic profile, antioxidant activity and enzyme inhibitory activities of extracts from aromatic plants used in mediterranean diet. *J. Food Sci. Technol.* 2017, 54, 219-227.
19. Mileo, A.M. Nisticò, P. Miccadei, S. Polyphenols: Immunomodulatory and therapeutic implication in colorectal cancer. *Front. Immunol.* 2019, 10, 729. [CrossRef] [PubMed]
20. Singh, A.K. Cabral, C. Kumar, R. Ganguly, R. Rana, H.K. Gupta, A. Lauro, M.R. Carbone, C. Reis, F. Pandey, A.K. Beneficial effects of dietary polyphenols on gut microbiota and strategies to improve delivery efficiency. *Nutrients* 2019, 11, 2216.
21. Firrman, J. Liu, L.S. Yam, K. A review on flavonoid apigenin: Dietary intake, ADME, antimicrobial effects, and interactions with human gut microbiota. *BioMed Res. Int.* 2019, 2019, 7010467.
22. Chua, L.S. A review on plant-based rutin extraction methods and its pharmacological activities. *J. Ethnopharmacol.* 2013, 150, 805-817.
23. Enogieru, A.B. Haylett, W. Hiss, D.C. Bardien, S. Ekpo, O.E. Rutin as a potent antioxidant: Implications for neurodegenerative disorders. *Oxid. Med. Cell Longev.* 2018, 2018, 6241017.
24. Edwards, P.N. Talent, N. Dickinson, T.A. Shipley, P.R. A review of the chemistry of the genus Crataegus. *Phytochemistry* 2012, 79, 5-26.
25. Yamaguchi, K.K.L. Pereira, L.F.R. Lamarão, C.V. Lima, E.S. da Veiga-Junior, V.F. Amazon acai: Chemistry and biological activities: A review. *Food Chem.* 2015, 179, 137-151.
26. Yuan, L. Wang, J. Wu, W. Liu, Q. Liu, X. Effect of isoorientin on intracellular antioxidant defence mechanisms in hepatoma and liver cell lines. *Biomed. Pharmacother.* 2016, 81, 356-362.
27. Mani, R. Natesan, V. Chrysin: Sources, beneficial pharmacological activities, and molecular mechanism of action. *Phytochemistry* 2018, 145, 187-196.
28. Knekt P, Jarvinen R, Seppanen R, Dietary flavonoids and the risk of lung cancer and other malignant neoplasms. *Am J Epidemiol.* (1997), 146:223-30
29. Robak J, Gryglewski RJ, *Pol J Pharmacol.* (1996), 48:555-64.
30. Gurdeep Chatwal, Organic Chemistry Of Natural Products, Vol.II, (2002), Himalaya Publishing House, 225-245
31. O.P.Agrawal, Chemistry of Organic Natural Products, Vol.II, (2001), Goel Publishing House, Meerut, 234-257.
32. O'zarowski, M. Piasecka, A. Paszel-Jaworska, A. Chaves, D.S. Romanik, A. Rybczyńska, M. Gryszczynska, A. Sawikowska, A. Kachlicki, P. Mikolajczak, P.L. et al. Comparison of bioactive compounds content in leaf extracts of Passiflora incarnata, P. caerulea and P. alata and in vitro cytotoxic potential on leukemia cell lines. *Rev. Bras. Farmacogn.* 2018, 28, 79-191.
33. Naz, S. Imran, M. Rauf, A. Orhan, I.E. Shariati, M.A. Haq, I.U. Yasmin, I. Shahbaz, M. Qaisrani, T.B. Shah, Z.A. et al. Chrysin: Pharmacological and therapeutic properties. *Life Sci.* 2019, 235, 116797.
34. Przybyłek, I. Karpiński, T.M. Antibacterial properties of propolis. *Molecules* 2019, 24, 2047.
35. Alam, M.A. Subhan, N. Rahman, M.M. Uddin, S.J. Reza, H.M. Sarker, S.D. Effect of Citrus flavonoids, naringin and naringenin, on metabolic syndrome and their mechanisms of action. *Adv. Nutr.* 2014, 5, 404-417.
36. Viljakainen, S. Visti, A. Laakso, S. Concentrations of organic acids and soluble sugars in juices from Nordic berries. *Acta Agric. Scand. Sect. B Soil Plant Sci.* 2002, 52, 101-109.
37. Pande, G. Akoh, C.C. Organic acids, antioxidant capacity, phenolic content and lipid characterisation of Georgia-grown underutilized fruit crops. *Food Chem.* 2010, 120, 1067-1075.
38. Nour, V. Trandafir, I. Ionica, M.E. Ascorbic acid, anthocyanins, organic acids and mineral content of some black and red currant cultivars. *Fruits* 2011, 66, 353-362. *J. Clin. Med.* 2020, 9, 109 14 of 17
39. Kaume, L. Howard, L.R. Devareddy, L. The blackberry fruit: A review on its composition and chemistry, metabolism and bioavailability, and health benefits. *J. Agric. Food Chem.* 2012, 60, 5716-5727.
40. Wang, Y. Johnson-Cicalese, J. Singh, A.P. Vorsa, N. Characterization and quantification of flavonoids and organic acids over fruit development in American cranberry (*Vaccinium macrocarpon*) cultivars using HPLC and APCI-MS/MS. *Plant Sci.* 2017, 262, 91-102.
41. Denev, P. Kratchanova, M. Petrova, I. Klisurova, D. Georgiev, Y. Ognyanov, M. Yanakieva, I. Black chokeberry (*Aronia melanocarpa* (Michx.) Elliot) fruits and functional drinks differ significantly in their chemical composition and antioxidant activity. *J. Chem.* 2018, 2018, 9574587.
42. Adamczak, A. Buchwald, W. Kozłowski, J. Variation in the content of flavonols and main organic acids in the fruit of European cranberry (*Oxycoccus palustris* Pers.) growing in peatlands of North-Western Poland. *Herba Pol.* 2011, 57, 5-15.
43. Jurikova, T. Mlcek, J. Skrovánková, S. Sumczynski, D. Sochor, J. Hlaváčová, I. Snopk, L. Oršávová, J. Fruits of black chokeberry *Aronia melanocarpa* in the prevention of chronic diseases. *Molecules* 2017, 22, 944.
44. Santana-Gálvez, J. Cisneros-Zevallos, L. Jacobo-Velázquez, D.A. Chlorogenic Acid: Recent advances on its dual role as a food additive and a nutraceutical against metabolic syndrome. *Molecules* 2017, 22, 358.
45. Meinhart, A.D. Damin, F.M. Caldeirao, L. Silveira, T.F.F. Filho, J.T. Godoy, H.T. Chlorogenic acid isomer contents in 100 plants commercialized in Brazil. *Food Res. Int.* 2017, 99, 522-530.
46. Shekarchi, M. Hajimehdipoor, H. Saeidnia, S. Gohari, A.R. Hamedani, M.P. Comparative study of rosmarinic acid content in some plants of Labiate family. *Pharmacogn. Mag.* 2012, 8, 37-41.
47. O'zarowski, M. Mikołajczak, P. Bogacz, A. Gryszczynska, A. Kujawska, M. Jodynis-Libert, J. Piasecka, A. Napieczynska, H. Szulc, M. Kujawski, R. et al. Rosmarinus officinalis L. leaf extract improves memory impairment and affects acetylcholinesterase and butyrylcholinesterase activities in rat brain. *Fitoterapia* 2013, 91, 261-271.
48. Ozarowski, M. Mikolajczak, P.L. Piasecka, A. Kachlicki, P. Kujawski, R. Bogacz, A. BartkowiakWieczorek, J. Szulc, M. Kaminska, E. Kujawska, M. et al. Influence of the *Melissa officinalis* leaf extract on long-term memory in scopolamine animal model with assessment of mechanism of action. *Evid. Based Complement. Alternat. Med.* 2016, 2016.
49. Balcke, G.U. Handrick, V. Bergau, N. Fichtner, M. Henning, A. Stellmach, H. Tissier, A. Hause, B. Frolov, A. An UPLC-MS/MS method for highly sensitive high-throughput analysis of phytohormones in plant tissues. *Plant Methods* 2012, 8,

50. Toiu, A. Vlase, L. Oniga, I. Benedec, D. Tămaș, M. HPLC analysis of salicylic derivatives from natural products. *Farmacia* 2011, 59, 106–112.

51. Bijnbeker, S. van der Auwera, A. Voorspoels, S. Noten, B. Hermans, N. Pieters, L. Apers, S. A first step in the quest for the active constituents in *Filipendula ulmaria* (meadowsweet): Comprehensive phytochemical identification by liquid chromatography coupled to quadrupole-orbitrap mass spectrometry. *Planta Med.* 2016, 82, 559–572.

52. Nabavi, S.F. Habtemariam, S. Ahmed, T. Sureda, A. Daglia, M. Sobarzo-Sánchez, E. Nabavi, S.M. Polyphenolic composition of *Crataegus monogyna* Jacq.: From chemistry to medical applications. *Nutrients* 2015, 7, 7708–7728.

53. Kerasioti, E. Apostolou, A. Kafantaris, I. Chronis, K. Kokka, E. Dimitriadou, C. Tzanetou, E.N. Priftis, A. Koulocheri, S.D. Haroutounian, S. et al. Polyphenolic composition of *Rosa canina*, *Rosa semperflorens* and *Pyrocantha coccinea* extracts and assessment of their antioxidant activity in human endothelial cells. *Antioxidants* 2019, 8, 92.

54. Siriwong, S. Thumanu, K. Hengpratoom, T. Eumkeb, G. Synergy and mode of action of ceftazidime plus quercetin or luteolin on *Streptococcus pyogenes*. *Evid. Based Complem. Altern. Med.* 2015.

55. Betts, J.W. Sharili, A.S. Phee, L.M. Wareham, D.W. In vitro activity of epigallocatechin gallate and quercetin alone and in combination versus clinical isolates of methicillin-resistant *Staphylococcus aureus*. *J. Nat. Prod.* 2015, 78, 2145–2148.

56. Hossion, A.M. Zamami, Y. Kandahary, R.K. Tsuchiya, T. Ogawa, W. Iwado, A. Quercetin diacylglycoside analogues showing dual inhibition of DNA gyrase and topoisomerase IV as novel antibacterial agents. *J. Med. Chem.* 2011, 54, 3686–3703.

57. Chen, C.C. Huang, C.Y. Inhibition of *Klebsiella pneumoniae* DnaB helicase by the flavonol galangin. *Protein J.* 2011, 30, 59–65.

58. Lin, H.H. Huang, C.Y. Characterization of flavonol inhibition of DnaB helicase: Real-time monitoring, structural modeling, and proposed mechanism. *J. Biomed. Biotechnol.* 2012, 2012, 735368.

59. Huang, Y.H. Huang, C.C. Chen, C.C. Yang, K.; Huang, C.Y. Inhibition of *Staphylococcus aureus* PriA helicase by flavonol kaempferol. *Protein J.* 2015, 34, 169–172.

60. Farhadi, F. Khameneh, B. Iranshahi, M. Iranshahy, M. Antibacterial activity of flavonoids and their structure-activity relationship: An update review. *Phytother. Res.* 2019, 33, 13–40.

61. Echeverría, J. Opazo, J. Mendoza, L. Urzúa, A. Wilkens, M. Structure-activity and lipophilicity relationships of selected antibacterial natural flavones and flavanones of chilean flora. *Molecules* 2017, 22, 608.

62. Xie, Y. Chen, J. Xiao, A. Liu, L. Antibacterial activity of polyphenols: Structure-activity relationship and influence of hyperglycemic condition. *Molecules* 2017, 22, 1913.

63. Bouarab-Chibane, L. Forquet, V. Lantéri, P. Clément. Antibacterial properties of polyphenols: Characterization and QSAR (Quantitative Structure-Activity Relationship) models. *Front. Microbiol.* 2019, 10, 829.

64. Morimoto, Y. Baba, T. Sasaki, T. Hiramatsu, K. Apigenin as an anti-quinolone-resistance antibiotic. *Int. J. Antimicrob. Agents.* 2015, 46, 666–673.

65. Amin, A. Vincent, R. Séverine, M. Rosmarinic acid and its methyl ester as antimicrobial components of the hydromethanolic extract of *Hyptis atrorubens* Poit. (Lamiaceae). *Evid. Based Complem. Alternat. Med.* 2013, 2013, 604536.

66. Ronald L Prior, Fruits and vegetables in the prevention of cellular oxidative damage, *American Journal of Clinical Nutrition*, (2003), 78(3): 570S-578S.

67. Finar, *Stereochemistry of Natural products*, 5th edition, Vol II, ELBS longman publication, 785.

68. Gary R. Beecher, Overview of Dietary Flavonoids: Nomenclature, Occurrence and Intake, *The American Society for Nutritional Sciences J. Nutr.*, (2003), 133:3248S-3254S.

69. Oyvind M. Andersen, Kenneth R. Markham, *Flavonoids: Chemistry, Biochemistry, and Applications*, (2006), CRC Press.

70. Lichard A. Larson, *Naturally Occurring Antioxidants*

71. De Groot H. Reactive oxygen species in tissue injury., *Hepatogastroenterology* (1994), 41:328–32.

72. Robak J, Gryglewski RJ., *Pol J Pharmacol*, (1996), 48:555–64.

73. Hollman PC, Katan MB. Dietary flavonoids: intake, health effects and bioavailability, *Food Chem Toxicol*, (1999), 37:937–42

74. Okushio K, Matsumoto N, Kohri T, Suzuki M, Nanjo F, Hara Y., Absorption of tea catechins into rat portal vein., *Biol Pharm Bull*, (1996), 19:326–9

75. Piskula MK, Terao J., Accumulation of (-)-epicatechin metabolites in rat plasma after oral administration and distribution of conjugation enzymes in rat tissues. *J Nutr*, (1998), 128:1172–8.