

RESEARCH ARTICLE

PHYTOCHEMICAL SCREENING AND *IN VITRO* ANTIOXIDANT POTENTIAL OF *MEMECYLON UMBELLATUM* BURM LEAF EXTRACTS

*Killedar Suresh¹, More Harinath¹, Mali Sachin², Nadaf Sameer², Salunkhe Sachin², Karade Raviraj².

¹Department of Pharmacognosy, Bharati Vidyapeeth College of Pharmacy, Kolhapur-416013, M.H., India

²Department of Pharmaceutical Technology, Bharati Vidyapeeth College of Pharmacy, Kolhapur-416013, M.H., India

ABSTRACT

Objective: Different dry extracts of *Memecylon umbellatum* Burm leaf obtained by various solvents such as petroleum ether, chloroform, ethyl acetate, acetone, methanol and chloroform water (IP) was screened to reap the benefits of its antioxidant and free radical scavenging properties using ascorbic acid as standard antioxidants. **Methods:** The *in vitro* free radical scavenging activity was evaluated using diphenyl picryl hydrazyl (DPPH) radical method using various concentrations of dry extract in distilled water (1, 2, 4, 8, 16, 20 µg/ml) against blank with ascorbic acid as a standard in same concentrations. **Results:** Among the all extracts, Methanol leaf extract has showed higher Antioxidant activity ($84.65 \pm 0.064\%$) having IC_{50} Value $11.81 \pm 0.033\text{ }\mu\text{g/ml}$ at $20\text{ }\mu\text{g/ml}$. While, IC_{50} value for ascorbic acid was found to be $8.91 \pm 0.084\text{ }\mu\text{g/ml}$. **Conclusion:** The results clearly indicate that Methanol leaf extract of *Memecylon umbellatum* is effective in free radical scavenging. So in future, this may emerge as promising natural herbal source of powerful antioxidant.

Keywords: *Memecylon umbellatum*, DPPH reagent, Antioxidant activity, Ascorbic acid, IC_{50} .

1 INTRODUCTION

The mammalian body has its own multifarious defense mechanism involving natural enzymatic (superoxide dismutase, catalase and glutathione peroxidase) and non-enzymatic (thioredoxin, thiols, and disulfide-bonding) antioxidants which counteract the harmful effects of free radicals and other oxidants¹. In normal metabolism, the levels of oxidants (i.e. free radicals) and antioxidants in humans are maintained in balance, which is necessary for sustaining optimal physiological conditions². Free radicals are generated as a result of impaired balance between reactive oxygen species (ROS) production and antioxidant enzymes³. These are chemically unstable atoms or molecules that cause extensive damage to cells, causes damage to DNA molecule, lipids and proteins⁴. If free radicals overcome the body's ability to regulate them, a condition known as oxidative stress ensues⁵. This could lead to number of life threatening diseases like cardiovascular disease⁶, Parkinson's disease⁷, cancer⁸, mild cognitive impairment⁹, neural disorders¹⁰, Alzheimer's disease¹¹, ulcerative colitis¹², aging¹³, diabetes mellitus¹⁴, anaemia¹⁵, atherosclerosis¹⁶, asthma¹⁷. Protection against this type of disease can be enhanced by ample intake of various dietary food supplements (containing α -tocopherol, b-carotene, and ascorbic acid etc) and synthetic antioxidants, but these synthetic antioxidant capsules and dietary supplements are found to be less effective in various cases. This has attracted a great deal of research interest in natural antioxidants. Several herbs and spices including *Ocimum sanctum*, *Cichorium intybus*, *Piper cubeba*, *Punica granatum*, *Allium sativum*, *Delonix regia*, *Terminalia chebula*, *Zingiber officinale* etc have been reported to exhibit antioxidant activity¹⁸. The majority of the antioxidant activity is due to flavonoids, isoflavones, flavones, anthocyanins, coumarins, lignans, catechins, and isocatechins^{19,20}.

Memecylon umbellatum Burm. (Family: Melastomataceae) is a small evergreen shrub or tree grows up to 8-14 m tall having young tree branches and bears numerous umbellate cymes. The plant is known as "Anjani" in Sanskrit and "Ironwood tree" in English. Plants are distributed mostly in coastal regions of the Deccan peninsula, the eastern and southern part of India all along the Western Ghats and in the Andaman islands^{21, 22}. The leaves have been reported to possess astringent properties and are administered to treat leucorrhoea and gonorrhoea²³. Different extracts of *Memecylon umbellatum* Burm Inflorescences²⁴ and bark²⁵ have been evaluated for its antimicrobial potential. Estimation of total content of tannin²⁶ and Seasonal Variation of Tannin Content in Different Parts has also been carried out²⁷. Estimation of Sugars and minerals in healthy and infected parts has also been carried out²⁸. Different root extract have been proved to possess antioxidant activity²⁹. The decoction of the root is used in the treatment of excessive menstrual discharge³⁰. Leaves are also reported to possess antiviral activity³¹. The literature survey reveals that the leaves and roots of *Memecylon umbellatum* have been investigated for its hypoglycemic activity using alloxan induced hyperglycemia Wistar albino rats^{32, 33}. Wound healing activity of ethanolic extract of the leaves has also been reported³⁴. Plant contains a wide variety of phytoconstituents such as umbellactone, β -amyrine, Oleanolic acid, ursolic acid, sitosterol and organic acids^{35, 36}.

* Corresponding Author

Dr. Suresh Killedar

Professor and Head, Department of Pharmacognosy,
Bharati Vidyapeeth College of Pharmacy,
Kolhapur-416013, M.H., India

Email: sureshgk_64@rediffmail.com,

Phone: 91-231-2637286, Fax: 91-231-2638833

2 MATERIAL AND METHODS

2.1 Plant material

The leaves of *Memecylon umbellatum* were collected in the month of March-April from Gaganbawda hills region, Maharashtra, India. The plant material was taxonomically identified by Dr. S. R. Yadav, Department of Botany, Shivaji University, Kolhapur, India (M.S.). The voucher herbarium specimen is deposited in the Department of Pharmacognosy, Bharati Vidyapeeth College of Pharmacy, Kolhapur.

2.2 Chemical

DPPH Reagent, Ascorbic acid (ACME Chemicals, Mumbai), All other chemicals are of analytical grade and procured from Loba Chem.

2.3 Methods

A standard curve was obtained using Ascorbic acid with the help of double beam UV/Visible spectrophotometer (Jasco-V-630).

2.4 Preparation of extract

Leaves were sorted for foreign matter and dried under shade by spreading in thin layers using aluminum trays for 10 days. Electric grinder (Bajaj-make) was used for powdering soft tissues of leaves. Coarse powder of leaves (#40) was used for extraction.

2.5 Soxhlet extraction process

Extraction was carried out by standard procedure³⁷⁻³⁹. One kg powder of roots of *Memecylon umbellatum* was used for extraction. Sample powder was packed

gently in previously washed and dried cloth bag and solvent was placed from the top with the help of funnel to moisten the drug sample. 3.5 liter of solvent (ethyl acetate, methanol, chloroform water, chloroform, and petroleum ether) was placed in distillation flask and assembly was made air tight with sealing wax. Solvents were selected on the basis of extractive values and with their increasing order of polarity. Extraction was carried out at or slightly above the boiling point of each solvent. Extraction was carried out for 18 hours or on the basis of clarity of dropping solvent (saturation). The solvent was collected every time after completion of the process and powder was dried in hot air oven for 24h at 450C. The process was repeated for all the next solvents and finally the dried powder was macerated with 3.5 liter of chloroform water IP (0.25% v/v) at room temperature with frequent shaking. All the liquid extracts were subjected for physical analysis and are concentrated in a rotary film vacuum evaporator (Dolphin, Mumbai) and finally dried under reduced pressure. The residue was weighed, % yield was calculated. All the extracts were further dried over anhydrous calcium chloride and preserved in vacuum desiccators for further studies. Different extracts were abbreviated according to solvent and part of the plant and used throughout the work.

2.6 Physical evaluation of different liquid extracts of leaf

All the extracts were studied for physical evaluation with respect to color, pH, fluorescence, density, specific gravity, viscosity along with nature of solid residue obtained after concentration of the extracts with % yield has shown in Table 1.

Table 1: Physical analysis of different liquid leaf extracts of *Memecylon umbellatum*

Name of extract	Extract color	pH	Fluorescence			Specific gravity	Density	Viscosity	yield of solids (g)	Nature of solid extract
			D	S	L					
PEEL	DG	6.8	R	O	DB	0.6908	0.8231	0.6124	8.810	Waxy
ChEL	G	6.2	G	R	YG	1.2990	1.5480	1.0290	2.062	Lumpy
SEEL	YG	6.5	Y	P	G	0.6582	0.7850	0.3474	0.375	Powder
EAEL	RB	5.4	-	R	BR	0.6437	0.7265	1.8547	3.065	Powder
ButEL	WR	5.8	-	MG	-	0.6760	0.8058	2.5793	4.604	Powder
AEL	BR	7.2	Y	MW	-	0.6994	0.7801	0.4835	2.243	Powder
EthEL	RB	7.1	-	G	-	0.7902	0.8185	0.9625	8.544	Powder
MEL	RB	6.0	BR	DG	YW	0.8240	0.8848	0.8556	28.36	Waxy
AqEL	RB	5.2	YG	B	G	1.0080	1.0124	1.0210	15.72	Powder

PEEL-Petroleum Ether extract leaf, ChEL-Chloroform Extract Leaf, SEEL- Solvent Ether Extract Leaf, EAEL Ethyl Acetate Extract Leaf, ButEL - n-Butanol Extract Leaf, AEL- Acetone Extract Leaf, EthEL-Ethanol Extract Leaf, MEL-Methanol Extract Leaf, AqEL-Aqueous Extract Leaf, DG -Dark Green, G-Green, YG – Yellowish Green, RB-Reddish Brown, WR-Wine Red, BR-Brownish Red, RB-Reddish Brown, R-Red, G-Green, MG-Milky Green, MW-Milky White, B-Blue, DB-Dark Blue, YW- Yellowish White.

2.7 Phytochemical screening

About 500 mg of each dried extract was dissolved in 100 ml of respective solvent and solution obtained was subjected for Phytochemical screening using different specific and general reagents. Samples were prepared as

per the requirement of procedure and tests were repeated for final confirmation of phytoconstituents. The positive phytoconstituents present in different parts with various solvents have shown in Table 2.

Table 2: Phytochemical screening of leaf extracts of *Memecylon umbellatum*

Extract	Sugars		Alk.	Tannins		Glycosides				steroids		proteins	Org. acids		
	R	NR		HT	CT	a	c	s	f	co	ST		C	O	T
AqEL	+	+	-	+	-	-	+	+	+	+	-	-	+	+	+
MEL	+	+	-	+	+	-	+	+	+	+	-	-	+	+	+
EthEL	+	+	-	+	+	+	-	+	+	-	-	-	+	+	+
ButEL	+	+	-	+	+	-	+	-	+	-	+	-	+	+	+
AEL	+	+	-	+	+	+	+	+	+	-	+	-	+	+	+
EAEL	+	+	-	+	+	+	-	-	+	-	-	+	+	+	-
ChEL	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-
SEEL	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-
PEEL	-	-	-	-	-	-	-	-	-	+	+	-	-	-	-

AqEL-Aqueous Extract Leaf, MEL-Methanol Extract Leaf, EthEL-Ethanol Extract Leaf, ButEL-n Butanol Extract Leaf, AEL-Acetone Extract Leaf, EAEL-Ethyl Acetate Extract Leaf, ChEL-Chloroform Extract Leaf, SEEL-Solvent Ether Extract Leaf, PEEL-Petroleum Ether Extract Leaf, Alk-Alkaloids, Gly-Glycosides, Org.acids-Organic acids, R-Reducing sugars, NR-Non Reducing sugars, HT-Hydrolysable Tannins, CT-Condensed Tannins, a-anthracene glycosides, c-Cardiac glycosides, s-Saponin glycosides, f-Flavanoidal glycosides, co-Coumarin glycosides, ST-Sesquiterpene, TT-Triterpene, +Positive, -Negative.

2.8 Screening of extracts for in-vitro antioxidant activity using DPPH Assay⁴⁰⁻⁴⁶

Extracts showing presence of triterpenes and polyphenolic compounds were screened for antioxidant activity using DPPH reagent. DPPH assay is most widely used method for determination antioxidant potential. Its use has been previously reported for species *Acacia caesia*^[47], *Aerva Lanata*^[48] etc.

2.8.1 Reagents for antioxidant activity


1. DPPH Reagent: Methanolic solution of DPPH (0.1 mM): 39.4 mg of DPPH was dissolved in one liter of analytical grade methanol.

2. Standard solution: Ascorbic acid was used as standard in following concentrations 1, 2, 4, 8, 16, 20 µg/ml in methanol.

3. Sample preparation: Test samples of each dry extract were prepared by dissolving in distilled water in the various concentrations as 1, 2, 4, 8, 16, 20 µg/ml.

2.8.2 Principle:

The scavenging reaction between (DPPH) and an antioxidant (H-A) can be written as:



DPPH (1, 1-diphenyl-2-picrylhydrazyl) is a stable free radical, characterised by the delocalisation of the spare electron over the molecule as a whole. So this does not dimerize unlike the other free radicals. The delocalization of electron also gives rise to the deep violet color. When antioxidants react with DPPH, which is a stable free radical is reduced to the DPPH radical i.e. 1-1 diphenyl - 2 - picryl hydrazine and as consequence there is loss of this violet color. The degree of discoloration indicates the scavenging potential of the antioxidant compounds or extracts in terms of hydrogen donating ability.

2.8.3 Procedure

The DPPH scavenging activity was performed using a solution of 0.1 mM DPPH in methanol solution and 1.0 ml solution was added in 3.0 ml of test samples of each dry extract having concentrations as 1, 2, 4, 8, 16 and 20 µg/ml in methanol and kept in darkness. Thirty

minutes later, the absorbance was measured at 517 nm. A blank was prepared without adding the extract. Ascorbic acid at concentration 1, 2, 4, 8, 16, 20 µg/ml was used as standard. Lower the absorbance of the reaction mixture indicates higher free radical scavenging activity. The capability to scavenge the DPPH radical was calculated using the following equation.

$$\text{DPPH Scavenged (\%)} = \frac{A_{\text{Control}} - A_{\text{Test}}}{A_{\text{Control}}} \times 100$$

Where 'A control' is the absorbance of the control reaction and 'A test' is the absorbance in the presence of the sample of the extracts. The antioxidant activity of the different extract was expressed in % DPPH radical scavenged and the results are given in table 3.

Table 3: Antioxidant activity and IC₅₀ value of different extracts of *Memecylon umbellatum**

Sr. No.	Solvent	% Antioxidant activity of different extracts at various concentrations (µg/ml)						IC ₅₀ (µg/ml)
		1	2	4	8	16	20	
1	Standard	12.36 ± 0.035	29.27 ± 0.055	37.42 ± 0.314	44.89 ± 0.121	74.92 ± 0.099	91.90 ± 0.054	8.91 ± 0.054
2	Petroleum ether	0.36 ± 0.248	0.87 ± 0.159	1.48 ± 0.097	2.39 ± 0.342	2.88 ± 0.463	3.17 ± 0.084	315.45 ± 0.059
3	Chloroform	0.29 ± 0.111	1.08 ± 0.058	1.97 ± 0.096	2.98 ± 0.278	3.86 ± 0.396	4.76 ± 0.036	210.08 ± 0.063
4	Ethyl acetate	8.32 ± 0.296	22.96 ± 0.398	41.36 ± 0.179	52.69 ± 0.224	63.89 ± 0.419	77.24 ± 0.059	12.94 ± 0.016
5	Acetone	11.23 ± 0.385	31.69 ± 0.054	47.65 ± 0.158	57.38 ± 0.269	68.93 ± 0.114	78.83 ± 0.167	12.68 ± 0.064
6	n-Butanol	14.39 ± 0.159	38.96 ± 0.178	48.36 ± 0.342	61.32 ± 0.329	71.88 ± 0.152	82.01 ± 0.018	12.19 ± 0.089
7	Ethanol	5.07 ± 0.118	18.65 ± 0.159	26.65 ± 0.277	33.33 ± 0.096	39.37 ± 0.518	49.20 ± 0.068	20.33 ± 0.018
8	Methanol**	9.56 ± 0.278	19.63 ± 0.152	45.69 ± 0.329	59.36 ± 0.114	79.69 ± 0.278	84.65 ± 0.064	11.81 ± 0.033
9	Aqueous	2.12 ± 0.114	4.48 ± 0.096	6.65 ± 0.342	7.08 ± 0.196	9.96 ± 0.059	11.11 ± 0.128	90.01 ± 0.055

* Indicates ± SD (n=5) ** indicates more potent extract & significance (p<0.05)

IC₅₀ value was determined to express antioxidant activity. It is the concentration of fractions that inhibits the formation of DPPH radicals by 50%. The lower IC₅₀ value represents the higher antioxidant activity of the tested sample.

3 RESULTS AND DISCUSSION

3.1 Evaluation of different liquid extracts of leaf

Most of the extracts have shown different color in different solvents. Some of the extracts have showed typical fluorescence either in day or short (254nm) and long (366nm) wavelengths. Leaf extract showed maximum pH 7.2 for acetone and minimum pH 5.2 for aqueous extract. Specific gravity was found highest (1.2990) for chloroform extract and lowest (0.6437) for ethyl acetate. Also maximum viscosity (2.5793cp) for n-butanol and minimum (0.3474cp) for solvent ether was observed. The maximum % yield 28.36 for leaf was found for methanol and minimum 0.375 % for solvent ether.

3.2 Phytochemical screening of different extracts

Polar solvents used in the process of extraction have shown the presence of polar constituents such as mono and disaccharides, proteins amino acids different glycosides like anthracene, cardiac, flavanoidal, saponin and coumarin type glycosides. Polyphenolics like tannins, organic acids, minerals and triterpenes were also found in most of the polar extracts while non polar solvents showed positive tests for sterols, aglycones of different glycosides, fatty acids and polysaccharides. Phytochemical screening of different liquid extracts showed the presence of reducing, nonreducing sugars, tannins and proteins (both hydrolysable and condensed)

in almost every extract except for Chloroform (ChEL), Solvent Ether (SEEL) and Petroleum Ether Extract (PEEL). No traces of alkaloids were detected in any of the extract. Chloroform (ChEL), Solvent Ether (SEEL) and Petroleum Ether Extract of leaf (PEEL) shows dearth of different glycosides, while other extract showed existence of cardiac, saponin, flavanoidal and coumarin glycosides. Being nonpolar in nature ChEL, SEEL and PEEL showed presence of Steroids and terpenes. Some extracts shows presence of organic acids also.

3.3 Screening of different extracts for In-vitro antioxidant activity

In the present study, in vitro antioxidant activity of the different leaf extract of *Memecylon umbellatum* was investigated by DPPH radical scavenging assays. It is probably due to the presence of phytochemicals like polyphenolics, steroids, glycosides and saponins, highly responsible secondary metabolite for antioxidant activities in these species. Methanol leaf extract showed 84.65 ± 0.064 % antioxidant activity which is higher than other extract [Pet. Ether (3.17 ± 0.084), Chloroform (4.76 ± 0.036), Ethyl acetate (77.24 ± 0.059), Acetone (78.83 ± 0.167), n-Butanol (82.01 ± 0.188) Ethanol (49.20 ± 0.068) and Aqueous extract (11.11 ± 0.058)] at 20 µg/ml. Pet. Ether, Chloroform and aqueous leaf extracts showed very feeble antioxidant activity. Antioxidant activity of different extracts was found to be in order as follow,

Methanol > n-butanol > Acetone > Ethyl acetate > Ethanol > Aqueous > Chloroform > Pet. Ether.

IC₅₀ value, a guide for antioxidant value was determined from % antioxidant activity has been shown in figure 1.




Figure 1: Plot of % antioxidant activity versus various concentrations of standard and different extracts.

Methanol leaf extract showed significant ($p<0.05$ Graphpad instat 3) IC_{50} value (11.81 $\mu\text{g/ml}$) compared to standard i.e. ascorbic acid (8.91 $\mu\text{g/ml}$). All other extracts showed higher IC_{50} value, indicate lesser antioxidant activity than standard and methanol leaf

extract. Pet. Ether and chloroform leaf extract showed higher IC_{50} value, 315.45 and 210.08 $\mu\text{g/ml}$ respectively. IC_{50} value of different extract has been shown in figure 2.




Figure 2: Graph showing IC_{50} value of different extracts of *Memecylon umbellatum* leaf.

4 CONCLUSIONS

From the findings of this study, it can be concluded that *Memecylon umbellatum* leaf extracts, emerging as promising natural herbal sources of antioxidants and can be used in nutritional or pharmaceutical fields for the prevention of free radical-mediated perilous diseases (oxidative stresses). However, in-vivo assays are essential to characterise it as biological antioxidants. In addition to this, flavonoids, mainly responsible for antioxidant activity need to be investigated in details.

ACKNOWLEDGEMENT

Authors are thankful to Dr. S. R. Yadav, Department of Botany, Shivaji University, Kolhapur, India (M.S.) for their valuable guidance.

CONFLICT OF INTEREST:

We declare that we have no conflict of interest.

REFERENCES

1. Halliwell B, Gutteridge JMC, Free radical in biology and medicine, 3rd ed, Oxford University Press: London; 1998.
2. Temple NJ, Antioxidants and disease: More questions than answers, *Nutr Res*, 2000, 20, 449-59.
3. Farber JL, Mechanisms of cell injury by activated oxygen species, *Environ health perspect* 1994, 102, 17-24.
4. Liu RH, Supplement quick fix fails to deliver, *Food Technol Int*, 2002, 1, 71-2.
5. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev*, 2010, 4(8), 118-26.
6. Dhalla NS, Temsah RM, Netticadan T, Role of oxidative stress in cardiovascular diseases, *J Hypertens*, 2000, 18(6), 655-73.
7. Jenner P, Oxidative stress in Parkinson's disease, *Ann Neurol*, 2003, 53(3), S26-36.
8. Church SL, Grant JW, Ridnourt LA, Oberleyt LW, Swansont PE, Meltzer PS, Increased manganese superoxide dismutase expression suppresses the malignant phenotype of human melanoma cells, *Proc Natl Acad Sci*, 1993, 90, 3113-17.
9. Praticò D, Clark CM, Liun F, Rokach J, Lee VY, Trojanowski JQ, Increase of brain oxidative stress in mild cognitive impairment: a possible predictor of Alzheimer disease, *Arch Neurol*, 2002, 59(6), 972-6.
10. Uttara B, Singh AV, Zamboni P, Mahajan RT, Oxidative Stress and Neurodegenerative Diseases: A Review of Upstream and Downstream Antioxidant Therapeutic Options, *Curr Neuropharmacol*, 2009, 7(1), 65-74.
11. Perry G, Cash AD, Smith MA, Alzheimer Disease and Oxidative Stress, *J Biomed Biotechnol*, 2002, 2(3), 120-23.
12. Seril DN, Liao J, Yang GY, Yang CS, Oxidative stress and ulcerative colitis-associated carcinogenesis: studies in humans and animal models. *Carcinogenesis*, 2003, 24(3), 353-62.
13. Romano AD, Serviddio G, de Matthaeis A, Bellanti F, Vendemiale G, Oxidative stress and aging. *J Nephrol*, 2010, 23(15), S29-36.
14. Yang H, Jin X, Kei Lam CW, Yan SK, Oxidative stress and diabetes mellitus, *Clin Chem Lab Med*, 2011, 49(11), 1773-82.
15. Fibach E, Rachmilewitz E, The role of oxidative stress in hemolytic anemia, *Curr Mol Med*, 2008, 8(7), 609-19.
16. Harrison D, Griendling KK, Landmesser U, Hornig B, Drexler H, Role of oxidative stress in atherosclerosis, *Am J Cardiol* 2003, 6,91(3A), 7A-11A.
17. Cho YS, Moon HB, The Role of Oxidative Stress in the Pathogenesis of Asthma, *Allergy Asthma Immunol Res*, 2010, 2(3), 183-87.
18. Farrukh A, Iqbal A, Zafar M, Antioxidant and Free Radical Scavenging Properties of Twelve Traditionally Used Indian Medicinal Plants, *Turk J Biol*, 2006, 30, 177-83.
19. Prior RL, Fruit and vegetables in the prevention of cellular oxidative damage, *Am J Clin Nutr*, 2003, 78, 570S-78S.
20. Kaur C, Kapoor HC, Anti-oxidant activity and total phenolics content of some Asian vegetables, *Int J Food Sci Technol*, 2002, 37, 153-61.
21. Kirtikar KR, Basu BD, Indian Medicinal Plants, 2nd ed, Popular Prakashan: New Delhi, 1996, 1063-67.
22. Nadkarni KM, Indian MateriaMedica, 1st ed. Bombay Prakashan: Bombay, 1982.
23. Padmavathy J, Raju D, Sai Saraswathi V, Kayalvizhi M, Saravanan D, Pharmacognostic Parameters for the Evaluation of the Leaves and Young Stem of *Memecylon umbellatum* Burm.f, *Int J PharmTech Research*, 2010, 2(3), 2001-6.
24. Killedar SG, More HN, Screening of Antimicrobial Potential and Phytoconstituents for Different Extracts of *Memecylon umbellatum* Burm Inflorescences, *Asian J Pharm Res*, 2011, 1(4), 114-18.
25. Killedar SG, More HN, Antimicrobial potential and phytochemical screening of different bark extracts of *Memecylon umbellatum* Burm, *Int J Appl Microb Sci*, 2012, 2(3), 50-8.
26. Killedar SG, More HN, Estimation of Tannins in different parts of *Memecylon Umbellatum* Burm, *J Pharm Research* 2010, 3(3), 554-6.
27. Killedar SG, More HN, Seasonal Variation of Tannin Content in Different Parts of *Memecylon umbellatum* Burm, *Research J Pharm and Tech*, 2011, 4(1), 78-81.
28. Killedar SG, More HN, Estimation of Sugars and minerals in healthy and infected parts of *Memecylon umbellatum* Burm, *Int J research in ayurveda & pharm*, 2010, 1(2), 582-85.
29. Killedar SG, More HN, Shah G, Gaikwad S, Phytochemical screening and in-vitro antioxidant Activity of *Memecylon umbellatum* root extracts, *World J pharm and pharm sci*, 2013, 2(6), 5988-96.
30. Sastri BN, The Wealth of India, Publication and Information Directorate, CSIR Hillside: New Delhi, India, 1962, 336.
31. Dhar ML, Dhar MM, Dhawan BN, Mehrotra BN, Ray C, Screening of Indian plants for biological activity: Part 1. *Indian J Exp Biol*, 1968, 6(4), 232-47.
32. Amalraj T, Ignacimuthu S, Evaluation of Hypoglycemic effect of *Memecylon umbellatum* in normal and alloxan diabetic mice, *J Ethanopharmacol*, 1998, 62, 247-50.
33. Joshi H, Hypoglycemic effect of *Memecylon umbellatum* root, *Phcog Mag*, 2008, 15, 1-4.
34. Puratchikody, Naglakshmi G, Wound Healing Activity of ethanolic extract of *Memecylon umbellatum* burm leaves, *J Plant Sci*, 2007, 2, 179-86.
35. Ram PR, Mehrotra BN, Compendium of Indian Medicinal Plants, (Drug Research Preparative: A CDRI Series). Central Drug Research Institute, Lucknow and Publications and Information Directorate: New Delhi, 1993, 453.
36. Asolkar LV, Kakkar KK and Chakre OJ, Glossary of Indian Medicinal Plants with active principal, part-I. Publications and Information Directorate, Council of Scientific and Industrial Research Publ: New Delhi; 1956, 165.
37. Peach EK, Tracy MV, Modern Methods of Plant Analysis. Springer Verlag Publication: Vol-III, 1915 , 332-50.
38. Harborne JB, A guide to Modern Techniques of Plant Analysis, Chapman and Hall: London, 1973.
39. British Pharmacopoeia, Appendix XID A153 Vol. II HMSO: London,1993.
40. Nikhat F, Satynarayana D, Subhramanyam EVS. Isolation, Charactrisation and Screening of Antioxidant Activity of the Roots of *Syzygiumcumini* (L) Skeel, *Asian J Research Chem* 2009, 2(2), 218-21.
41. Erasto P, Mbawando ZH, Antioxidant activity and HPTLC profile of *Lagenaria siceraria* fruits. *Tanzania J Health Research*, 2009, 11(2), 79-83.
42. Rice-Evans CA, Miller NJ, Paganga G, Structure-antioxidant activity relationships of flavonoids and phenolic acids, *Free Radical Biol Med*, 1996, 20, 933-56.
43. Gao XL, Bjork V, Trajkovski, Uggla M, Evaluation of antioxidant activities of rosehip ethanol extracts in different test systems, *J Sci Food Agr*, 2000, 80, 2021-27.
44. Mariani C, Flavonoid characterization and in vitro antioxidant activity of *Aconitum anthora* L, (Ranunculaceae) *Phytochemistry*, 2008, 69, 1220-26.
45. Md. Nur Alam, Bristi NJ, Md. Rafiquzzaman. Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharm J*, 2013, 21, 143-52.
46. Sochor J, Ryvolova M, Krystofova O, Salas P, Hubalek J, Adam V, Fully Automated Spectrometric Protocols for Determination of Antioxidant Activity: Advantages and Disadvantages, *Molecules* 2010, 15, 8618-40.
47. Thambiraj J, Paulsamy S, In vitro antioxidant potential of methanol extract of the medicinal plant, *Acacia caesia* (L.) Willd, *Asian Pacific J Tropical Biomed*, 2012, S732-S736.
48. Battu GR, Kumar BM, *In-Vitro Antioxidant Activity of Leaf Extract of Aerva Lanata Linn*, *Int J Pharm Sci*, 2012, 2(4), 74-78.